

Sevinc Can Sandikci

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/441158/publications.pdf>

Version: 2024-02-01

18
papers

141
citations

1307594

7
h-index

1281871

11
g-index

18
all docs

18
docs citations

18
times ranked

244
citing authors

#	ARTICLE	IF	CITATIONS
1	Can the Thiol/Disulfide Imbalance Be a Predictor of Colchicine Resistance in Familial Mediterranean Fever?. <i>Journal of Korean Medical Science</i> , 2017, 32, 1588.	2.5	19
2	An evaluation of depression, anxiety and fatigue in patients with Behçet's disease. <i>International Journal of Rheumatic Diseases</i> , 2019, 22, 974-979.	1.9	19
3	Fatigue in rheumatic diseases. <i>European Journal of Rheumatology</i> , 2015, 2, 109-113.	0.6	19
4	Evaluation of restless legs syndrome and sleep disorders in patients with psoriatic arthritis. <i>Zeitschrift Fur Rheumatologie</i> , 2019, 78, 987-995.	1.0	17
5	Do the European League Against Rheumatism (EULAR) Sjögren's syndrome outcome measures correlate with impaired quality of life, fatigue, anxiety and depression in primary Sjögren's syndrome?. <i>Archives of Medical Science</i> , 2018, 14, 830-837.	0.9	13
6	Serum calprotectin and ischemia modified albumin levels as markers of disease activity in Behçet's disease. <i>Postepy Dermatologii i Alergologii</i> , 2018, 35, 609-613.	0.9	11
7	The relationship of serum VEGF and sVEGFR-1 levels with vascular involvement in patients with Behçet's disease. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2018, 78, 443-449.	1.2	9
8	Thiol/Disulfide homeostasis in patients with rheumatoid arthritis. <i>Romanian Journal of Internal Medicine = Revue Roumaine De Medecine Interne</i> , 2019, 57, 30-36.	0.6	7
9	Serum neopterin and ischemia modified albumin levels are associated with the disease activity of adult immunoglobulin A vasculitis (Henoch-Schönlein purpura). <i>International Journal of Rheumatic Diseases</i> , 2019, 22, 1920-1925.	1.9	6
10	Short report: correlates of functional disability with disease activity in elderly patients with rheumatoid arthritis. <i>Psychology, Health and Medicine</i> , 2018, 23, 668-673.	2.4	4
11	Determination of serum vascular endothelial growth factor (VEGF) and VEGF receptor levels with VEGF gene polymorphisms in patients with Behçet's uveitis. <i>Advances in Clinical and Experimental Medicine</i> , 2022, 31, 231-240.	1.4	4
12	Is there a relationship between serum omentin level and acute phase response in patients with familial Mediterranean fever?. <i>Clinical Rheumatology</i> , 2021, 40, 669-674.	2.2	3
13	Follow-Up of Treatment Response With Dynamic Doppler Ultrasound in Raynaud Phenomenon. <i>American Journal of Roentgenology</i> , 2017, 209, W388-W394.	2.2	2
14	The relationship between bristol rheumatoid arthritis fatigue scales and disease activity of patients with rheumatoid arthritis. <i>Clinical Rheumatology</i> , 2018, 37, 2927-2932.	2.2	2
15	Evaluation of red blood cell distribution width in patients with psoriatic arthritis. <i>Egyptian Rheumatologist</i> , 2020, 42, 309-312.	1.0	2
16	An investigation of thiol/disulfide homeostasis in patients with Behçet's disease. <i>Archives of Medical Science</i> , 2020, 16, 1353-1359.	0.9	2
17	Lipoprotein-associated phospholipase A2 and carotid intima-media thickness in primary Sjögren syndrome. <i>Archives of Rheumatology</i> , 0, , .	0.9	2
18	AB0605â€¦SERUM NEOPTERIN AND ISCHEMIA MODIFIED ALBUMIN LEVELS ARE ASSOCIATED WITH THE DISEASE ACTIVITY OF ADULT IMMUNOGLOBULIN A VASCULITIS (HENOCH-SCHONLEIN PURPURA)., 2019, , .		0