
Koichi Amano

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4420304/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Integrated safety analysis of filgotinib treatment for rheumatoid arthritis in patients from Japan over a median of 1.5 years. Modern Rheumatology, 2023, 33, 64-72.	1.8	2
2	Efficacy and safety of filgotinib in Japanese patients with refractory rheumatoid arthritis: Subgroup analyses of a global phase 3 study (FINCH 2). Modern Rheumatology, 2022, 32, 59-67.	1.8	6
3	Nation-wide survey of the treatment trend of microscopic polyangiitis and granulomatosis with polyangiitis in Japan using the Japanese Ministry of Health, Labour and Welfare Database. Modern Rheumatology, 2022, 32, 915-922.	1.8	7
4	Novel susceptibility loci for steroid-associated osteonecrosis of the femoral head in systemic lupus erythematosus. Human Molecular Genetics, 2022, 31, 1082-1095.	2.9	1
5	Efficacy and safety of filgotinib in combination with methotrexate in Japanese patients with active rheumatoid arthritis who have an inadequate response to methotrexate: Subpopulation analyses of 24-week data of a global phase 3 study (FINCH 1). Modern Rheumatology, 2022, 32, 263-272.	1.8	10
6	Efficacy and safety of filgotinib alone and in combination with methotrexate in Japanese patients with active rheumatoid arthritis and limited or no prior exposure to methotrexate: Subpopulation analyses of 24-week data of a global phase 3 study (FINCH 3). Modern Rheumatology, 2022, 32, 273-283.	1.8	3
7	Concurrent Takayasu Arteritis and Vascular Ehlers–Danlos Syndrome: A Case Report. Frontiers in Cardiovascular Medicine, 2022, 9, 805505.	2.4	1
8	Selection of treatment regimens based on shared decision-making in patients with rheumatoid arthritis on remission in the FREE-J study. Rheumatology, 2022, 61, 4273-4285.	1.9	3
9	Clinical impact of urinary CD11b and CD163 on the renal outcomes of anti-neutrophil cytoplasmic antibody-associated glomerulonephritis. Nephrology Dialysis Transplantation, 2021, 36, 1452-1463.	0.7	5
10	Meta-analysis of 208370 East Asians identifies 113 susceptibility loci for systemic lupus erythematosus. Annals of the Rheumatic Diseases, 2021, 80, 632-640.	0.9	103
11	Vascular endothelial growth factor (VEGF)-A and VEGF-A ₁₆₅ b are associated with time to remission of granulomatosis with polyangiitis in a nationwide Japanese prospective cohort study. Annals of Clinical Biochemistry, 2021, 58, 86-94.	1.6	1
12	Kidney GATA3+ regulatory T cells play roles in the convalescence stage after antibody-mediated renal injury. Cellular and Molecular Immunology, 2021, 18, 1249-1261.	10.5	31
13	Rationale of concomitant cyclophosphamide for remission-induction in patients with antineutrophil cytoplasmic antibody-associated vasculitis: A propensity score-matched analysis of two nationwide prospective cohort studies. Modern Rheumatology, 2021, 31, 205-213.	1.8	3
14	Usefulness of tissue inhibitor of metalloproteinase 1 as a predictor of sustained remission in patients with antineutrophil cytoplasmic antibody-associated vasculitis. Arthritis Research and Therapy, 2021, 23, 91.	3.5	5
15	Exploratory classification of clinical phenotypes in Japanese patients with antineutrophil cytoplasmic antibody-associated vasculitis using cluster analysis. Scientific Reports, 2021, 11, 5223.	3.3	5
16	Effect of Reduced-Dose vs High-Dose Glucocorticoids Added to Rituximab on Remission Induction in ANCA-Associated Vasculitis. JAMA - Journal of the American Medical Association, 2021, 325, 2178.	7.4	103
17	Impact of the HLA-DRB1 shared epitope on responses to treatment with tofacitinib or abatacept in patients with rheumatoid arthritis. Arthritis Research and Therapy, 2021, 23, 228.	3.5	9
18	Potential Triggers for Thrombocytopenia and/or Hemorrhage by the BNT162b2 Vaccine, Pfizer-BioNTech. Frontiers in Medicine, 2021, 8, 751598.	2.6	9

#	Article	IF	CITATIONS
19	Safety and tolerability of sifalimumab, an anti-interferon-α monoclonal antibody, in Japanese patients with systemic lupus erythematosus: A multicenter, phase 2, open-label study. Modern Rheumatology, 2020, 30, 93-100.	1.8	23
20	Predictive value of serum amyloid a levels for requirement of concomitant methotrexate in tocilizumab initiation: A <i>post hoc</i> analysis of the SURPRISE study. Modern Rheumatology, 2020, 30, 442-449.	1.8	2
21	Tocilizumab monotherapy for large vessel vasculitis: results of 104-week treatment of a prospective, single-centre, open study. Rheumatology, 2020, 59, 1617-1621.	1.9	25
22	Sustained discontinuation of infliximab with a raising-dose strategy after obtaining remission in patients with rheumatoid arthritis: the RRRR study, a randomised controlled trial. Annals of the Rheumatic Diseases, 2020, 79, 94-102.	0.9	13
23	Treatment-related damage in elderly-onset ANCA-associated vasculitis: safety outcome analysis of two nationwide prospective cohort studies. Arthritis Research and Therapy, 2020, 22, 236.	3.5	14
24	A case of polyangiitis overlap syndrome of giant cell arteritis and granulomatosis with polyangiitis successfully treated with rituximab. Modern Rheumatology Case Reports, 2020, 5, 1-5.	0.7	3
25	Identification of molecules associated with response to abatacept in patients with rheumatoid arthritis. Arthritis Research and Therapy, 2020, 22, 46.	3.5	19
26	JCS 2017 Guideline on Management of Vasculitis Syndrome ― Digest Version ―. Circulation Journal, 2020, 84, 299-359.	1.6	59
27	The clinical impact of absolute lymphocyte count in peripheral blood among patients with methotrexate - associated lymphoproliferative disorders. Journal of Clinical and Experimental Hematopathology: JCEH, 2020, 60, 41-50.	0.8	9
28	2017 Clinical practice guidelines of the Japan Research Committee of the Ministry of Health, Labour, and Welfare for Intractable Vasculitis for the management of ANCA-associated vasculitis. Modern Rheumatology, 2019, 29, 20-30.	1.8	47
29	Tocilizumab monotherapy for polymyalgia rheumatica: A prospective, singleâ€center, openâ€label study. International Journal of Rheumatic Diseases, 2019, 22, 2151-2157.	1.9	21
30	Clinicopathological features of clinical methotrexate-related lymphoproliferative disorders. Leukemia and Lymphoma, 2019, 60, 2508-2515.	1.3	8
31	FRI0099â€PREDICTIVE FACTORS FOR REMISSION ACHIEVEMENT BY TOCILIZUMAB MONOTHERAPY IN PATIENT WITH RHEUMATOID ARTHRITIS AFTER INADEQUATE RESPONSE TO METHOTREXATE: A POST HOC ANALYSIS OF THE SURPRISE STUDY. , 2019, , .	S	0
32	Histopathological classification of anti-neutrophil cytoplasmic antibody-associated glomerulonephritis in a nationwide Japanese prospective 2-year follow-up cohort study. Clinical and Experimental Nephrology, 2019, 23, 387-394.	1.6	9
33	Efficacy and tolerability of six-week extended dosing interval with tocilizumab therapy in a prospective cohort as remission maintenance in patients with rheumatoid arthritis. Modern Rheumatology, 2018, 28, 444-451.	1.8	6
34	Clinicopathologic investigation of methotrexate-induced lymphoproliferative disorders, with a focus on regression. Leukemia and Lymphoma, 2018, 59, 1143-1152.	1.3	49
35	Efficacy and safety of baricitinib in Japanese patients with rheumatoid arthritis: Subgroup analyses of four multinational phase 3 randomized trials. Modern Rheumatology, 2018, 28, 583-591.	1.8	36
36	Association Between Reappearance of Myeloperoxidase–Antineutrophil Cytoplasmic Antibody and Relapse in Antineutrophil Cytoplasmic Antibody–Associated Vasculitis. Arthritis and Rheumatology, 2018, 70, 1626-1633.	5.6	34

#	Article	IF	CITATIONS
37	Tocilizumab discontinuation after attaining remission in patients with rheumatoid arthritis who were treated with tocilizumab alone or in combination with methotrexate: results from a prospective randomised controlled study (the second year of the SURPRISE study). Annals of the Rheumatic Diseases, 2018, 77, 1268-1275.	0.9	43
38	Successful treatment with tocilizumab monotherapy for Takayasu arteritis developing during infliximab therapy in a patient with ulcerative colitis. Modern Rheumatology Case Reports, 2018, 2, 174-176.	0.7	4
39	Restoration of Decreased T Helper 1 and CD8+ T Cell Subsets Is Associated With Regression of Lymphoproliferative Disorders Developed During Methotrexate Treatment. Frontiers in Immunology, 2018, 9, 621.	4.8	21
40	Era of steroid sparing in the management of immune-mediated inflammatory diseases. Immunological Medicine, 2018, 41, 6-11.	2.6	13
41	Discontinuation of tofacitinib after achieving low disease activity in patients with rheumatoid arthritis: a multicentre, observational study. Rheumatology, 2017, 56, 1293-1301.	1.9	19
42	Prediction of response to remission induction therapy by gene expression profiling of peripheral blood in Japanese patients with microscopic polyangiitis. Arthritis Research and Therapy, 2017, 19, 117.	3.5	10
43	Achieving simplified disease activity index remission in patients with active rheumatoid arthritis is associated with subsequent good functional and structural outcomes in a real-world clinical setting under a treat-to-target strategy. Modern Rheumatology, 2017, 27, 811-819.	1.8	10
44	Current clinical evidence of tocilizumab for the treatment of ANCA-associated vasculitis: a prospective case series for microscopic polyangiitis in a combination with corticosteroids and literature review. Clinical Rheumatology, 2017, 36, 2383-2392.	2.2	36
45	Genome-wide Association Study of Idiopathic Osteonecrosis of the Femoral Head. Scientific Reports, 2017, 7, 15035.	3.3	23
46	Clinical characteristics of and risk factors for serious infection in Japanese patients within six months of remission induction therapy for antineutrophil cytoplasmic antibody-associated vasculitis registered in a nationwide, prospective, inception cohort study. Modern Rheumatology, 2017, 27, 646-651.	1.8	25
47	Simplified Disease Activity Index remission at month 6 is an independent predictor of functional and structural remissions at month 12 during abatacept treatment in patients with rheumatoid arthritis: A multi-center, prospective cohort study in Japan. Modern Rheumatology, 2017, 27, 787-794.	1.8	5
48	Low-dose glucocorticoids plus rituximab versus high-dose glucocorticoids plus rituximab for remission induction in ANCA-associated vasculitis (LoVAS): protocol for a multicentre, open-label, randomised controlled trial. BMJ Open, 2017, 7, e018748.	1.9	16
49	Targeted proteomics reveals promising biomarkers of disease activity and organ involvement in antineutrophil cytoplasmic antibody-associated vasculitis. Arthritis Research and Therapy, 2017, 19, 218.	3.5	40
50	Discussion on Management of Rheumatic Diseases for Elderly Patients. The Journal of the Japanese Society of Internal Medicine, 2017, 106, 2170-2181.	0.0	0
51	Identification of baseline gene expression signatures predicting therapeutic responses to three biologic agents in rheumatoid arthritis: a retrospective observational study. Arthritis Research and Therapy, 2016, 18, 159.	3.5	38
52	Comparison of adding tocilizumab to methotrexate with switching to tocilizumab in patients with rheumatoid arthritis with inadequate response to methotrexate: 52-week results from a prospective, randomised, controlled study (SURPRISE study). Annals of the Rheumatic Diseases, 2016, 75, 1917-1923.	0.9	81
53	Comparison of severity classification in Japanese patients with antineutrophil cytoplasmic antibody-associated vasculitis in a nationwide, prospective, inception cohort study. Modern Rheumatology, 2016, 26, 730-737.	1.8	39
54	Corticosteroid-free treatment of tocilizumab monotherapy for microscopic polyangiitis: a single-arm, single-center, clinical trial. Modern Rheumatology, 2016, 26, 900-907.	1.8	18

#	Article	IF	CITATIONS
55	V. Giant Cell Arteritis. The Journal of the Japanese Society of Internal Medicine, 2015, 104, 2139-2142.	0.0	1
56	Different responses to treatment across classified diseases and severities in Japanese patients with microscopic polyangiitis and granulomatosis with polyangiitis: a nationwide prospective inception cohort study. Arthritis Research and Therapy, 2015, 17, 305.	3.5	41
57	Head-to-head comparison of the safety of tocilizumab and tumor necrosis factor inhibitors in rheumatoid arthritis patients (RA) in clinical practice: results from the registry of Japanese RA patients on biologics for long-term safety (REAL) registry. Arthritis Research and Therapy, 2015, 17, 74.	3.5	53
58	Long-term safety and efficacy of treatment with subcutaneous abatacept in Japanese patients with rheumatoid arthritis who are methotrexate inadequate responders. Modern Rheumatology, 2015, 25, 665-671.	1.8	11
59	Infliximab and etanercept have distinct actions but similar effects on cytokine profiles in rheumatoid arthritis. Cytokine, 2015, 75, 222-227.	3.2	16
60	Longterm Safety and Efficacy of Subcutaneous Tocilizumab Monotherapy: Results from the 2-year Open-label Extension of the MUSASHI Study. Journal of Rheumatology, 2015, 42, 799-809.	2.0	39
61	Biologic-free remission of established rheumatoid arthritis after discontinuation of abatacept: a prospective, multicentre, observational study in Japan. Rheumatology, 2015, 54, 683-691.	1.9	39
62	Drug free REmission/low disease activity after cessation of tocilizumab (Actemra) Monotherapy (DREAM) study. Modern Rheumatology, 2014, 24, 17-25.	1.8	105
63	Retreatment efficacy and safety of tocilizumab in patients with rheumatoid arthritis in recurrence (RESTORE) study. Modern Rheumatology, 2013, , 1.	1.8	1
64	Drug free REmission/low disease activity after cessation of tocilizumab (Actemra) Monotherapy (DREAM) study. Modern Rheumatology, 2013, , 1.	1.8	5
65	No increased mortality in patients with rheumatoid arthritis treated with biologics: results from the biologics register of six rheumatology institutes in Japan. Modern Rheumatology, 2013, 23, 945-952.	1.8	20
66	Effectiveness and safety of adalimumab in Japanese patients with rheumatoid arthritis: retrospective analyses of data collected during the first year of adalimumab treatment in routine clinical practice (HARMONY study). Modern Rheumatology, 2012, 22, 327-338.	1.8	38
67	Structural damages disturb functional improvement in patients with rheumatoid arthritis treated with etanercept. Modern Rheumatology, 2012, 22, 186-194.	1.8	6
68	Differences between the Health Assessment Questionnaire Disability Index (HAQ-DI) and the modified HAQ (mHAQ) score before and after infliximab treatment in patients with rheumatoid arthritis. Modern Rheumatology, 2010, 20, 337-342.	1.8	9
69	Japan College of Rheumatology 2009 guidelines for the use of tocilizumab, a humanized anti-interleukin-6 receptor monoclonal antibody, in rheumatoid arthritis. Modern Rheumatology, 2009, 19, 351-357.	1.8	75
70	Molecular effects to human regulatory T cells by treatment with T cell activation inhibitor and TNF blockade. FASEB Journal, 2008, 22, 1073.10.	0.5	0
71	Prospective study of low-dose cyclosporine A in patients with refractory lupus nephritis. Modern Rheumatology, 2007, 17, 92-97.	1.8	26
72	Clinical significance of elevated serum levels of matrix metalloproteinaseâ€3 and Câ€reactive protein in patients with rheumatoid arthritis. APLAR Journal of Rheumatology, 2007, 10, 295-299.	0.2	3

#	Article	IF	CITATIONS
73	Development and validation of handy rheumatoid activity score with 38 joints (HRAS38) in rheumatoid arthritis patients receiving infliximab. Modern Rheumatology, 2006, 16, 381-388.	1.8	16
74	Impact of TNF inhibitors on rheumatoid arthritis. Inflammation and Regeneration, 2006, 26, 148-159.	3.7	3
75	Amebiasis in Acquired Immunodeficiency Syndrome. Internal Medicine, 2001, 40, 563-564.	0.7	8
76	Candida pneumonia in a case of polymyositis complicated with interstitial pneumonitis. Japanese Journal of Rheumatology, 1999, 9, 397-402.	0.0	0
77	Candidapneumonia in a case of polymyositis complicated with interstitial pneumonitis. Japanese Journal of Rheumatology, 1999, 9, 397-402.	0.0	0
78	Nosocomial Pneumonia Likely Caused by Stenotrophomonas Maltophilia in Two Patients with Polymyositis Internal Medicine, 1999, 38, 910-916.	0.7	4
79	Integrin VLA-5 Negative Primary Plasma Cell Leukemia Internal Medicine, 1993, 32, 565-568.	0.7	5
80	Clinical studies of renal disorders in patients with rheumatoid arthritis Japanese Journal of Clinical Immunology, 1992, 15, 135-141.	0.0	0