## Noble Selasi Gati

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4420824/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Variation among Staphylococcus aureus membrane vesicle proteomes affects cytotoxicity of host cells. Microbial Pathogenesis, 2016, 93, 185-193.	2.9	76
2	Differences in Biofilm Mass, Expression of Biofilm-Associated Genes, and Resistance to Desiccation between Epidemic and Sporadic Clones of Carbapenem-Resistant Acinetobacter baumannii Sequence Type 191. PLoS ONE, 2016, 11, e0162576.	2.5	36
3	Origin and Evolution of Hybrid Shiga Toxin-Producing and Uropathogenic Escherichia coli Strains of Sequence Type 141. Journal of Clinical Microbiology, 2019, 58, .	3.9	31
4	Genetic basis of antimicrobial resistance and clonal dynamics of carbapenem-resistant Acinetobacter baumannii sequence type 191 in a Korean hospital. Infection, Genetics and Evolution, 2015, 36, 1-7.	2.3	24
5	<i>Stenotrophomonas maltophilia</i> outer membrane vesicles elicit a potent inflammatory response <i>in vitro</i> and <i>in vivo</i> . Pathogens and Disease, 2016, 74, ftw104.	2.0	23
6	Clostridium difficile -derived membrane vesicles induce the expression of pro-inflammatory cytokine genes and cytotoxicity in colonic epithelial cells inÂvitro. Microbial Pathogenesis, 2017, 107, 6-11.	2.9	21
7	Molecular epidemiology and antimicrobial susceptibility of Clostridium difficile isolates from two Korean hospitals. PLoS ONE, 2017, 12, e0174716.	2.5	17
8	Comparative phenotypic characterization of hybrid Shiga toxin-producing / uropathogenic Escherichia coli, canonical uropathogenic and Shiga toxin-producing Escherichia coli. International Journal of Medical Microbiology, 2021, 311, 151533.	3.6	8
9	Antimicrobial activity of novel 4 H -4-oxoquinolizine compounds against extensively drug-resistant Acinetobacter baumannii strains. International Journal of Antimicrobial Agents, 2017, 49, 107-111.	2.5	4