## Takeshi Nagasaka

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4424240/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Serum miR-21 as a Diagnostic and Prognostic Biomarker in Colorectal Cancer. Journal of the National Cancer Institute, 2013, 105, 849-859.	3.0	425
2	Fecal MicroRNAs as Novel Biomarkers for Colon Cancer Screening. Cancer Epidemiology Biomarkers and Prevention, 2010, 19, 1766-1774.	1.1	310
3	The CpG Island Methylator Phenotype and Chromosomal Instability Are Inversely Correlated in Sporadic Colorectal Cancer. Gastroenterology, 2007, 132, 127-138.	0.6	264
4	Colorectal Cancer With Mutation in BRAF, KRAS, and Wild-Type With Respect to Both Oncogenes Showing Different Patterns of DNA Methylation. Journal of Clinical Oncology, 2004, 22, 4584-4594.	0.8	202
5	Somatic Hypermethylation of <i>MSH2</i> Is a Frequent Event in Lynch Syndrome Colorectal Cancers. Cancer Research, 2010, 70, 3098-3108.	0.4	167
6	An Optimized Pentaplex PCR for Detecting DNA Mismatch Repair-Deficient Colorectal Cancers. PLoS ONE, 2010, 5, e9393.	1.1	136
7	Analysis of Fecal DNA Methylation to Detect Gastrointestinal Neoplasia. Journal of the National Cancer Institute, 2009, 101, 1244-1258.	3.0	122
8	Mutations in Both KRAS and BRAF May Contribute to the Methylator Phenotype in Colon Cancer. Gastroenterology, 2008, 134, 1950-1960.e1.	0.6	114
9	AZIN1 RNA editing confers cancer stemness and enhances oncogenic potential in colorectal cancer. JCI Insight, 2018, 3, .	2.3	91
10	Methylation pattern of theO6-methylguanine-DNA methyltransferase gene in colon during progressive colorectal tumorigenesis. International Journal of Cancer, 2008, 122, 2429-2436.	2.3	62
11	Poor prognosis of <i>KRAS</i> or <i>BRAF</i> mutant colorectal liver metastasis without microsatellite instability. Journal of Hepato-Biliary-Pancreatic Sciences, 2013, 20, 223-233.	1.4	50
12	Activation of AZIN1 RNA editing is a novel mechanism that promotes invasive potential of cancer-associated fibroblasts in colorectal cancer. Cancer Letters, 2019, 444, 127-135.	3.2	40
13	Randomised phase II trial of mFOLFOX6 plus bevacizumab versus mFOLFOX6 plus cetuximab as first-line treatment for colorectal liver metastasis (ATOM trial). British Journal of Cancer, 2019, 121, 222-229.	2.9	37
14	PD-L1 expression combined with microsatellite instability/CD8+ tumor infiltrating lymphocytes as a useful prognostic biomarker in gastric cancer. Scientific Reports, 2019, 9, 4633.	1.6	37
15	Clinical impact of endometrial cancer stratified by genetic mutational profiles, POLE mutation, and microsatellite instability. PLoS ONE, 2018, 13, e0195655.	1.1	30
16	Molecular diagnosis and therapy for occult peritoneal metastasis in gastric cancer patients. World Journal of Gastroenterology, 2014, 20, 17796-17803.	1.4	28
17	Fluorescence virus-guided capturing system of human colorectal circulating tumour cells for non-invasive companion diagnostics. Gut, 2015, 64, 627-635.	6.1	27
18	The rare BRAF VK600-601E mutation as a possible indicator of poor prognosis in rectal carcinoma – a report of a case. BMC Medical Genetics, 2015, 16, 1.	2.1	24

#	Article	IF	CITATIONS
19	Clinical outcomes of women with ovarian metastases of colorectal cancer treated with oophorectomy with respect to their somatic mutation profiles. Oncotarget, 2018, 9, 16477-16488.	0.8	23
20	Detection of circulating microRNAs with Ago2 complexes to monitor the tumor dynamics of colorectal cancer patients during chemotherapy. International Journal of Cancer, 2019, 144, 2169-2180.	2.3	22
21	Accuracy of four mononucleotide-repeat markers for the identification of DNA mismatch-repair deficiency in solid tumors. Journal of Translational Medicine, 2018, 16, 5.	1.8	21
22	Expansion of CpG methylation in the SFRP2 promoter region during colorectal tumorigenesis. Acta Medica Okayama, 2011, 65, 169-77.	0.1	18
23	Heterogeneity of Epigenetic and Epithelial Mesenchymal Transition Marks in Hepatocellular Carcinoma with Keratin 19 Proficiency. Liver Cancer, 2019, 8, 239-254.	4.2	14
24	Multicenter phase II study of biweekly CAPIRI plus bevacizumab as second-line therapy in patients with metastatic colorectal cancer (JSWOG-C3 study). International Journal of Clinical Oncology, 2019, 24, 1223-1230.	1.0	9
25	Clinical and epigenetic features of colorectal cancer patients with somatic POLE proofreading mutations. Clinical Epigenetics, 2021, 13, 117.	1.8	8
26	Comparison of outcomes between symptomatic and asymptomatic patients with colorectal cancer: a propensity score-matched analysis of surgical invasiveness, medical costs and oncological outcomes. BMJ Open Gastroenterology, 2017, 4, e000146.	1.1	7
27	Protocol of a randomised phase III clinical trial of sequential capecitabine or 5-fluorouracil plus bevacizumab (Cape/5-FU-Bmab) to capecitabine or 5-fluorouracil plus oxaliplatin plus bevacizumab (CapeOX/mFOLFOX6-Bmab) versus combination CapeOX/mFOLFOX6-Bmab in advanced colorectal cancer: the C-cubed (C <sup>3</sup> ) study. BMI Open, 2016, 6, e011454.	0.8	6
28	BRAF V600E mutation is a predictive indicator of upfront chemotherapy for stageÃ <sup>-</sup> ¿½IV colorectal cancer. Oncology Letters, 2017, 15, 2195-2201.	0.8	6
29	Upregulation of microRNA‑31 is associated with poor prognosis in patients with advanced colorectal cancer. Oncology Letters, 2020, 19, 2685-2694.	0.8	6
30	Expansion of epigenetic alterations in EFEMP1 promoter predicts malignant formation in pancreatobiliary intraductal papillary mucinous neoplasms. Journal of Cancer Research and Clinical Oncology, 2016, 142, 1557-1569.	1.2	5
31	Detection of fecal DNA methylation for colorectal neoplasia: does it lead to an optimal screening test?. Acta Medica Okayama, 2006, 60, 249-56.	0.1	5
32	A multicenter single‑arm Phase II clinical trial of second‑line FOLFIRI plus panitumumab after first‑line treatment with FOLFOX plus panitumumab for initial RAS wild‑type colorectal cancer with evaluation of circulating tumor DNA: A protocol study. Oncology Letters, 2018, 17, 1980-1985.	0.8	3
33	Analysis of fecal DNA methylation to detect gastrointestinal neoplasia. Okayama Igakkai Zasshi, 2010, 122, 107-112.	0.0	1
34	Multicenter open-label randomized phase II study of second-line panitumumab and irinotecan with or without fluoropyrimidines in patients with KRAS wild-type metastatic colorectal cancer (PACIFIC) Tj ETQq0 0 0 rg	;BT1/Øverlc	ock1 10 Tf 50

35	Concordance of acquired mutations between metastatic lesions and liquid biopsy in metastatic colorectal cancer. Future Science OA, 2021, 7, FSO757.	0.9	1
36	Technique of vessel-skeletonized parenchyma-sparing hepatectomy for the oncological treatment of bilobar colorectal liver metastases. Langenbeck's Archives of Surgery, 2021, , 1.	0.8	1

Τακέςμι Νασαδάκα

#	Article	IF	CITATIONS
37	Prognostic Nutritional Index as a Predictor of Postoperative Outcome in Patients Aged 85ÂYears or Older After Colorectal Cancer Surgery. Indian Journal of Surgery, 2020, 82, 874-878.	0.2	0
38	Natural history of epithelioid hemangioendothelioma that progressed over 20 years. Pediatric Blood and Cancer, 2020, 67, e28261.	0.8	0
39	Adenocarcinoma in the jejunum 20 years after surgery for familial adenomatous polyposis. Okayama Igakkai Zasshi, 2017, 129, 111-114.	0.0	Ο
40	Analysis of UGT1A's polymorphisms and RAS: RAF mutations based on phase II study of biweekly XELIRI plus bevacizumab as a second-line therapy in patients with metastatic colorectal cancer Journal of Clinical Oncology, 2019, 37, 565-565.	0.8	0
41	Phase 3 trial of sequential versus combination treatment in colorectal cancer: The C-cubed study. European Journal of Cancer, 2022, 169, 166-178.	1.3	0