

MaÃ-ra Pombo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4426392/publications.pdf>

Version: 2024-02-01

13
papers

137
citations

1478505

6
h-index

1199594

12
g-index

13
all docs

13
docs citations

13
times ranked

121
citing authors

#	ARTICLE	IF	CITATIONS
1	Issues to Be Considered in Counting Burrows as a Measure of Atlantic Ghost Crab Populations, an Important Bioindicator of Sandy Beaches. PLoS ONE, 2013, 8, e83792.	2.5	38
2	The burrow resetting method, an easy and effective approach to improve indirect ghost-crab population assessments. Ecological Indicators, 2019, 104, 422-428.	6.3	19
3	Seasonality, Dietary Overlap and the Role of Taxonomic Resolution in the Study of the Diet of Three Congeneric Fishes from a Tropical Bay. PLoS ONE, 2013, 8, e56107.	2.5	17
4	Continuous, video-recording assessment of daily activity cycle of the ghost crab <i>Ocypode quadrata</i> Fabricius, 1787 (Brachyura: Ocypodidae) in southeastern Brazil. Journal of Crustacean Biology, 2018, 38, 133-139.	0.8	12
5	Population biology of <i>Stellifer rastrifer</i> , <i>S. brasiliensis</i> and <i>S. stellifer</i> in Caraguatatuba Bay, northern coast of São Paulo, Brazil. Brazilian Journal of Oceanography, 2012, 60, 271-282.	0.6	9
6	Ghost crab burrows simulation shows differential across-shore persistence. Crustaceana, 2018, 91, 821-830.	0.3	8
7	Population structure and use of space by ghost crabs (Brachyura: Ocypodidae) on an equatorial, macrotidal sandy beach. Estuarine, Coastal and Shelf Science, 2021, 258, 107376.	2.1	8
8	Variation in the body growth parameters of the ghost crab <i>Ocypode quadrata</i> from morphodynamically distinct sandy beaches. Brazilian Journal of Oceanography, 2017, 65, 656-665.	0.6	8
9	Population Dynamics and Diet of the Madamango Sea Catfish <i>Cathorops spixii</i> (Agassiz, 1829) (Siluriformes: Ariidae) in a Tropical Bight in Southeastern Brazil. PLoS ONE, 2013, 8, e81257.	2.5	6
10	The barred grunt <i>Conodon nobilis</i> (Perciformes: Haemulidae) in shallow areas of a tropical bight: spatial and temporal distribution, body growth and diet. Helgoland Marine Research, 2014, .	1.3	5
11	A new species of philichthyid copepod (Crustacea: Cyclopoida) parasitic on <i>Stellifer</i> spp. (Perciformes: Sciaenidae) from southeastern Brazil. Zootaxa, 2015, 3925, 438-44.	0.5	5
12	Assessing the efficiency of low-cost cleaning methods for diatoms from intertidal sediment samples. Journal of Microbiological Methods, 2021, 187, 106255.	1.6	2
13	Novel structure in sciaenid fish skulls indicates continuous production of the cephalic neuromast cupula. Scientific Reports, 2016, 6, 37523.	3.3	0