

# MaÃ-ra Pombo

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4426392/publications.pdf>

Version: 2024-02-01

13  
papers

137  
citations

1478505

6  
h-index

1199594

12  
g-index

13  
all docs

13  
docs citations

13  
times ranked

121  
citing authors

#	ARTICLE	IF	CITATIONS
1	Assessing the efficiency of low-cost cleaning methods for diatoms from intertidal sediment samples. <i>Journal of Microbiological Methods</i> , 2021, 187, 106255.	1.6	2
2	Population structure and use of space by ghost crabs (Brachyura: Ocypodidae) on an equatorial, macrotidal sandy beach. <i>Estuarine, Coastal and Shelf Science</i> , 2021, 258, 107376.	2.1	8
3	The burrow resetting method, an easy and effective approach to improve indirect ghost-crab population assessments. <i>Ecological Indicators</i> , 2019, 104, 422-428.	6.3	19
4	Continuous, video-recording assessment of daily activity cycle of the ghost crab <i>Ocypode quadrata</i> Fabricius, 1787 (Brachyura: Ocypodidae) in southeastern Brazil. <i>Journal of Crustacean Biology</i> , 2018, 38, 133-139.	0.8	12
5	Ghost crab burrows simulation shows differential across-shore persistence. <i>Crustaceana</i> , 2018, 91, 821-830.	0.3	8
6	Variation in the body growth parameters of the ghost crab <i>Ocypode quadrata</i> from morphodynamically distinct sandy beaches. <i>Brazilian Journal of Oceanography</i> , 2017, 65, 656-665.	0.6	8
7	Novel structure in sciaenid fish skulls indicates continuous production of the cephalic neuromast cupula. <i>Scientific Reports</i> , 2016, 6, 37523.	3.3	0
8	A new species of philichthyid copepod (Crustacea: Cyclopoida) parasitic on <i>Stellifer</i> spp. (Perciformes: Sciaenidae) from southeastern Brazil. <i>Zootaxa</i> , 2015, 3925, 438-44.	0.5	5
9	The barred grunt <i>Conodon nobilis</i> (Perciformes: Haemulidae) in shallow areas of a tropical bight: spatial and temporal distribution, body growth and diet. <i>Helgoland Marine Research</i> , 2014, , .	1.3	5
10	Seasonality, Dietary Overlap and the Role of Taxonomic Resolution in the Study of the Diet of Three Congeneric Fishes from a Tropical Bay. <i>PLoS ONE</i> , 2013, 8, e56107.	2.5	17
11	Population Dynamics and Diet of the Madamango Sea Catfish <i>Cathorops spixii</i> (Agassiz, 1829) (Siluriformes: Ariidae) in a Tropical Bight in Southeastern Brazil. <i>PLoS ONE</i> , 2013, 8, e81257.	2.5	6
12	Issues to Be Considered in Counting Burrows as a Measure of Atlantic Ghost Crab Populations, an Important Bioindicator of Sandy Beaches. <i>PLoS ONE</i> , 2013, 8, e83792.	2.5	38
13	Population biology of <i>Stellifer rastrifer</i> , <i>S. brasiliensis</i> and <i>S. stellifer</i> in Caraguatatuba Bay, northern coast of São Paulo, Brazil. <i>Brazilian Journal of Oceanography</i> , 2012, 60, 271-282.	0.6	9