

Herman Mann

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4427493/publications.pdf>

Version: 2024-02-01

101
papers

2,884
citations

279487

23
h-index

174990

52
g-index

116
all docs

116
docs citations

116
times ranked

3808
citing authors

#	ARTICLE	IF	CITATIONS
1	Hsp90 Levels in Idiopathic Inflammatory Myopathies and Their Association With Muscle Involvement and Disease Activity: A Cross-Sectional and Longitudinal Study. <i>Frontiers in Immunology</i> , 2022, 13, 811045.	2.2	2
2	The beneficial effect of csDMARDs co-medication on drug persistence of first-line TNF inhibitor in rheumatoid arthritis patients: data from Czech ATTRA registry. <i>Rheumatology International</i> , 2022, 42, 803-814.	1.5	3
3	The impact of a csDMARD in combination with a TNF inhibitor on drug retention and clinical remission in axial spondyloarthritis. <i>Rheumatology</i> , 2022, 61, 4741-4751.	0.9	6
4	Cardiovascular Risk in Myositis Patients Compared to the General Population: Preliminary Data From a Single-Center Cross-Sectional Study. <i>Frontiers in Medicine</i> , 2022, 9, .	1.2	5
5	The dysregulation of monocyte subpopulations in individuals at risk of developing rheumatoid arthritis. <i>Rheumatology</i> , 2021, 60, 1823-1831.	0.9	4
6	Interleukin-35 in idiopathic inflammatory myopathies. <i>Cytokine</i> , 2021, 137, 155350.	1.4	3
7	Sexual function in patients with idiopathic inflammatory myopathies: a cross-sectional study. <i>Rheumatology</i> , 2021, 60, 5060-5072.	0.9	6
8	Clinical Trials in Myositis: Where Do We Stand?. <i>Current Treatment Options in Rheumatology</i> , 2021, 7, 222-242.	0.6	0
9	The effect of a 24-week training focused on activities of daily living, muscle strengthening, and stability in idiopathic inflammatory myopathies: a monocentric controlled study with follow-up. <i>Arthritis Research and Therapy</i> , 2021, 23, 173.	1.6	8
10	Clusterin is upregulated in serum and muscle tissue in idiopathic inflammatory myopathies and associates with clinical disease activity and cytokine profile. <i>Clinical and Experimental Rheumatology</i> , 2021, 39, 1021-1032.	0.4	1
11	Retention and response rates in 14 261 PsA patients starting TNF inhibitor treatment—results from 12 countries in EuroSpA. <i>Rheumatology</i> , 2020, 59, 1640-1650.	0.9	18
12	Earlier cancer diagnosis after myositis onset is associated with improved long term survival: results from UK, French and Czech cohorts. <i>Rheumatology</i> , 2020, 59, .	0.9	0
13	Alterations in activin A—myostatin—follistatin system associate with disease activity in inflammatory myopathies. <i>Rheumatology</i> , 2020, 59, 2491-2501.	0.9	15
14	Treatment response and drug retention rates in 24 195 biologic-naïve patients with axial spondyloarthritis initiating TNFi treatment: routine care data from 12 registries in the EuroSpA collaboration. <i>Annals of the Rheumatic Diseases</i> , 2019, 78, 1536-1544.	0.5	58
15	Efficacy and Safety of Subcutaneous Secukinumab 150mg with or Without Loading Regimen in Psoriatic Arthritis: Results from the FUTURE 4 Study. <i>Rheumatology and Therapy</i> , 2019, 6, 393-407.	1.1	53
16	Serum visfatin levels in patients with axial spondyloarthritis and their relationship to disease activity and spinal radiographic damage: a cross-sectional study. <i>Rheumatology International</i> , 2019, 39, 1037-1043.	1.5	6
17	FRI0403—HIGH BASELINE PATIENTS COMPARED WITH EVALUATOR'S GLOBAL ASSESSMENT IS ASSOCIATED WITH LOWER RETENTION AND REMISSION RATES OF FIRST TNF INHIBITOR IN AXSPA PATIENTS — DATA FROM THE EUROSPA COLLABORATION. , 2019, , .		0
18	FRI0404—POOLED 6-MONTH TREATMENT OUTCOMES AND DRUG RETENTION RATES IN 1556 PATIENTS WITH AXIAL SPONDYLOARTHRITIS TREATED WITH SECUKINUMAB IN ROUTINE CLINICAL PRACTICE IN 12 EUROPEAN COUNTRIES IN THE EUROSPA RESEARCH COLLABORATION. , 2019, , .		1

#	ARTICLE	IF	CITATIONS
19	AB0752â€¦DRUG RETENTION AND REMISSION RATES IN 14,261 BIOLOGIC-NAÏVE PATIENTS WITH PSORIATIC ARTHRITIS STARTING TNF INHIBITOR TREATMENT IN ROUTINE CARE â€œ RESULTS FROM 12 REGISTRIES IN THE EUROSPA RESEARCH COLLABORATION. , 2019, , .		0
20	AB0687â€¦EFFECTIVENESS OF SPECIALIZED AND INTENSIVE ADL TRAINING IN PATIENTS WITH IDIOPATHIC INFLAMMATORY MYOPATHIES â€œ PRELIMINARY RESULTS OF A ONE-YEAR CONTROLLED STUDY. , 2019, , .		0
21	SAT0391â€¦REMISSION AND DRUG RETENTION RATES OF SECUKINUMAB IN 1549 PATIENTS WITH PSORIATIC ARTHRITIS TREATED IN ROUTINE CARE â€œ POOLED DATA FROM THE OBSERVATIONAL EUROSPA RESEARCH COLLABORATION NETWORK. , 2019, , .		0
22	FRI0107â€¦THE EFFECT OF CO-MEDICATION WITH METHOTREXATE OR OTHER CONVENTIONAL SYNTHETIC DISEASE MODIFYING ANTI-RHEUMATIC DRUGS ON FIRST TUMOR NECROSIS INHIBITOR DRUG SURVIVAL IN PATIENTS WITH RHEUMATOID ARTHRITIS: DATA FROM THE CZECH ATTRA REGISTRY. , 2019, , .		0
23	AB0676â€¦BODY COMPOSITION IN MYOSITIS PATIENTS AND THE ASSOCIATION WITH DISEASE SPECIFIC FEATURES, PHYSICAL ACTIVITY AND PLASMA LEVELS OF INFLAMMATORY CYTOKINES. , 2019, , .		0
24	FRI0378â€¦DOES DRUG EFFECTIVENESS OF 2ND AND 3RD TNF INHIBITORS IN PATIENTS WITH AXIAL SPONDYLOARTHRITIS DEPEND ON THE REASON FROM WITHDRAWAL FROM THE PREVIOUS TREATMENT? â€œ RESULTS FROM THE EUROSPA RESEARCH COLLABORATION. , 2019, , .		3
25	AB0660â€¦SEXUAL HEALTH IMPAIRMENT IN WOMEN WITH IDIOPATHIC INFLAMMATORY MYOPATHIES. , 2019, , .		0
26	FRI0453â€¦DOES DISCORDANCE BETWEEN BASELINE PATIENTâ€™S AND EVALUATORâ€™S GLOBAL ASSESSMENT OF DISEASE ACTIVITY IMPACT RETENTION AND REMISSION RATES OF TNF INHIBITORS IN PATIENTS WITH PSORIATIC ARTHRITIS? DATA FROM THE EUROSPA RESEARCH COLLABORATION. , 2019, , .		0
27	Arthritis in Idiopathic Inflammatory Myopathies. <i>Current Rheumatology Reports</i> , 2019, 21, 70.	2.1	10
28	Serum calprotectin may reflect inflammatory activity in patients with active rheumatoid arthritis despite normal to low C-reactive protein. <i>Clinical Rheumatology</i> , 2018, 37, 2055-2062.	1.0	23
29	Splicing variant of <i>WDFY4</i> augments MDA5 signalling and the risk of clinically amyopathic dermatomyositis. <i>Annals of the Rheumatic Diseases</i> , 2018, 77, 602-611.	0.5	51
30	Abatacept in the treatment of adult dermatomyositis and polymyositis: a randomised, phase IIb treatment delayed-start trial. <i>Annals of the Rheumatic Diseases</i> , 2018, 77, 55-62.	0.5	100
31	CNT06785, a Fully Human Antiinterleukin 17 Monoclonal Antibody, in Patients with Rheumatoid Arthritis with Inadequate Response to Methotrexate: A Randomized, Placebo-controlled, Phase II, Dose-ranging Study. <i>Journal of Rheumatology</i> , 2018, 45, 22-31.	1.0	17
32	The role of imaging in evaluating patients with idiopathic inflammatory myopathies. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 114, 74-81.	0.4	4
33	Immune Array Analysis in Sporadic Inclusion Body Myositis Reveals HLAâ€™DRB1 Amino Acid Heterogeneity Across the Myositis Spectrum. <i>Arthritis and Rheumatology</i> , 2017, 69, 1090-1099.	2.9	41
34	2016 American College of Rheumatology/European League Against Rheumatism Criteria for Minimal, Moderate, and Major Clinical Response in Adult Dermatomyositis and Polymyositis: An International Myositis Assessment and Clinical Studies Group/Paediatric Rheumatology International Trials Organisation Collaborative Initiative. <i>Arthritis and Rheumatology</i> , 2017, 69, 898-910.	2.9	52
35	08.48â€¦Increased body fat but decreased lean body mass and bone mineral density in myositis patients are associated with disease duration, inflammatory status, skeletal muscle involvement and physical activity. , 2017, , .		0
36	THU0113â€¦Serum calprotectin may reflect inflammatory activity in patients with active rheumatoid arthritis despite normal c-reactive protein. , 2017, , .		0

#	ARTICLE	IF	CITATIONS
37	08.49â€¦Efficacy of an intensive 24-week physiotherapy programme in scleroderma patients â€œ preliminary data from a single-centre controlled study. , 2017, , .		0
38	FRIO450â€¦Commonalities and differences in data collection across european spondyloarthritis registries. , 2017, , .		0
39	FRIO399â€¦Efficacy of an intensive 24-week physiotherapy programme in patients with systemic sclerosis - preliminary data from a single-center controlled study. , 2017, , .		0
40	SAT0358â€¦Decreased body fat, lean body mass and bone mineral density in patients with systemic sclerosis are associated with disease activity and physical activity. , 2017, , .		0
41	MRI scoring methods used in evaluation of muscle involvement in patients with idiopathic inflammatory myopathies. Current Opinion in Rheumatology, 2017, 29, 623-631.	2.0	23
42	The relationship between synovitis quantified by an ultrasound 7-joint inflammation score and physical disability in rheumatoid arthritis â€œ a cohort study. Arthritis Research and Therapy, 2017, 19, 5.	1.6	15
43	08.01â€¦Heat shock protein 90 is increased in muscle tissue and plasma in idiopathic inflammatory myopathies. , 2017, , .		0
44	08.47â€¦Decreased body fat, lean body mass and bone mineral density in scleroderma patients are associated with disease activity and physical activity. , 2017, , .		0
45	02.39â€¦Clusterin is increased in idiopathic inflammatory myopathies and is associated with disease activity. , 2017, , .		0
46	THU0058â€¦S100A11 protein is up-regulated in patients with idiopathic inflammatory myopathies and is associated with disease activity. , 2017, , .		0
47	AB0653â€¦Efficacy of an intensive 24-week physiotherapy programme in myositis patients - preliminary data from a single-center controlled study. , 2017, , .		0
48	02.30â€¦High levels of mir-451a differentiate patients with clinically suspect arthralgia from healthy controls. , 2017, , .		0
49	AB1034â€¦Scoring systems of muscle mri in idiopathic inflammatory myopathies. , 2017, , .		0
50	Relationship between serum calprotectin (S100A8/9) and clinical, laboratory and ultrasound parameters of disease activity in rheumatoid arthritis: A large cohort study. PLoS ONE, 2017, 12, e0183420.	1.1	32
51	A tailored approach to reduce dose of anti-TNF drugs may be equally effective, but substantially less costly than standard dosing in patients with ankylosing spondylitis over 1â€¦year: a propensity score-matched cohort study. Annals of the Rheumatic Diseases, 2016, 75, 96-102.	0.5	61
52	AB0952â€¦Serum Visfatin and Resistin, but Not Adiponectin or Leptin, Are Associated with Ultrasound Synovitis in Patients with Rheumatoid Arthritis. Annals of the Rheumatic Diseases, 2016, 75, 1227.1-1227.	0.5	0
53	Systematic protein-protein interaction and pathway analyses in the idiopathic inflammatory myopathies. Arthritis Research and Therapy, 2016, 18, 156.	1.6	4
54	SAT0068â€¦Circulating MIR-126 Is A Marker of Disease Activity in Patients with Early RA but Does Not Differentiate Patients at Risk of Developing RA. Annals of the Rheumatic Diseases, 2016, 75, 689.2-689.	0.5	0

#	ARTICLE	IF	CITATIONS
55	FRI0544â€¦Impact of Education in Musculoskeletal Ultrasonography: Good To Excellent Reliability Results in Modified US7 Scoring System after One Year of Continuing Education since Basic or Intermediate Level of Training. <i>Annals of the Rheumatic Diseases</i> , 2016, 75, 637.1-637.	0.5	0
56	AB0599â€¦Interleukin-35 in Idiopathic Inflammatory Myopathies. <i>Annals of the Rheumatic Diseases</i> , 2016, 75, 1109.2-1109.	0.5	0
57	FRI0532â€¦Association between The 7-Joint Ultrasound Score (US7S) and Physical Function in Rheumatoid Arthritis: Table 1.. <i>Annals of the Rheumatic Diseases</i> , 2016, 75, 632.1-632.	0.5	0
58	MicroRNA-125b: association with disease activity and the treatment response of patients with early rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2016, 18, 124.	1.6	48
59	Dense genotyping of immune-related loci in idiopathic inflammatory myopathies confirms HLA alleles as the strongest genetic risk factor and suggests different genetic background for major clinical subgroups. <i>Annals of the Rheumatic Diseases</i> , 2016, 75, 1558-1566.	0.5	127
60	Serum Calprotectin Discriminates Subclinical Disease Activity from Ultrasound-Defined Remission in Patients with Rheumatoid Arthritis in Clinical Remission. <i>PLoS ONE</i> , 2016, 11, e0165498.	1.1	14
61	Increased visfatin levels are associated with higher disease activity in anti-Jo-1-positive myositis patients. <i>Clinical and Experimental Rheumatology</i> , 2016, 34, 222-9.	0.4	2
62	SAT0436â€¦Abatacept in the Treatment of Adult Dermatomyositis and Polymyositis: Artemis, a Randomized, Treatment Delayed-Start Trial. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 817.3-818.	0.5	3
63	SAT0250â€¦A Tailored Approach to Reduce Dose of Anti-TNF Drugs is Equally Effective and Less Costly Than Standard Dosing in Patients with Ankylosing Spondylitis Over Two Years: A Propensity Score-Matched Cohort Study. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 749.2-749.	0.5	2
64	FRI0592â€¦Serum Calprotectin is Associated with Ultrasound-Determined Active Synovitis in Patients with Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 642.2-642.	0.5	0
65	Serum calprotectin (S100A8/9): an independent predictor of ultrasound synovitis in patients with rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2015, 17, 252.	1.6	60
66	Secukinumab, a human anti-interleukin-17A monoclonal antibody, in patients with psoriatic arthritis (FUTURE 2): a randomised, double-blind, placebo-controlled, phase 3 trial. <i>Lancet</i> , The, 2015, 386, 1137-1146.	6.3	722
67	High levels of metastasis-inducing S100A4 protein and treatment outcome in early rheumatoid arthritis: data from the PERAC cohort. <i>Biomarkers</i> , 2015, 20, 47-51.	0.9	12
68	Increasing incidence of immune-mediated necrotizing myopathy: single-centre experience. <i>Rheumatology</i> , 2015, 54, 2010-2014.	0.9	55
69	Dr. VencovskÃ½, <i>et al</i> reply. <i>Journal of Rheumatology</i> , 2015, 42, 1061.2-1061.	1.0	0
70	Interleukin 35 Synovial Fluid Levels Are Associated with Disease Activity of Rheumatoid Arthritis. <i>PLoS ONE</i> , 2015, 10, e0132674.	1.1	28
71	Decreased Circulating Visfatin Is Associated with Improved Disease Activity in Early Rheumatoid Arthritis: Data from the PERAC Cohort. <i>PLoS ONE</i> , 2014, 9, e103495.	1.1	25
72	THU0168â€¦Monotherapy with Biological Disease-Modifying Anti-Rheumatic Drugs-Data from the Czech Registry Attra.. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 238.2-238.	0.5	0

#	ARTICLE	IF	CITATIONS
73	THU0528â€¦Decrease in Circulating Visfatin Levels is Associated with Disease Activity Improvement in Early Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 366.1-366.	0.5	1
74	A1.48â€¦Enhanced expression of CD11c on non-classical CD16+peripheral blood monocytes in early rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, A20.2-A21.	0.5	1
75	SAT0349â€¦A Tailored Approach to Reduce Dose of Anti-TNF Drugs is Equally Effective, but Substantially Less Costly than Standard Dosing in Patients with Ankylosing Spondylitis over One Year: A Matched Observational Study: Table 1. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 719.1-719.	0.5	1
76	Arthritis in Idiopathic Inflammatory Myopathy: Clinical Features and Autoantibody Associations. <i>Journal of Rheumatology</i> , 2014, 41, 1133-1139.	1.0	23
77	The metastasis promoting protein S100A4 levels associate with disease activity rather than cancer development in patients with idiopathic inflammatory myopathies. <i>Arthritis Research and Therapy</i> , 2014, 16, 468.	1.6	13
78	Association of circulating miR-223 and miR-16 with disease activity in patients with early rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 1898-1904.	0.5	165
79	A3.30â€¦Plasma level of HSP70 protein is increased in czech patients with idiopathic inflammatory myopathy. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, A54.1-A54.	0.5	1
80	Inflammatory myopathy associated with statins: report of three cases. <i>Modern Rheumatology</i> , 2014, 24, 366-371.	0.9	7
81	OP0289â€¦A Prospective, Randomized, Open-Label, Assessor-Blind, Multicenter Study of Efficacy and Safety of Combined Treatment of Methotrexate + Glucocorticoids versus Glucocorticoids Alone in Patients with Polymyositis and Dermatomyositis (Prometheus Trial). <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 171.1-171.	0.5	3
82	FRI0520â€¦Association Study of the BAFF Genetic Variations in Two Independent Cohorts with Idiopathic Inflammatory Myopathies. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 575.3-576.	0.5	0
83	AB0009â€¦Association of MHC Haplotypes with Idiopathic Inflammatory Myopathy in Cohort of Czech Patients. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 806.3-806.	0.5	0
84	SAT0188â€¦Serum Calprotectin (S100A8/9) Correlates with Clinical and Ultrasound Outcomes in Patients with Early Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 658.1-658.	0.5	0
85	A3.29â€¦Micro-RNA molecules are clearly involved in pathogenesis of idiopathic inflammatory myopathy. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, A53.3-A54.	0.5	1
86	Rituximab in the treatment of refractory adult and juvenile dermatomyositis and adult polymyositis: A randomized, placeboâ€¦phase trial. <i>Arthritis and Rheumatism</i> , 2013, 65, 314-324.	6.7	514
87	A10.3â€¦Arthritis in Patients with Idiopathic Inflammatory Myopathies. <i>Annals of the Rheumatic Diseases</i> , 2013, 72, A72.2-A72.	0.5	1
88	A7.3â€¦Association of Circulating miR-223 and miR-16 with Disease Activity in Patients with Early Rheumatoid Arthritis. <i>Annals of the Rheumatic Diseases</i> , 2013, 72, A48.3-A49.	0.5	0
89	Inflammatory myopathy associated with statins: report of three cases. <i>Modern Rheumatology</i> , 2012, , 1.	0.9	0
90	Anti-Ro52 epitope mapping in inflammatory myopathies. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, A50.1-A50.	0.5	0

#	ARTICLE	IF	CITATIONS
91	Resistin in idiopathic inflammatory myopathies. <i>Arthritis Research and Therapy</i> , 2012, 14, R111.	1.6	21
92	Decreases in serum levels of S100A8/9 (calprotectin) correlate with improvements in total swollen joint count in patients with recent-onset rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2011, 13, R122.	1.6	69
93	Clinical trials roundup in idiopathic inflammatory myopathies. <i>Current Opinion in Rheumatology</i> , 2011, 23, 605-611.	2.0	9
94	The expression regulation of the HSPA1B gene in patients with myositis is not dependent on the presence of HLA-DRB1*03 risk allele. <i>Annals of the Rheumatic Diseases</i> , 2011, 70, A19-A20.	0.5	1
95	Serum levels of interferon γ do not correlate with disease activity in patients with dermatomyositis/polymyositis. <i>Annals of the Rheumatic Diseases</i> , 2011, 70, 879-880.	0.5	17
96	The metastasis promoting protein S100A4 is increased in idiopathic inflammatory myopathies. <i>Rheumatology</i> , 2011, 50, 1766-1772.	0.9	35
97	Delays in assessment of patients with rheumatoid arthritis: variations across Europe. <i>Annals of the Rheumatic Diseases</i> , 2011, 70, 1822-1825.	0.5	112
98	Treatment-resistant inflammatory myopathy. <i>Best Practice and Research in Clinical Rheumatology</i> , 2010, 24, 427-440.	1.4	10
99	Increased serum concentration of immune cell derived microparticles in polymyositis/dermatomyositis. <i>Immunology Letters</i> , 2010, 128, 124-130.	1.1	35
100	Increased serum levels of B cell activating factor (BAFF) in subsets of patients with idiopathic inflammatory myopathies. <i>Annals of the Rheumatic Diseases</i> , 2009, 68, 836-843.	0.5	95
101	Factors influencing health status and disability of patients with ankylosing spondylitis in the Czech Republic. <i>Clinical Rheumatology</i> , 2008, 27, 1005-1013.	1.0	25