## Anna Sankiewicz

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4429729/publications.pdf Version: 2024-02-01



ANNA SANKIEWICZ

#	Article	IF	CITATIONS
1	An SPR imaging immunosensor for leptin determination in blood plasma. Analytical Methods, 2021, 13, 642-646.	2.7	12
2	Two SPRi biosensors for the determination of cathepsin S in blood plasma. Talanta, 2021, 225, 121900.	5.5	7
3	Application of SPRi Biosensors for Determination of 20S Proteasome and UCH-L1 Levels in the Serum and Urine of Transitional Bladder Cancer Patients. Applied Sciences (Switzerland), 2021, 11, 7835.	2.5	3
4	Levels of Selected Matrix Metalloproteinases—MMP-1, MMP-2 and Fibronectin in the Saliva of Patients Planned for Endodontic Treatment or Surgical Extraction. Journal of Clinical Medicine, 2020, 9, 3971.	2.4	5
5	Plasma level of laminin 5 and collagen IV in cryptorchidism. Advances in Medical Sciences, 2020, 65, 176-181.	2.1	1
6	Plasma concentration of MMP-1 and MMP-2 in boys with cryptorchidism and its lack of correlation with INSL3 and inhibin B. Scandinavian Journal of Clinical and Laboratory Investigation, 2019, 79, 412-418.	1.2	4
7	A New Analytical Method for Determination of Cathepsin L Based on the Surface Plasmon Resonance Imaging Biosensor. International Journal of Molecular Sciences, 2019, 20, 2166.	4.1	6
8	Matrix metalloproteinase-2 and its correlation with basal membrane components laminin-5 and collagen type IV in paediatric burn patients measured with Surface Plasmon Resonance Imaging (SPRI) biosensors. Burns, 2018, 44, 931-940.	1.9	19
9	Concentration of Proteasome in the Blood Plasma of Children with Acute Appendicitis, Before and After Surgery, and Its Correlation with CRP. World Journal of Surgery, 2018, 42, 2259-2264.	1.6	3
10	Immunoproteasome in the Plasma of Pediatric Patients With Moderate and Major Burns, and Its Correlation With Proteasome and UCHL1 Measured by SPR Imaging Biosensors. Journal of Burn Care and Research, 2018, 39, 948-953.	0.4	6
11	SPR imaging biosensor for the quantitation of fibronectin concentration in blood samples. Journal of Pharmaceutical and Biomedical Analysis, 2018, 150, 1-8.	2.8	37
12	Determination of the concentration of cathepsin B bySPRI biosensor in children with appendicitis, and its correlation with proteasomes. Advances in Clinical and Experimental Medicine, 2018, 27, 1529-1534.	1.4	0
13	Overexpression of ubiquitin carboxyl-terminal hydrolase L1 (UCHL1) in serum of children after thermal injury. Advances in Medical Sciences, 2017, 62, 83-86.	2.1	24
14	Methods for 20S Immunoproteasome and 20S Constitutive Proteasome Determination Based on SPRI Biosensors. Cellular and Molecular Bioengineering, 2017, 10, 174-185.	2.1	14
15	Podoplanin serum and urine concentration in transitional bladder cancer. Cancer Biomarkers, 2016, 16, 343-350.	1.7	25
16	Application of SPR Imaging Biosensor for the Measurement of 20S Proteasomes in Blood Plasma of Children with Thermal Injury. Annals of Clinical and Laboratory Science, 2016, 46, 407-11.	0.2	11
17	Development of surface plasmon resonance imaging biosensors for detection of ubiquitin carboxyl-terminal hydrolase L1. Analytical Biochemistry, 2015, 469, 4-11.	2.4	33
18	Surface plasmon resonance imaging biosensors for aromatase based on a potent inhibitor and a specific antibody: Sensor development and application for biological material. Open Chemistry, 2014, 12, 557-567.	1.9	18

#	Article	IF	CITATIONS
19	Analytical applications of electrode sensitive to labetalol in pharmaceuticals. Open Chemistry, 2003, 1, 242-259.	1.9	4