

Robert L Hanson

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4431973/robert-l-hanson-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

269
papers

18,317
citations

72
h-index

127
g-index

277
ext. papers

20,340
ext. citations

6.7
avg, IF

5.96
L-index

#	Paper	IF	Citations
269	Epidemiology of Type 2 Diabetes in Indigenous Communities in the United States. <i>Current Diabetes Reports</i> , 2021 , 21, 47	5.6	
268	A missense variant Arg611Cys in LIPE which encodes hormone sensitive lipase decreases lipolysis and increases risk of type 2 diabetes in American Indians. <i>Diabetes/Metabolism Research and Reviews</i> , 2021 , e3504	7.5	0
267	Incidence of diabetes in South Asian young adults compared to Pima Indians. <i>BMJ Open Diabetes Research and Care</i> , 2021 , 9,	4.5	2
266	Next generation sequencing for HLA loci in full heritage Pima Indians of Arizona, Part II: HLA-A, -B, and -C with selected non-classical loci at 4-field resolution from whole genome sequences. <i>Human Immunology</i> , 2021 , 82, 385-403	2.3	
265	Further evidence supporting a potential role for ADH1B in obesity. <i>Scientific Reports</i> , 2021 , 11, 1932	4.9	1
264	Whole genome sequence analyses of eGFR in 23,732 people representing multiple ancestries in the NHLBI trans-omics for precision medicine (TOPMed) consortium. <i>EBioMedicine</i> , 2021 , 63, 103157	8.8	3
263	Exome Sequencing of 21 Bardet-Biedl Syndrome (BBS) Genes to Identify Obesity Variants in 6,851 American Indians. <i>Obesity</i> , 2021 , 29, 748-754	8	0
262	Weight tracking in childhood and adolescence and type 2 diabetes risk. <i>Diabetologia</i> , 2020 , 63, 1753-1763	10.3	2
261	Characterization of Exome Variants and Their Metabolic Impact in 6,716 American Indians from the Southwest US. <i>American Journal of Human Genetics</i> , 2020 , 107, 251-264	11	3
260	Low Serum Insulinlike Growth FactorIII Levels Correlate with High BMI in American Indian Adults. <i>Obesity</i> , 2020 , 28, 676-682	8	
259	Exome Sequencing Identifies A Nonsense Variant in DAO Associated With Reduced Energy Expenditure in American Indians. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020 , 105,	5.6	1
258	Assessment of the potential role of natural selection in type 2 diabetes and related traits across human continental ancestry groups: comparison of phenotypic with genotypic divergence. <i>Diabetologia</i> , 2020 , 63, 2616-2627	10.3	2
257	Racial/ethnic differences in the burden of type 2 diabetes over the life course: a focus on the USA and India. <i>Diabetologia</i> , 2019 , 62, 1751-1760	10.3	21
256	Birthweight and early-onset type 2 diabetes in American Indians: differential effects in adolescents and young adults and additive effects of genotype, BMI and maternal diabetes. <i>Diabetologia</i> , 2019 , 62, 1628-1637	10.3	3
255	Kidney cytosine methylation changes improve renal function decline estimation in patients with diabetic kidney disease. <i>Nature Communications</i> , 2019 , 10, 2461	17.4	30
254	Assessing the Role of 98 Established Loci for BMI in American Indians. <i>Obesity</i> , 2019 , 27, 845-854	8	8
253	Use of graded Semmes Weinstein monofilament testing for ascertaining peripheral neuropathy in people with and without diabetes. <i>Diabetes Research and Clinical Practice</i> , 2019 , 151, 1-10	7.4	7

252	Association of CREBRF variants with obesity and diabetes in Pacific Islanders from Guam and Saipan. <i>Diabetologia</i> , 2019 , 62, 1647-1652	10.3	15
251	A trans-ancestral meta-analysis of genome-wide association studies reveals loci associated with childhood obesity. <i>Human Molecular Genetics</i> , 2019 , 28, 3327-3338	5.6	30
250	Glycemia affects glomerular filtration rate in people with type 2 diabetes. <i>BMC Nephrology</i> , 2019 , 20, 397	2.7	2
249	Next generation sequencing and the classical HLA loci in full heritage Pima Indians of Arizona: Defining the core HLA variation for North American Paleo-Indians. <i>Human Immunology</i> , 2019 , 80, 955-965	2.3	7
248	Identification and functional analysis of a novel G310D variant in the insulin-like growth factor 1 receptor (IGF1R) gene associated with type 2 diabetes in American Indians. <i>Diabetes/Metabolism Research and Reviews</i> , 2018 , 34, e2994	7.5	2
247	Effect of severe obesity in childhood and adolescence on risk of type 2 diabetes in youth and early adulthood in an American Indian population. <i>Pediatric Diabetes</i> , 2018 , 19, 622-629	3.6	17
246	Functional and association analysis of an Amerindian-derived population-specific p.(Thr280Met) variant in RBPJL, a component of the PTF1 complex. <i>European Journal of Human Genetics</i> , 2018 , 26, 238-246	5.3	6
245	Cytosine methylation predicts renal function decline in American Indians. <i>Kidney International</i> , 2018 , 93, 1417-1431	9.9	27
244	White blood cell fractions correlate with lesions of diabetic kidney disease and predict loss of kidney function in Type 2 diabetes. <i>Nephrology Dialysis Transplantation</i> , 2018 , 33, 1001-1009	4.3	11
243	Serum lipids and mortality in an American Indian population: A longitudinal study. <i>Journal of Diabetes and Its Complications</i> , 2018 , 32, 18-26	3.2	6
242	Analysis of type 2 diabetes and obesity genetic variants in Mexican Pima Indians: Marked allelic differentiation among Amerindians at HLA. <i>Annals of Human Genetics</i> , 2018 , 82, 287-299	2.2	5
241	Effect of different methods of accounting for antihypertensive treatment when assessing the relationship between diabetes or obesity and systolic blood pressure. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 693-699	3.2	4
240	Differential methylation of genes in individuals exposed to maternal diabetes in utero. <i>Diabetologia</i> , 2017 , 60, 645-655	10.3	49
239	Admixture Mapping Identifies an Amerindian Ancestry Locus Associated with Albuminuria in Hispanics in the United States. <i>Journal of the American Society of Nephrology: JASN</i> , 2017 , 28, 2211-2220	12.7	17
238	A Genome-Wide Association Study of IVGTT-Based Measures of First-Phase Insulin Secretion Refines the Underlying Physiology of Type 2 Diabetes Variants. <i>Diabetes</i> , 2017 , 66, 2296-2309	0.9	69
237	Associations between persistent organic pollutants, type 2 diabetes, diabetic nephropathy and mortality. <i>Occupational and Environmental Medicine</i> , 2017 , 74, 521-527	2.1	27
236	A Loss-of-Function Splice Acceptor Variant in Is Protective for Type 2 Diabetes. <i>Diabetes</i> , 2017 , 66, 2903-2914	2.9	32
235	Autoantibodies against PFDN2 are associated with an increased risk of type 2 diabetes: A case-control study. <i>Diabetes/Metabolism Research and Reviews</i> , 2017 , 33, e2922	7.5	11

234	Large meta-analysis of genome-wide association studies identifies five loci for lean body mass. <i>Nature Communications</i> , 2017 , 8, 80	17.4	88
233	Growth Tracking in Severely Obese or Underweight Children. <i>Pediatrics</i> , 2017 , 140,	7.4	10
232	One-hour and two-hour postload plasma glucose concentrations are comparable predictors of type 2 diabetes mellitus in Southwestern Native Americans. <i>Diabetologia</i> , 2017 , 60, 1704-1711	10.3	29
231	HbA1c and the Prediction of Type 2 Diabetes in Children and Adults. <i>Diabetes Care</i> , 2017 , 40, 16-21	14.6	54
230	Assessing variation across 8 established East Asian loci for type 2 diabetes mellitus in American Indians: Suggestive evidence for new sex-specific diabetes signals in GLIS3 and ZFAND3. <i>Diabetes/Metabolism Research and Reviews</i> , 2017 , 33, e2869	7.5	12
229	Identity-by-Descent Mapping Identifies Major Locus for Serum Triglycerides in Amerindians Largely Explained by a Founder Mutation. <i>Circulation: Cardiovascular Genetics</i> , 2017 , 10,		10
228	Admixture mapping in the Hispanic Community Health Study/Study of Latinos reveals regions of genetic associations with blood pressure traits. <i>PLoS ONE</i> , 2017 , 12, e0188400	3.7	17
227	A Genome-Wide Association Study Using a Custom Genotyping Array Identifies Variants in Associated With Reduced Energy Expenditure in American Indians. <i>Diabetes</i> , 2017 , 66, 2284-2295	0.9	20
226	Long-term Effect of Losartan on Kidney Disease in American Indians With Type 2 Diabetes: A Follow-up Analysis of a Randomized Clinical Trial. <i>Diabetes Care</i> , 2016 , 39, 2004-2010	14.6	8
225	Selecting SNPs informative for African, American Indian and European Ancestry: application to the Family Investigation of Nephropathy and Diabetes (FIND). <i>BMC Genomics</i> , 2016 , 17, 325	4.5	1
224	Genome-wide Association Study of Platelet Count Identifies Ancestry-Specific Loci in Hispanic/Latino Americans. <i>American Journal of Human Genetics</i> , 2016 , 98, 229-42	11	54
223	Assessment of established HDL-C loci for association with HDL-C levels and type 2 diabetes in Pima Indians. <i>Diabetologia</i> , 2016 , 59, 481-91	10.3	13
222	Analysis of SLC16A11 Variants in 12,811 American Indians: Genotype-Obesity Interaction for Type 2 Diabetes and an Association With RNASEK Expression. <i>Diabetes</i> , 2016 , 65, 510-9	0.9	16
221	Metabolic Risk Factors and Type 2 Diabetes Incidence in American Indian Children. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016 , 101, 1437-44	5.6	16
220	The Arg59Trp variant in ANGPTL8 (betatrophin) is associated with total and HDL-cholesterol in American Indians and Mexican Americans and differentially affects cleavage of ANGPTL3. <i>Molecular Genetics and Metabolism</i> , 2016 , 118, 128-37	3.7	26
219	Role of Established Type 2 Diabetes-Susceptibility Genetic Variants in a High Prevalence American Indian Population. <i>Diabetes</i> , 2015 , 64, 2646-57	0.9	23
218	A cis-eQTL in PFKFB2 is associated with diabetic nephropathy, adiposity and insulin secretion in American Indians. <i>Human Molecular Genetics</i> , 2015 , 24, 2985-96	5.6	11
217	Urinary monocyte chemoattractant protein-1 and hepcidin and early diabetic nephropathy lesions in type 1 diabetes mellitus. <i>Nephrology Dialysis Transplantation</i> , 2015 , 30, 599-606	4.3	21

216	Environmentally Driven Increases in Type 2 Diabetes and Obesity in Pima Indians and Non-Pimas in Mexico Over a 15-Year Period: The Maycoba Project. <i>Diabetes Care</i> , 2015 , 38, 2075-82	14.6	28
215	The transcriptional landscape of age in human peripheral blood. <i>Nature Communications</i> , 2015 , 6, 8570	17.4	335
214	ABCC8 R1420H Loss-of-Function Variant in a Southwest American Indian Community: Association With Increased Birth Weight and Doubled Risk of Type 2 Diabetes. <i>Diabetes</i> , 2015 , 64, 4322-32	0.9	37
213	Association of urinary KIM-1, L-FABP, NAG and NGAL with incident end-stage renal disease and mortality in American Indians with type 2 diabetes mellitus. <i>Diabetologia</i> , 2015 , 58, 188-98	10.3	68
212	Assessing FOXO1A as a potential susceptibility locus for type 2 diabetes and obesity in American Indians. <i>Obesity</i> , 2015 , 23, 1960-5	8	6
211	The effect of differing patterns of childhood body mass index gain on adult physiology in American Indians. <i>Obesity</i> , 2015 , 23, 1872-80	8	6
210	Genome-Wide Association and Trans-ethnic Meta-Analysis for Advanced Diabetic Kidney Disease: Family Investigation of Nephropathy and Diabetes (FIND). <i>PLoS Genetics</i> , 2015 , 11, e1005352	6	84
209	Use of a High-Density Protein Microarray to Identify Autoantibodies in Subjects with Type 2 Diabetes Mellitus and an HLA Background Associated with Reduced Insulin Secretion. <i>PLoS ONE</i> , 2015 , 10, e0143551	3.7	11
208	Prospective association of a genetic risk score and lifestyle intervention with cardiovascular morbidity and mortality among individuals with type 2 diabetes: the Look AHEAD randomised controlled trial. <i>Diabetologia</i> , 2015 , 58, 1803-13	10.3	15
207	Association Studies to Map Genes for Disease-Related Traits in Humans 2015 , 53-66		
206	A genome-wide association study in American Indians implicates DNER as a susceptibility locus for type 2 diabetes. <i>Diabetes</i> , 2014 , 63, 369-76	0.9	49
205	The influence of rare genetic variation in SLC30A8 on diabetes incidence and β -cell function. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014 , 99, E926-30	5.6	19
204	Common genetic variation in the glucokinase gene (GCK) is associated with type 2 diabetes and rates of carbohydrate oxidation and energy expenditure. <i>Diabetologia</i> , 2014 , 57, 1382-90	10.3	27
203	Potential epigenetic dysregulation of genes associated with MODY and type 2 diabetes in humans exposed to a diabetic intrauterine environment: an analysis of genome-wide DNA methylation. <i>Metabolism: Clinical and Experimental</i> , 2014 , 63, 654-60	12.7	44
202	Weight maintenance from young adult weight predicts better health outcomes. <i>Obesity</i> , 2014 , 22, 2361-9		10
201	Assessing accuracy of genotype imputation in American Indians. <i>PLoS ONE</i> , 2014 , 9, e102544	3.7	2
200	Variants associated with type 2 diabetes identified by the transethnic meta-analysis study: assessment in American Indians and evidence for a new signal in LPP. <i>Diabetologia</i> , 2014 , 57, 2334-8	10.3	7
199	Study design of the Maycoba Project: obesity and diabetes in Mexican Pimas. <i>American Journal of Health Behavior</i> , 2014 , 38, 370-8	1.9	5

198	Whole exome sequencing identifies variation in CYB5A and RNF10 associated with adiposity and type 2 diabetes. <i>Obesity</i> , 2014 , 22, 984-8	8	28
197	Common genetic variation in and near the melanocortin 4 receptor gene (MC4R) is associated with body mass index in American Indian adults and children. <i>Human Genetics</i> , 2014 , 133, 1431-41	6.3	17
196	Identification of genetic variation that determines human trehalase activity and its association with type 2 diabetes. <i>Human Genetics</i> , 2013 , 132, 697-707	6.3	13
195	Effect of losartan on prevention and progression of early diabetic nephropathy in American Indians with type 2 diabetes. <i>Diabetes</i> , 2013 , 62, 3224-31	0.9	61
194	Strong parent-of-origin effects in the association of KCNQ1 variants with type 2 diabetes in American Indians. <i>Diabetes</i> , 2013 , 62, 2984-91	0.9	51
193	Comparison of serum cystatin C, serum creatinine, measured GFR, and estimated GFR to assess the risk of kidney failure in American Indians with diabetic nephropathy. <i>American Journal of Kidney Diseases</i> , 2013 , 62, 33-41	7.4	27
192	MAP2K3 is associated with body mass index in American Indians and Caucasians and may mediate hypothalamic inflammation. <i>Human Molecular Genetics</i> , 2013 , 22, 4438-49	5.6	19
191	Evidence for a role of LPGAT1 in influencing BMI and percent body fat in Native Americans. <i>Obesity</i> , 2013 , 21, 193-202	8	15
190	An ACACB variant implicated in diabetic nephropathy associates with body mass index and gene expression in obese subjects. <i>PLoS ONE</i> , 2013 , 8, e56193	3.7	9
189	A genome-wide search for linkage of estimated glomerular filtration rate (eGFR) in the Family Investigation of Nephropathy and Diabetes (FIND). <i>PLoS ONE</i> , 2013 , 8, e81888	3.7	23
188	Variants in the LEPR gene are nominally associated with higher BMI and lower 24-h energy expenditure in Pima Indians. <i>Obesity</i> , 2012 , 20, 2426-30	8	17
187	New susceptibility loci associated with kidney disease in type 1 diabetes. <i>PLoS Genetics</i> , 2012 , 8, e1002921	17.6	
186	Greater impact of melanocortin-4 receptor deficiency on rates of growth and risk of type 2 diabetes during childhood compared with adulthood in Pima Indians. <i>Diabetes</i> , 2012 , 61, 250-7	0.9	46
185	Association of variants in the carnosine peptidase 1 gene (CNDP1) with diabetic nephropathy in American Indians. <i>Molecular Genetics and Metabolism</i> , 2011 , 103, 185-90	3.7	14
184	A genome-wide association study of BMI in American Indians. <i>Obesity</i> , 2011 , 19, 2102-6	8	27
183	Higher energy expenditure in humans predicts natural mortality. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011 , 96, E972-6	5.6	31
182	SIRT1 is associated with a decrease in acute insulin secretion and a sex specific increase in risk for type 2 diabetes in Pima Indians. <i>Molecular Genetics and Metabolism</i> , 2011 , 104, 661-5	3.7	53
181	HLA-DRB1 reduces the risk of type 2 diabetes mellitus by increased insulin secretion. <i>Diabetologia</i> , 2011 , 54, 1684-92	10.3	28

180	Bimodal distribution of RNA expression levels in human skeletal muscle tissue. <i>BMC Genomics</i> , 2011 , 12, 98	4.5	25
179	Genomewide linkage scan for diabetic renal failure and albuminuria: the FIND study. <i>American Journal of Nephrology</i> , 2011 , 33, 381-9	4.6	48
178	Albuminuria and estimated glomerular filtration rate as predictors of diabetic end-stage renal disease and death. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2011 , 6, 2444-51	6.9	94
177	Secular trends in treatment and control of type 2 diabetes in an American Indian population: a 30-year longitudinal study. <i>Diabetes Care</i> , 2010 , 33, 2383-9	14.6	10
176	Functional variants in MBL2 are associated with type 2 diabetes and pre-diabetes traits in Pima Indians and the old order Amish. <i>Diabetes</i> , 2010 , 59, 2080-5	0.9	14
175	Variants in ASK1 are associated with skeletal muscle ASK1 expression, in vivo insulin resistance, and type 2 diabetes in Pima Indians. <i>Diabetes</i> , 2010 , 59, 1276-82	0.9	17
174	Predictive value of albuminuria in American Indian youth with or without type 2 diabetes. <i>Pediatrics</i> , 2010 , 125, e844-51	7.4	19
173	Effect of intrauterine diabetes exposure on the incidence of end-stage renal disease in young adults with type 2 diabetes. <i>Diabetes Care</i> , 2010 , 33, 2396-8	14.6	15
172	Evaluation of A2BP1 as an obesity gene. <i>Diabetes</i> , 2010 , 59, 2837-45	0.9	35
171	ELMO1 variants and susceptibility to diabetic nephropathy in American Indians. <i>Molecular Genetics and Metabolism</i> , 2010 , 101, 383-90	3.7	36
170	The separate and joint effects of prolonged QT interval and heart rate on mortality. <i>Atherosclerosis</i> , 2010 , 209, 539-44	3.1	3
169	Common variants in 40 genes assessed for diabetes incidence and response to metformin and lifestyle intervention in the diabetes prevention program. <i>Diabetes</i> , 2010 , 59, 2672-81	0.9	200
168	Childhood obesity, other cardiovascular risk factors, and premature death. <i>New England Journal of Medicine</i> , 2010 , 362, 485-93	59.2	903
167	Variants in ACAD10 are associated with type 2 diabetes, insulin resistance and lipid oxidation in Pima Indians. <i>Diabetologia</i> , 2010 , 53, 1349-53	10.3	27
166	Linkage disequilibrium mapping of the replicated type 2 diabetes linkage signal on chromosome 1q. <i>Diabetes</i> , 2009 , 58, 1704-9	0.9	23
165	The association of ENPP1 K121Q with diabetes incidence is abolished by lifestyle modification in the diabetes prevention program. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2009 , 94, 449-55	5.6	45
164	Common variation in SIM1 is reproducibly associated with BMI in Pima Indians. <i>Diabetes</i> , 2009 , 58, 1682-9	0.9	40
163	Genome-wide linkage scans for type 2 diabetes mellitus in four ethnically diverse populations-significant evidence for linkage on chromosome 4q in African Americans: the Family Investigation of Nephropathy and Diabetes Research Group. <i>Diabetes/Metabolism Research and Reviews</i> , 2009 , 25, 710-7	7.5	9

162	The effect of salsalate on insulin action and glucose tolerance in obese non-diabetic patients: results of a randomised double-blind placebo-controlled study. <i>Diabetologia</i> , 2009 , 52, 385-93	10.3	102
161	Association analysis of variation in/near FTO, CDKAL1, SLC30A8, HHEX, EXT2, IGF2BP2, LOC387761, and CDKN2B with type 2 diabetes and related quantitative traits in Pima Indians. <i>Diabetes</i> , 2009 , 58, 478-88	0.9	120
160	Physical activity levels in American-Indian adults: the Strong Heart Family Study. <i>American Journal of Preventive Medicine</i> , 2009 , 37, 481-7	6.1	29
159	Predictive power of sequential measures of albuminuria for progression to ESRD or death in Pima Indians with type 2 diabetes. <i>American Journal of Kidney Diseases</i> , 2008 , 51, 759-66	7.4	36
158	Changing course of diabetic nephropathy in the Pima Indians. <i>Diabetes Research and Clinical Practice</i> , 2008 , 82 Suppl 1, S10-4	7.4	6
157	Genome-wide scan for estimated glomerular filtration rate in multi-ethnic diabetic populations: the Family Investigation of Nephropathy and Diabetes (FIND). <i>Diabetes</i> , 2008 , 57, 235-43	0.9	75
156	PCLO variants are nominally associated with early-onset type 2 diabetes and insulin resistance in Pima Indians. <i>Diabetes</i> , 2008 , 57, 3156-60	0.9	10
155	Extension of type 2 diabetes genome-wide association scan results in the diabetes prevention program. <i>Diabetes</i> , 2008 , 57, 2503-10	0.9	86
154	Plasma glucose regulation and mortality in pima Indians. <i>Diabetes Care</i> , 2008 , 31, 488-92	14.6	14
153	Lower metabolic rate in individuals heterozygous for either a frameshift or a functional missense MC4R variant. <i>Diabetes</i> , 2008 , 57, 3267-72	0.9	45
152	Association analysis of Krüppel-like factor 11 variants with type 2 diabetes in Pima Indians. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008 , 93, 3644-9	5.6	9
151	Relation of central adiposity and body mass index to the development of diabetes in the Diabetes Prevention Program. <i>American Journal of Clinical Nutrition</i> , 2008 , 87, 1212-8	7	177
150	Heritability of the severity of diabetic retinopathy: the FIND-Eye study 2008 , 49, 3839-45		133
149	Design and analysis of genetic association studies to finely map a locus identified by linkage analysis: assessment of the extent to which an association can account for the linkage. <i>Annals of Human Genetics</i> , 2008 , 72, 126-39	2.2	6
148	Diabetic nephropathy in American Indians, with a special emphasis on the Pima Indians. <i>Current Diabetes Reports</i> , 2008 , 8, 486-93	5.6	16
147	Identification of PVT1 as a candidate gene for end-stage renal disease in type 2 diabetes using a pooling-based genome-wide single nucleotide polymorphism association study. <i>Diabetes</i> , 2007 , 56, 975-83	6.9	156
146	Homocysteine and vitamin B(12) concentrations and mortality rates in type 2 diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2007 , 23, 193-201	7.5	19
145	Meta-analysis of genome-wide linkage studies in BMI and obesity. <i>Obesity</i> , 2007 , 15, 2263-75	8	122

144	Protein tyrosine phosphatase 1B is not a major susceptibility gene for type 2 diabetes mellitus or obesity among Pima Indians. <i>Diabetologia</i> , 2007 , 50, 985-9	10.3	14
143	The Pro12Ala variant at the peroxisome proliferator-activated receptor gamma gene and change in obesity-related traits in the Diabetes Prevention Program. <i>Diabetologia</i> , 2007 , 50, 2451-60	10.3	62
142	A search for variants associated with young-onset type 2 diabetes in American Indians in a 100K genotyping array. <i>Diabetes</i> , 2007 , 56, 3045-52	0.9	79
141	Progression to type 2 diabetes characterized by moderate then rapid glucose increases. <i>Diabetes</i> , 2007 , 56, 2054-61	0.9	64
140	Variants in ARHGEF11, a candidate gene for the linkage to type 2 diabetes on chromosome 1q, are nominally associated with insulin resistance and type 2 diabetes in Pima Indians. <i>Diabetes</i> , 2007 , 56, 1454-9	0.9	29
139	TCF7L2 is not a major susceptibility gene for type 2 diabetes in Pima Indians: analysis of 3,501 individuals. <i>Diabetes</i> , 2007 , 56, 3082-8	0.9	64
138	Childhood predictors of young-onset type 2 diabetes. <i>Diabetes</i> , 2007 , 56, 2964-72	0.9	123
137	Genome-wide linkage analyses to identify Loci for diabetic retinopathy. <i>Diabetes</i> , 2007 , 56, 1160-6	0.9	92
136	Variants in the Ca V 2.3 (alpha 1E) subunit of voltage-activated Ca ²⁺ channels are associated with insulin resistance and type 2 diabetes in Pima Indians. <i>Diabetes</i> , 2007 , 56, 3089-94	0.9	27
135	Prediction of diabetic nephropathy using urine proteomic profiling 10 years prior to development of nephropathy. <i>Diabetes Care</i> , 2007 , 30, 638-43	14.6	106
134	Changing patterns of type 2 diabetes incidence among Pima Indians. <i>Diabetes Care</i> , 2007 , 30, 1758-63	14.6	88
133	Common variation in the LMNA gene (encoding lamin A/C) and type 2 diabetes: association analyses in 9,518 subjects. <i>Diabetes</i> , 2007 , 56, 879-83	0.9	27
132	Meta-analysis of genome-wide linkage studies of quantitative lipid traits in families ascertained for type 2 diabetes. <i>Diabetes</i> , 2007 , 56, 890-6	0.9	27
131	Body size and shape changes and the risk of diabetes in the diabetes prevention program. <i>Diabetes</i> , 2007 , 56, 1680-5	0.9	86
130	A genomewide single-nucleotide-polymorphism panel for Mexican American admixture mapping. <i>American Journal of Human Genetics</i> , 2007 , 80, 1014-23	11	113
129	Genome-wide scans for diabetic nephropathy and albuminuria in multiethnic populations: the family investigation of nephropathy and diabetes (FIND). <i>Diabetes</i> , 2007 , 56, 1577-85	0.9	125
128	Design and analysis of genetic association studies to finely map a locus identified by linkage analysis: sample size and power calculations. <i>Annals of Human Genetics</i> , 2006 , 70, 332-49	2.2	18
127	Gestational glucose tolerance and risk of type 2 diabetes in young Pima Indian offspring. <i>Diabetes</i> , 2006 , 55, 460-5	0.9	180

126	A functional Tyr1306Cys variant in LARG is associated with increased insulin action in vivo. <i>Diabetes</i> , 2006 , 55, 1497-503	0.9	14
125	Variation within the gene encoding the upstream stimulatory factor 1 does not influence susceptibility to type 2 diabetes in samples from populations with replicated evidence of linkage to chromosome 1q. <i>Diabetes</i> , 2006 , 55, 2541-8	0.9	33
124	IL6 gene promoter polymorphisms and type 2 diabetes: joint analysis of individual participantsO data from 21 studies. <i>Diabetes</i> , 2006 , 55, 2915-21	0.9	91
123	Electrocardiographic abnormalities predict deaths from cardiovascular disease and ischemic heart disease in Pima Indians with type 2 diabetes. <i>American Heart Journal</i> , 2006 , 151, 1080-6	4.9	10
122	Variants in the gene encoding aldose reductase (AKR1B1) and diabetic nephropathy in American Indians. <i>Diabetic Medicine</i> , 2006 , 23, 367-76	3.5	20
121	Genetic variation in UCP2 (uncoupling protein-2) is associated with energy metabolism in Pima Indians. <i>Diabetologia</i> , 2005 , 48, 2292-5	10.3	51
120	Meprin beta metalloprotease gene polymorphisms associated with diabetic nephropathy in the Pima Indians. <i>Human Genetics</i> , 2005 , 118, 12-22	6.3	32
119	Habitual physical activity in children: the role of genes and the environment. <i>American Journal of Clinical Nutrition</i> , 2005 , 82, 901-8	7	83
118	Variants in hepatocyte nuclear factor 4alpha are modestly associated with type 2 diabetes in Pima Indians. <i>Diabetes</i> , 2005 , 54, 3035-9	0.9	43
117	Periodontal disease and mortality in type 2 diabetes. <i>Diabetes Care</i> , 2005 , 28, 27-32	14.6	299
116	Common Polymorphisms in the Adiponectin Gene ACDC Are Not Associated With Diabetes in Pima Indians. <i>Diabetes</i> , 2005 , 54, 284-9	0.9	75
115	Variations in peptide YY and Y2 receptor genes are associated with severe obesity in Pima Indian men. <i>Diabetes</i> , 2005 , 54, 1598-602	0.9	43
114	Adiponectin concentrations are influenced by renal function and diabetes duration in Pima Indians with type 2 diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2004 , 89, 4010-7	5.6	111
113	Validity of the SMAST in two American Indian tribal populations. <i>Substance Use and Misuse</i> , 2004 , 39, 601-24	2.2	9
112	A novel missense substitution (Val1483Ile) in the fatty acid synthase gene (FAS) is associated with percentage of body fat and substrate oxidation rates in nondiabetic Pima Indians. <i>Diabetes</i> , 2004 , 53, 1915-9	0.9	45
111	Depression, diabetes, and glycemic control in Pima Indians. <i>Diabetes Care</i> , 2004 , 27, 618-9	14.6	18
110	Hematocrit and the incidence of type 2 diabetes in the pima indians. <i>Diabetes Care</i> , 2004 , 27, 2245-6	14.6	22
109	Genetic studies of the etiology of type 2 diabetes in Pima Indians: hunting for pieces to a complicated puzzle. <i>Diabetes</i> , 2004 , 53, 1181-6	0.9	78

108	Interaction between an 11betaHSD1 gene variant and birth era modifies the risk of hypertension in Pima Indians. <i>Hypertension</i> , 2004 , 44, 681-8	8.5	55
107	Both subcutaneous and visceral adipose tissue correlate highly with insulin resistance in african americans. <i>Obesity</i> , 2004 , 12, 1352-9		77
106	Mexican American ancestry-informative markers: examination of population structure and marker characteristics in European Americans, Mexican Americans, Amerindians and Asians. <i>Human Genetics</i> , 2004 , 114, 263-71	6.3	87
105	Alcohol consumption predicts hypertension but not diabetes. <i>Journal of Studies on Alcohol and Drugs</i> , 2004 , 65, 184-90		39
104	The role of insulin receptor substrate-1 gene (IRS1) in type 2 diabetes in Pima Indians. <i>Diabetes</i> , 2003 , 52, 3005-9	0.9	52
103	Genome-wide linkage analysis of serum adiponectin in the Pima Indian population. <i>Diabetes</i> , 2003 , 52, 2419-25	0.9	84
102	Longitudinal studies of incidence and progression of diabetic retinopathy assessed by retinal photography in pima indians. <i>Diabetes Care</i> , 2003 , 26, 320-6	14.6	46
101	Incidence of retinopathy and nephropathy in youth-onset compared with adult-onset type 2 diabetes. <i>Diabetes Care</i> , 2003 , 26, 76-81	14.6	102
100	The insulin gene variable number tandem repeat class I/III polymorphism is in linkage disequilibrium with birth weight but not Type 2 diabetes in the Pima population. <i>Diabetes</i> , 2003 , 52, 187-93	0.9	60
99	Homocysteine as a risk factor for nephropathy and retinopathy in Type 2 diabetes. <i>Diabetologia</i> , 2003 , 46, 766-72	10.3	78
98	Quantitative trait linkage studies of diabetes-related traits. <i>Current Diabetes Reports</i> , 2003 , 3, 176-83	5.6	11
97	The interleukin-6 (-174) G/C promoter polymorphism is associated with type-2 diabetes mellitus in Native Americans and Caucasians. <i>Human Genetics</i> , 2003 , 112, 409-13	6.3	145
96	Inflammatory markers, adiponectin, and risk of type 2 diabetes in the Pima Indian. <i>Diabetes Care</i> , 2003 , 26, 1745-51	14.6	279
95	Higher prevalence of type 2 diabetes, metabolic syndrome and cardiovascular diseases in gypsies than in non-gypsies in Slovakia. <i>Diabetes Research and Clinical Practice</i> , 2003 , 62, 95-103	7.4	66
94	A C-reactive protein promoter polymorphism is associated with type 2 diabetes mellitus in Pima Indians. <i>Molecular Genetics and Metabolism</i> , 2003 , 78, 136-44	3.7	78
93	Physical activity, obesity, and the incidence of type 2 diabetes in a high-risk population. <i>American Journal of Epidemiology</i> , 2003 , 158, 669-75	3.8	152
92	Metabolic effects of the Gly1057Asp polymorphism in IRS-2 and interactions with obesity. <i>Diabetes</i> , 2003 , 52, 1544-50	0.9	38
91	Do measures of body fat distribution provide information on the risk of type 2 diabetes in addition to measures of general obesity? Comparison of anthropometric predictors of type 2 diabetes in Pima Indians. <i>Diabetes Care</i> , 2003 , 26, 2556-61	14.6	128

90	Weight, adiposity, and physical activity as determinants of an insulin sensitivity index in pima Indian children. <i>Diabetes Care</i> , 2003 , 26, 2524-30	14.6	35
89	Identification of susceptibility genes for complex metabolic diseases. <i>Annals of the New York Academy of Sciences</i> , 2002 , 967, 1-6	6.5	12
88	Early excess weight gain of children in the Pima Indian population. <i>Pediatrics</i> , 2002 , 109, E33	7.4	31
87	Genome-wide linkage analysis assessing parent-of-origin effects in the inheritance of birth weight. <i>Human Genetics</i> , 2002 , 110, 503-9	6.3	63
86	Gene for susceptibility to diabetic nephropathy in type 2 diabetes maps to 18q22.3-23. <i>Kidney International</i> , 2002 , 62, 2176-83	9.9	121
85	Currently identified genes affecting insulin resistance are not associated with birth weight in the Pima population. <i>Diabetic Medicine</i> , 2002 , 19, 882-4	3.5	4
84	The Gln223Arg polymorphism of the leptin receptor in Pima Indians: influence on energy expenditure, physical activity and lipid metabolism. <i>International Journal of Obesity</i> , 2002 , 26, 1629-32	5.5	85
83	Plasma concentrations of macrophage migration inhibitory factor are elevated in Pima Indians compared to Caucasians and are associated with insulin resistance. <i>Diabetologia</i> , 2002 , 45, 1739-41	10.3	39
82	Components of the "metabolic syndrome" and incidence of type 2 diabetes. <i>Diabetes</i> , 2002 , 51, 3120-7	0.9	456
81	Adiponectin and development of type 2 diabetes in the Pima Indian population. <i>Lancet, The</i> , 2002 , 360, 57-8	40	904
80	Ethnic-difference markers for use in mapping by admixture linkage disequilibrium. <i>American Journal of Human Genetics</i> , 2002 , 70, 737-50	11	113
79	Molecular analysis of KCNJ10 on 1q as a candidate gene for Type 2 diabetes in Pima Indians. <i>Diabetes</i> , 2002 , 51, 3342-6	0.9	13
78	Validity of the CAGE questionnaire in an American Indian population. <i>Journal of Studies on Alcohol and Drugs</i> , 2001 , 62, 294-300		23
77	Association of physical activity and serum insulin concentrations in two populations at high risk for type 2 diabetes but differing by BMI. <i>Diabetes Care</i> , 2001 , 24, 1175-80	14.6	55
76	Genetics of diabetic nephropathy in the Pima Indians. <i>Current Diabetes Reports</i> , 2001 , 1, 275-81	5.6	36
75	Analysis of the association between diabetic nephropathy and polymorphisms in the aldose reductase gene in Type 1 and Type 2 diabetes mellitus. <i>Diabetic Medicine</i> , 2001 , 18, 906-14	3.5	37
74	Family and genetic studies of indices of insulin sensitivity and insulin secretion in Pima Indians. <i>Diabetes/Metabolism Research and Reviews</i> , 2001 , 17, 296-303	7.5	37
73	Joint swelling as a predictor of death from cardiovascular disease in a population study of Pima Indians. <i>Arthritis and Rheumatism</i> , 2001 , 44, 1170-6		50

72	Analysis of the lamin A/C gene as a candidate for type II diabetes susceptibility in Pima Indians. <i>Diabetologia</i> , 2001 , 44, 779-82	10.3	24
71	Changes in BMI and weight before and after the development of type 2 diabetes. <i>Diabetes Care</i> , 2001 , 24, 1917-22	14.6	94
70	Comparison of the effect of plasma glucose concentrations on microvascular disease between Pima Indian youths and adults. <i>Diabetes Care</i> , 2001 , 24, 1023-8	14.6	11
69	The effect of Indian or Anglo dietary preference on the incidence of diabetes in Pima Indians. <i>Diabetes Care</i> , 2001 , 24, 811-6	14.6	41
68	Genome-wide linkage analysis assessing parent-of-origin effects in the inheritance of type 2 diabetes and BMI in Pima Indians. <i>Diabetes</i> , 2001 , 50, 2850-7	0.9	98
67	Body mass index as a measure of adiposity in children and adolescents: relationship to adiposity by dual energy x-ray absorptiometry and to cardiovascular risk factors. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001 , 86, 4061-7	5.6	129
66	Gamma globulin levels predict type 2 diabetes in the Pima Indian population. <i>Diabetes</i> , 2001 , 50, 1598-603	0.9	54
65	Subcutaneous abdominal adipocyte size, a predictor of type 2 diabetes, is linked to chromosome 1q21-q23 and is associated with a common polymorphism in LMNA in Pima Indians. <i>Molecular Genetics and Metabolism</i> , 2001 , 72, 231-8	3.7	62
64	Analysis of linkage disequilibrium between polymorphisms in the KCNJ9 gene with type 2 diabetes mellitus in Pima Indians. <i>Molecular Genetics and Metabolism</i> , 2001 , 73, 97-103	3.7	21
63	Assessment of parent-of-origin effects in linkage analysis of quantitative traits. <i>American Journal of Human Genetics</i> , 2001 , 68, 951-62	11	65
62	Family and genetic studies of indices of insulin sensitivity and insulin secretion in Pima Indians 2001 , 17, 296		1
61	A calpain-10 gene polymorphism is associated with reduced muscle mRNA levels and insulin resistance. <i>Journal of Clinical Investigation</i> , 2000 , 106, R69-73	15.9	216
60	A high fasting plasma insulin concentration predicts type 2 diabetes independent of insulin resistance: evidence for a pathogenic role of relative hyperinsulinemia. <i>Diabetes</i> , 2000 , 49, 2094-101	0.9	212
59	An autosomal genomic scan for loci linked to plasma leptin concentration in Pima Indians. <i>International Journal of Obesity</i> , 2000 , 24, 559-65	5.5	26
58	Novel polymorphisms in the neuropeptide-Y Y5 receptor associated with obesity in Pima Indians. <i>International Journal of Obesity</i> , 2000 , 24, 580-4	5.5	43
57	Association of dopamine D2 receptor polymorphisms Ser311Cys and TaqIA with obesity or type 2 diabetes mellitus in Pima Indians. <i>International Journal of Obesity</i> , 2000 , 24, 1233-8	5.5	50
56	Type 2 diabetes and low birth weight: the role of paternal inheritance in the association of low birth weight and diabetes. <i>Diabetes</i> , 2000 , 49, 445-9	0.9	160
55	The 1997 American Diabetes Association and 1999 World Health Organization criteria for hyperglycemia in the diagnosis and prediction of diabetes. <i>Diabetes Care</i> , 2000 , 23, 1108-12	14.6	388

54	Segregation analysis of diabetic nephropathy in Pima Indians. <i>Diabetes</i> , 2000 , 49, 1049-56	0.9	74
53	Mutations in the genes for hepatocyte nuclear factor (HNF)-1alpha, -4alpha, -1beta, and -3beta; the dimerization cofactor of HNF-1; and insulin promoter factor 1 are not common causes of early-onset type 2 diabetes in Pima Indians. <i>Diabetes Care</i> , 2000 , 23, 302-4	14.6	13
52	Intrauterine exposure to diabetes conveys risks for type 2 diabetes and obesity: a study of discordant sibships. <i>Diabetes</i> , 2000 , 49, 2208-11	0.9	910
51	Evaluation of simple indices of insulin sensitivity and insulin secretion for use in epidemiologic studies. <i>American Journal of Epidemiology</i> , 2000 , 151, 190-8	3.8	379
50	Plasma glucose and prediction of microvascular disease and mortality: evaluation of 1997 American Diabetes Association and 1999 World Health Organization criteria for diagnosis of diabetes. <i>Diabetes Care</i> , 2000 , 23, 1113-8	14.6	157
49	A locus influencing total serum cholesterol on chromosome 19p: results from an autosomal genomic scan of serum lipid concentrations in Pima Indians. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2000 , 20, 2651-6	9.4	65
48	Secular trends in birth weight, BMI, and diabetes in the offspring of diabetic mothers. <i>Diabetes Care</i> , 2000 , 23, 1249-54	14.6	91
47	Individual estimates of European genetic admixture associated with lower body-mass index, plasma glucose, and prevalence of type 2 diabetes in Pima Indians. <i>American Journal of Human Genetics</i> , 2000 , 66, 527-38	11	95
46	HDL cholesterol subfractions and risk of developing type 2 diabetes among Pima Indians. <i>Diabetes Care</i> , 1999 , 22, 271-4	14.6	23
45	Birth weight, type 2 diabetes, and insulin resistance in Pima Indian children and young adults. <i>Diabetes Care</i> , 1999 , 22, 944-50	14.6	202
44	Evidence for genetic linkage to alcohol dependence on chromosomes 4 and 11 from an autosome-wide scan in an American Indian population. <i>American Journal of Medical Genetics Part A</i> , 1998 , 81, 216-21		303
43	Analytic strategies to detect linkage to a common disorder with genetically determined age of onset: diabetes mellitus in Pima Indians. <i>Genetic Epidemiology</i> , 1998 , 15, 299-315	2.6	16
42	Increasing prevalence of Type II diabetes in American Indian children. <i>Diabetologia</i> , 1998 , 41, 904-10	10.3	329
41	Autosomal genomic scan for loci linked to obesity and energy metabolism in Pima Indians. <i>American Journal of Human Genetics</i> , 1998 , 62, 659-68	11	172
40	An autosomal genomic scan for loci linked to type II diabetes mellitus and body-mass index in Pima Indians. <i>American Journal of Human Genetics</i> , 1998 , 63, 1130-8	11	432
39	Dietary calcium and blood pressure in a Native American population. <i>Journal of the American College of Nutrition</i> , 1998 , 17, 59-64	3.5	4
38	Association between uncoupling protein polymorphisms (UCP2-UCP3) and energy metabolism/obesity in Pima Indians. <i>Human Molecular Genetics</i> , 1998 , 7, 1431-5	5.6	230
37	The relationship of menstrual irregularity to type 2 diabetes in Pima Indian women. <i>Diabetes Care</i> , 1998 , 21, 346-9	14.6	40

36	Sib-pair linkage analysis for susceptibility genes for microvascular complications among Pima Indians with type 2 diabetes. Pima Diabetes Genes Group. <i>Diabetes</i> , 1998 , 47, 821-30	0.9	282
35	Variant in the regulatory subunit of phosphatidylinositol 3-kinase (p85alpha): preliminary evidence indicates a potential role of this variant in the acute insulin response and type 2 diabetes in Pima women. <i>Diabetes</i> , 1998 , 47, 973-5	0.9	35
34	An autosomal genomic scan for loci linked to prediabetic phenotypes in Pima Indians. <i>Journal of Clinical Investigation</i> , 1998 , 101, 1757-64	15.9	169
33	Plasma lipoproteins and incidence of non-insulin-dependent diabetes mellitus in Pima Indians: protective effect of HDL cholesterol in women. <i>Atherosclerosis</i> , 1997 , 128, 113-9	3.1	35
32	Breastfeeding and incidence of non-insulin-dependent diabetes mellitus in Pima Indians. <i>Lancet, The</i> , 1997 , 350, 166-8	40	253
31	Diagnosing diabetes mellitus--do we need new criteria?. <i>Diabetologia</i> , 1997 , 40, 247-55	10.3	71
30	Serum cholesterol and mortality rates in a Native American population with low cholesterol concentrations: a U-shaped association. <i>Circulation</i> , 1997 , 96, 1408-15	16.7	15
29	A two-step strategy for identification of high-risk subjects for a clinical trial of prevention of NIDDM. <i>Diabetes Care</i> , 1996 , 19, 972-8	14.6	20
28	Familial aggregation of medial arterial calcification in Pima Indians with and without diabetes. <i>Diabetes Care</i> , 1996 , 19, 968-71	14.6	9
27	Survival during renal replacement therapy for diabetic end-stage renal disease in Pima Indians. <i>Diabetes Care</i> , 1996 , 19, 1333-7	14.6	14
26	Incidence of diabetes mellitus in women following impaired glucose tolerance in pregnancy is lower than following impaired glucose tolerance in the non-pregnant state. <i>Diabetologia</i> , 1996 , 39, 1334-7	10.3	10
25	HLA*A2 confers mortality risk for cardiovascular disease in Pimans. <i>Tissue Antigens</i> , 1996 , 47, 188-93		5
24	Parental hypertension and proteinuria in Pima Indians with NIDDM. <i>Diabetologia</i> , 1996 , 39, 433-8	10.3	28
23	Early and late insulin response as predictors of NIDDM in Pima Indians with impaired glucose tolerance. <i>Diabetologia</i> , 1995 , 38, 187-92	10.3	25
22	Diabetic nephropathy: a risk factor for diabetes mellitus in offspring. <i>Diabetologia</i> , 1995 , 38, 221-6	10.3	35
21	Familial relationships between obesity and NIDDM. <i>Diabetes</i> , 1995 , 44, 418-22	0.9	23
20	Incidence and determinants of elevated urinary albumin excretion in Pima Indians with NIDDM. <i>Diabetes Care</i> , 1995 , 18, 182-7	14.6	85
19	Comparison of body size measurements as predictors of NIDDM in Pima Indians. <i>Diabetes Care</i> , 1995 , 18, 435-9	14.6	48

18	Rate of weight gain, weight fluctuation, and incidence of NIDDM. <i>Diabetes</i> , 1995 , 44, 261-6	0.9	77
17	The U-shaped association between body mass index and mortality: relationship with weight gain in a Native American population. <i>Journal of Clinical Epidemiology</i> , 1995 , 48, 903-16	5.7	55
16	Which test for diagnosing diabetes?. <i>Diabetes Care</i> , 1995 , 18, 1042-4	14.6	32
15	Rate of weight gain, weight fluctuation, and incidence of NIDDM. <i>Diabetes</i> , 1995 , 44, 261-266	0.9	20
14	Familial relationships between obesity and NIDDM. <i>Diabetes</i> , 1995 , 44, 418-422	0.9	6
13	Preventing non-insulin-dependent diabetes. <i>Diabetes</i> , 1995 , 44, 483-488	0.9	40
12	Familial and metabolic factors related to blood pressure in Pima Indian children. <i>American Journal of Epidemiology</i> , 1994 , 140, 123-31	3.8	32
11	Comparison of World Health Organization and National Diabetes Data Group procedures to detect abnormalities of glucose tolerance during pregnancy. <i>Diabetes Care</i> , 1994 , 17, 1264-8	14.6	60
10	Glucose, insulin concentrations and obesity in childhood and adolescence as predictors of NIDDM. <i>Diabetologia</i> , 1994 , 37, 617-23	10.3	92
9	Decreasing incidence and prevalence of rheumatoid arthritis in Pima Indians over a twenty-five-year period. <i>Arthritis and Rheumatism</i> , 1994 , 37, 1158-65		84
8	A cross-sectional and longitudinal comparison of the Rome criteria for active rheumatoid arthritis (equivalent to the American College of Rheumatology 1958 criteria) and the American College of Rheumatology 1987 criteria for rheumatoid arthritis. <i>Arthritis and Rheumatism</i> , 1994 , 37, 1479-86		30
7	Gravidity, obesity, and non-insulin-dependent diabetes among Pima Indian women. <i>American Journal of Medicine</i> , 1994 , 97, 250-5	2.4	25
6	Rheumatoid arthritis and mortality. A longitudinal study in Pima Indians. <i>Arthritis and Rheumatism</i> , 1993 , 36, 1045-53		139
5	Glycated haemoglobin, plasma glucose and diabetic retinopathy: cross-sectional and prospective analyses. <i>Diabetologia</i> , 1993 , 36, 428-32	10.3	53
4	Determinants of end-stage renal disease in Pima Indians with type 2 (non-insulin-dependent) diabetes mellitus and proteinuria. <i>Diabetologia</i> , 1993 , 36, 1087-93	10.3	71
3	Comparison of screening tests for non-insulin-dependent diabetes mellitus. <i>Archives of Internal Medicine</i> , 1993 , 153, 2133-2140		29
2	Evidence for a Role of LPGAT1 in Influencing BMI and Percent Body Fat in Native Americans. <i>Obesity</i> ,	8	2
1	Non-Caucasian North American Populations: Native Americans		181-194 1

