

Ænal GeÆgel

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4443537/publications.pdf>

Version: 2024-02-01

12
papers

544
citations

933447

10
h-index

1199594

12
g-index

13
all docs

13
docs citations

13
times ranked

669
citing authors

#	ARTICLE	IF	CITATIONS
1	Comparisons of activated carbons produced from sycamore balls, ripe black locust seed pods, and Nerium oleander fruits and also their H ₂ storage studies. <i>Carbon Letters</i> , 2021, 31, 75-92.	5.9	10
2	Adsorptive removal of diclofenac sodium from aqueous solution onto sycamore ball activated carbon " isotherms, kinetics, and thermodynamic study. <i>Surfaces and Interfaces</i> , 2021, 24, 101097.	3.0	20
3	Preparation and characterization of mesoporous activated carbons from waste watermelon rind by using the chemical activation method with zinc chloride. <i>Arabian Journal of Chemistry</i> , 2019, 12, 3621-3627.	4.9	56
4	Adsorptive Removal of Rhodamine B with Activated Carbon Obtained from Okra Wastes. <i>Chemical Engineering Communications</i> , 2017, 204, 772-783.	2.6	53
5	Adsorption of Methylene Blue by an Efficient Activated Carbon Prepared from Citrullus lanatus Rind: Kinetic, Isotherm, Thermodynamic, and Mechanism Analysis. <i>Water, Air, and Soil Pollution</i> , 2016, 227, 1.	2.4	98
6	Active carbon/graphene hydrogel nanocomposites as a symmetric device for supercapacitors. <i>Fullerenes Nanotubes and Carbon Nanostructures</i> , 2016, 24, 427-434.	2.1	14
7	Adsorption of cationic dyes on activated carbon obtained from waste <i>Elaeagnus</i> stone. <i>Adsorption Science and Technology</i> , 2016, 34, 512-525.	3.2	64
8	Adsorptive Removal of Methylene Blue from Aqueous Solution by the Activated Carbon Obtained from the Fruit of <i>Catalpa bignonioides</i> . <i>Water, Air, and Soil Pollution</i> , 2015, 226, 1.	2.4	22
9	Investigating Fatty Acid Composition of Samples were Homogenized Various Meat and Offal Products from Turkey. <i>JAOCS, Journal of the American Oil Chemists' Society</i> , 2015, 92, 659-665.	1.9	5
10	Removal of Methylene Blue from Aqueous Solution by Activated Carbon Prepared from Pea Shells (<i>Pisum sativum</i>). <i>Journal of Chemistry</i> , 2013, 2013, 1-9.	1.9	116
11	Adsorption of Remazol Brilliant Blue R on activated carbon prepared from a pine cone. <i>Natural Product Research</i> , 2012, 26, 659-664.	1.8	30
12	Adsorption behavior of Cr(VI) on activated hazelnut shell ash and activated bentonite. <i>Microporous and Mesoporous Materials</i> , 2006, 91, 107-110.	4.4	56