

Hyung Jung Oh

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4445737/hyung-jung-oh-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

78
papers

1,070
citations

18
h-index

29
g-index

82
ext. papers

1,305
ext. citations

3.9
avg. IF

3.88
L-index

#	Paper	IF	Citations
78	The dietary intake of chronic kidney disease according to stages: Findings from the Korean National Health and Nutritional Examination Survey. <i>PLoS ONE</i> , 2021 , 16, e0260242	3.7	
77	Metformin use and cardiovascular outcomes in patients with diabetes and chronic kidney disease: a nationwide cohort study.. <i>Kidney Research and Clinical Practice</i> , 2021 , 40, 660-672	3.6	1
76	Clinical outcomes among hemodialysis patients with atrial fibrillation: a Korean nationwide population-based study. <i>Kidney Research and Clinical Practice</i> , 2021 , 40, 99-108	3.6	1
75	Quantitative Analysis of Microperfusion in Contrast-Induced Nephropathy Using Contrast-Enhanced Ultrasound: An Animal Study. <i>Korean Journal of Radiology</i> , 2021 , 22, 801-810	6.9	1
74	Mortality predictors in critically ill patients with acute kidney injury requiring continuous renal replacement therapy. <i>Kidney Research and Clinical Practice</i> , 2021 , 40, 401-410	3.6	5
73	Is there association between statin usage and contrast-associated acute kidney injury after intravenous administration of iodine-based contrast media in enhanced computed tomography?. <i>European Radiology</i> , 2020 , 30, 5261-5271	8	3
72	Comparison of hemodialysis and peritoneal dialysis patients's dietary behaviors. <i>BMC Nephrology</i> , 2020 , 21, 91	2.7	5
71	The Combined Impact of Chronic Kidney Disease and Diabetes on the Risk of Colorectal Cancer Depends on Sex: A Nationwide Population-Based Study. <i>Yonsei Medical Journal</i> , 2020 , 61, 506-514	3	1
70	Plasma Fibroblast Growth Factor 23 Concentration Is Associated with Intracranial Cerebral Atherosclerosis in Acute Ischemic Stroke Patients. <i>Journal of Clinical Neurology (Korea)</i> , 2020 , 16, 29-36	1.7	4
69	Comparison of dietary intake patterns in hemodialysis patients by nutritional status: A cross-sectional analysis. <i>Kidney Research and Clinical Practice</i> , 2020 , 39, 202-212	3.6	2
68	Sex differences in the blood pressure level associated with increased risks of cardiovascular events: a Korean nationwide population-based cohort study. <i>Journal of Clinical Hypertension</i> , 2020 , 22, 1638-1646 ^{2,3}	4.3	2
67	Age and sex specific target of blood pressure for the prevention of cardiovascular event among the treatment naive hypertensive patients. <i>Scientific Reports</i> , 2020 , 10, 21538	4.9	4
66	Health Insurance Status Is Related to Risk of Mortality and Hospitalization in Korean Maintenance Hemodialysis Patients: A Longitudinal Cohort Study. <i>American Journal of Nephrology</i> , 2020 , 51, 975-981	4.6	0
65	The Effect of Aspirin on Preventing Vascular Access Dysfunction in Incident Hemodialysis Patients: A Prospective Cohort Study in Korean Clinical Research Centers for End-Stage Renal Disease (CRC for ESRD). <i>Journal of Clinical Medicine</i> , 2019 , 8,	5.1	3
64	The protective effect of klotho against contrast-associated acute kidney injury via the antioxidative effect. <i>American Journal of Physiology - Renal Physiology</i> , 2019 , 317, F881-F889	4.3	18
63	The interactive effects of input and output on managing fluid balance in patients with acute kidney injury requiring continuous renal replacement therapy. <i>Critical Care</i> , 2019 , 23, 329	10.8	4
62	Fracture risk in chronic kidney disease: A Korean population-based cohort study. <i>Kidney Research and Clinical Practice</i> , 2019 , 38, 220-228	3.6	13

61	Impact of chronic kidney disease on mortality: A nationwide cohort study. <i>Kidney Research and Clinical Practice</i> , 2019 , 38, 382-390	3.6	38
60	Effect of Renin-Angiotensin System Blockade on Mortality in Korean Hypertensive Patients with Proteinuria. <i>Electrolyte and Blood Pressure</i> , 2019 , 17, 25-35	1.1	1
59	Increased Thrombogenicity in Chronic Renal Failure in a Rat Model Induced by 5/6 Ablation/Infarction. <i>Yonsei Medical Journal</i> , 2018 , 59, 754-759	3	2
58	The impact of disease severity on paradoxical association between body mass index and mortality in patients with acute kidney injury undergoing continuous renal replacement therapy. <i>BMC Nephrology</i> , 2018 , 19, 32	2.7	7
57	Klotho plays a protective role against glomerular hypertrophy in a cell cycle-dependent manner in diabetic nephropathy. <i>American Journal of Physiology - Renal Physiology</i> , 2018 , 315, F791-F805	4.3	15
56	SP662 THE DISCREPANCY IN THE PREDICTABILITY OF SUBJECTIVE GLOBAL ASSESSMENT FOR MORTALITY ACCORDING TO DIALYSIS VINTAGE IN HEMODIALYSIS PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2018 , 33, i569-i569	4.3	
55	Incidence risk of various types of digestive cancers in patients with pre-dialytic chronic kidney disease: A nationwide population-based cohort study. <i>PLoS ONE</i> , 2018 , 13, e0207756	3.7	9
54	VolumE maNagement Under body composition monitoring in critically ill patientS on CRRT: study protocol for a randomized controlled trial (VENUS trial). <i>Trials</i> , 2018 , 19, 681	2.8	4
53	Gender-specific discrepancy in subjective global assessment for mortality in hemodialysis patients. <i>Scientific Reports</i> , 2018 , 8, 17846	4.9	6
52	Factors Associated with Early Mortality in Critically Ill Patients Following the Initiation of Continuous Renal Replacement Therapy. <i>Journal of Clinical Medicine</i> , 2018 , 7,	5.1	13
51	FP233THE PROTECTIVE EFFECT OF KLOTHO AGAINST CONTRAST ASSOCIATED ACUTE KIDNEY INJURY VIA THE ANTIOXIDATIVE EFFECT. <i>Nephrology Dialysis Transplantation</i> , 2018 , 33, i108-i108	4.3	
50	FP012EFFECT OF LONG-TERM TUBULAR OVEREXPRESSION OF HYPOXIA-INDUCIBLE FACTOR-2A ON THE PROGRESSIVE RENAL FIBROSIS IN CHRONIC KIDNEY DISEASE MODEL. <i>Nephrology Dialysis Transplantation</i> , 2018 , 33, i53-i53	4.3	1
49	FP113THE INCIDENCE RISK OF DIGESTIVE CANCER IN PATIENTS WITH PRE-DIALYTIC CHRONIC KIDNEY DISEASE: A NATIONWIDE COHORT STUDY IN KOREA. <i>Nephrology Dialysis Transplantation</i> , 2018 , 33, i15-i15	4.3	
48	Association of blood pressure components with mortality and cardiovascular events in prehypertensive individuals: a nationwide population-based cohort study. <i>Annals of Medicine</i> , 2018 , 50, 443-452	1.5	7
47	Prognostic Value of Residual Urine Volume, GFR by 24-hour Urine Collection, and eGFR in Patients Receiving Dialysis. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2017 , 12, 426-434	6.9	30
46	Characteristics and Clinical Outcomes of End-Stage Renal Disease Patients on Peritoneal Dialysis for Over 15 Years: A Single-Center Experience. <i>Peritoneal Dialysis International</i> , 2017 , 37, 535-541	2.8	4
45	Delta neutrophil index is an independent predictor of mortality in septic acute kidney injury patients treated with continuous renal replacement therapy. <i>BMC Nephrology</i> , 2017 , 18, 94	2.7	6
44	Blood Pressure Control During Chronic Kidney Disease Progression. <i>American Journal of Hypertension</i> , 2017 , 30, 610-616	2.3	11

43	Selective tubular activation of hypoxia-inducible factor-2 α has dual effects on renal fibrosis. <i>Scientific Reports</i> , 2017 , 7, 11351	4.9	19
42	Warfarin Use in Patients With Atrial Fibrillation Undergoing Hemodialysis: A Nationwide Population-Based Study. <i>Stroke</i> , 2017 , 48, 2472-2479	6.7	25
41	An increased chloride level in hypochloremia is associated with decreased mortality in patients with severe sepsis or septic shock. <i>Scientific Reports</i> , 2017 , 7, 15883	4.9	16
40	Dementia is a risk factor for major adverse cardiac and cerebrovascular events in elderly Korean patients initiating hemodialysis: a Korean national population-based study. <i>BMC Nephrology</i> , 2017 , 18, 128	2.7	4
39	Clinical usefulness of the Oxford classification in determining immunosuppressive treatment in IgA nephropathy. <i>Annals of Medicine</i> , 2017 , 49, 217-229	1.5	7
38	Baseline Chloride Levels are Associated with the Incidence of Contrast-Associated Acute Kidney Injury. <i>Scientific Reports</i> , 2017 , 7, 17431	4.9	10
37	Body mass index is inversely associated with mortality in patients with acute kidney injury undergoing continuous renal replacement therapy. <i>Kidney Research and Clinical Practice</i> , 2017 , 36, 39-47 ^{3.6}	3.6	14
36	The locally activated renin-angiotensin system is involved in albumin permeability in glomerular endothelial cells under high glucose conditions. <i>Nephrology Dialysis Transplantation</i> , 2017 , 32, 61-72	4.3	4
35	Early initiation of continuous renal replacement therapy improves survival of elderly patients with acute kidney injury: a multicenter prospective cohort study. <i>Critical Care</i> , 2016 , 20, 260	10.8	25
34	Association between Vitamin D Deficiency and Anemia in Patients with End-Stage Renal Disease: A Cross-Sectional Study. <i>Yonsei Medical Journal</i> , 2016 , 57, 1159-64	3	7
33	Can early initiation of continuous renal replacement therapy improve patient survival with septic acute kidney injury when enrolled in early goal-directed therapy?. <i>Journal of Critical Care</i> , 2016 , 35, 51-6 ⁴	4	14
32	Low serum intact parathyroid hormone level is an independent risk factor for overall mortality and major adverse cardiac and cerebrovascular events in incident dialysis patients. <i>Osteoporosis International</i> , 2016 , 27, 2717-2726	5.3	18
31	High-Dose Versus Conventional-Dose Continuous Venovenous Hemodiafiltration and Patient and Kidney Survival and Cytokine Removal in Sepsis-Associated Acute Kidney Injury: A Randomized Controlled Trial. <i>American Journal of Kidney Diseases</i> , 2016 , 68, 599-608	7.4	64
30	Glomerular IgG deposition predicts renal outcome in patients with IgA nephropathy. <i>Modern Pathology</i> , 2016 , 29, 743-52	9.8	25
29	Decreased circulating klotho levels in patients undergoing dialysis and relationship to oxidative stress and inflammation. <i>Peritoneal Dialysis International</i> , 2015 , 35, 43-51	2.8	40
28	An increase in mean platelet volume from baseline is associated with mortality in patients with severe sepsis or septic shock. <i>PLoS ONE</i> , 2015 , 10, e0119437	3.7	78
27	Irisin, a novel myokine is an independent predictor for sarcopenia and carotid atherosclerosis in dialysis patients. <i>Atherosclerosis</i> , 2015 , 242, 476-82	3.1	55
26	Can mean platelet volume predict the prognosis of patients with acute kidney injury requiring continuous renal replacement therapy? Response to comments. <i>Journal of Critical Care</i> , 2015 , 30, 208	4	

25	Rifampicin-induced minimal change disease is improved after cessation of rifampicin without steroid therapy. <i>Yonsei Medical Journal</i> , 2015 , 56, 582-5	3	7
24	The effect of specialized continuous renal replacement therapy team in acute kidney injury patients treatment. <i>Yonsei Medical Journal</i> , 2015 , 56, 658-65	3	14
23	Diastolic dysfunction is an independent predictor of cardiovascular events in incident dialysis patients with preserved systolic function. <i>PLoS ONE</i> , 2015 , 10, e0118694	3.7	29
22	Which Biomarker is the Best for Predicting Mortality in Incident Peritoneal Dialysis Patients: NT-ProBNP, Cardiac TnT, or hsCRP?: A Prospective Observational Study. <i>Medicine (United States)</i> , 2015 , 94, e1636	1.8	13
21	Double transduction of a Cre/LoxP lentiviral vector: a simple method to generate kidney cell-specific knockdown mice. <i>American Journal of Physiology - Renal Physiology</i> , 2015 , 309, F1060-9	4.3	9
20	Comparison of the Haas and the Oxford classifications for prediction of renal outcome in patients with IgA nephropathy. <i>Human Pathology</i> , 2014 , 45, 236-43	3.7	26
19	Mean platelet volume is a prognostic factor in patients with acute kidney injury requiring continuous renal replacement therapy. <i>Journal of Critical Care</i> , 2014 , 29, 1016-21	4	15
18	NT-proBNP: is it a more significant risk factor for mortality than troponin T in incident hemodialysis patients?. <i>Medicine (United States)</i> , 2014 , 93, e241	1.8	6
17	Optimal proteinuria target for renoprotection in patients with IgA nephropathy. <i>PLoS ONE</i> , 2014 , 9, e101935	3.7	19
16	The effect of renin-angiotensin system blockade on renal protection in chronic kidney disease patients with hyperkalemia. <i>JRAAS - Journal of the Renin-Angiotensin-Aldosterone System</i> , 2014 , 15, 491-7	3	9
15	Using the Oxford classification of IgA nephropathy to predict long-term outcomes of Henoch-Schönlein purpura nephritis in adults. <i>Modern Pathology</i> , 2014 , 27, 972-82	9.8	44
14	Insulin resistance is associated with new-onset cardiovascular events in nondiabetic patients undergoing peritoneal dialysis. <i>Kidney Research and Clinical Practice</i> , 2014 , 33, 192-8	3.6	7
13	The relationship of initial transferrin saturation to cardiovascular parameters and outcomes in patients initiating dialysis. <i>PLoS ONE</i> , 2014 , 9, e87231	3.7	11
12	Membranous glomerulonephritis in a patient with myelodysplastic syndrome-refractory cytopenia with multilineage dysplasia. <i>Kidney Research and Clinical Practice</i> , 2013 , 32, 134-7	3.6	5
11	Urine output is associated with prognosis in patients with acute kidney injury requiring continuous renal replacement therapy. <i>Journal of Critical Care</i> , 2013 , 28, 379-88	4	20
10	A case of membranous nephropathy as a manifestation of graft-versus-host disease. <i>Kidney Research and Clinical Practice</i> , 2013 , 32, 39-42	3.6	3
9	Minimal Change Disease in Systemic Lupus: Another Renal Manifestation of Lupus?. <i>The Ewha Medical Journal</i> , 2013 , 36, 139	0.1	1
8	Clinical Remission of Renal Amyloidosis after Autologous Peripheral Blood Stem Cell Transplantation. <i>The Ewha Medical Journal</i> , 2013 , 36, S25	0.1	

7	Cardiovascular disease in end-stage renal disease. <i>Journal of the Korean Medical Association</i> , 2013 , 56, 576	0.5	3
6	A low serum bicarbonate concentration as a risk factor for mortality in peritoneal dialysis patients. <i>PLoS ONE</i> , 2013 , 8, e82912	3.7	13
5	Early initiation of continuous renal replacement therapy improves patient survival in severe progressive septic acute kidney injury. <i>Journal of Critical Care</i> , 2012 , 27, 743.e9-18	4	27
4	Syndrome of inappropriate antidiuretic hormone secretion in a patient with large cell neuroendocrine carcinoma. <i>Yonsei Medical Journal</i> , 2012 , 53, 667-9	3	3
3	Red blood cell distribution width is an independent predictor of mortality in acute kidney injury patients treated with continuous renal replacement therapy. <i>Nephrology Dialysis Transplantation</i> , 2012 , 27, 589-94	4.3	104
2	Clinical outcomes, when matched at presentation, do not vary between adult-onset Henrich-Schleim purpura nephritis and IgA nephropathy. <i>Kidney International</i> , 2012 , 82, 1304-12	9.9	36
1	A Case of Bronchioloalveolar Carcinoma Presenting with Initial Symptom of Visual Disturbance due to Intraocular Metastasis. <i>Tuberculosis and Respiratory Diseases</i> , 2005 , 59, 93	3.2	