

Martin Espinosa-Bravo

List of Publications by Citations

Source: <https://exaly.com/author-pdf/4451354/martin-espinosa-bravo-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

47
papers

828
citations

14
h-index

28
g-index

62
ext. papers

1,296
ext. citations

4
avg, IF

5.45
L-index

#	Paper	IF	Citations
47	Timing of surgery following SARS-CoV-2 infection: an international prospective cohort study. <i>Anaesthesia</i> , 2021 , 76, 748-758	6.6	115
46	SARS-CoV-2 vaccination modelling for safe surgery to save lives: data from an international prospective cohort study. <i>British Journal of Surgery</i> , 2021 , 108, 1056-1063	5.3	77
45	The superparamagnetic iron oxide is equivalent to the Tc99 radiotracer method for identifying the sentinel lymph node in breast cancer. <i>European Journal of Surgical Oncology</i> , 2015 , 41, 46-51	3.6	73
44	Intraoperative molecular analysis of total tumor load in sentinel lymph node: a new predictor of axillary status in early breast cancer patients. <i>Breast Cancer Research and Treatment</i> , 2013 , 139, 87-93	4.4	72
43	Intraoperative Ultrasound-Guided Excision of Axillary Clip in Patients with Node-Positive Breast Cancer Treated with Neoadjuvant Therapy (ILINA Trial) : A New Tool to Guide the Excision of the Clipped Node After Neoadjuvant Treatment. <i>Annals of Surgical Oncology</i> , 2018 , 25, 784-791	3.1	67
42	Prediction of non-sentinel lymph node metastasis in early breast cancer by assessing total tumoral load in the sentinel lymph node by molecular assay. <i>European Journal of Surgical Oncology</i> , 2013 , 39, 766-73	3.6	40
41	Intraoperative Ultrasound-Guided Lumpectomy Versus Mammographic Wire Localization for Breast Cancer Patients After Neoadjuvant Treatment. <i>Annals of Surgical Oncology</i> , 2016 , 23, 38-43	3.1	38
40	Nomogram including the total tumoral load in the sentinel nodes assessed by one-step nucleic acid amplification as a new factor for predicting nonsentinel lymph node metastasis in breast cancer patients. <i>Breast Cancer Research and Treatment</i> , 2014 , 147, 371-80	4.4	33
39	Anxiety and depression in women with breast cancer: Social and clinical determinants and influence of the social network and social support (DAMA cohort). <i>Cancer Epidemiology</i> , 2018 , 55, 123-129	2.8	30
38	Global variation in postoperative mortality and complications after cancer surgery: a multicentre, prospective cohort study in 82 countries. <i>Lancet, The</i> , 2021 , 397, 387-397	4.0	29
37	Extensive nodal involvement increases the positivity of blue nodes in the axillary reverse mapping procedure in patients with breast cancer. <i>Journal of Surgical Oncology</i> , 2012 , 106, 89-93	2.8	24
36	Different Prognostic Implications of Residual Disease After Neoadjuvant Treatment: Impact of Ki 67 and Site of Response. <i>Annals of Surgical Oncology</i> , 2016 , 23, 3831-3837	3.1	21
35	Role of total tumour load of sentinel lymph node on survival in early breast cancer patients. <i>Breast</i> , 2017 , 33, 8-13	3.6	20
34	Electrothermal bipolar vessel sealing system in axillary dissection: a prospective randomized clinical study. <i>International Journal of Surgery</i> , 2011 , 9, 636-40	7.5	19
33	The Second Generation Antibody-Drug Conjugate SYD985 Overcomes Resistances to T-DM1. <i>Cancers</i> , 2020 , 12,	6.6	14
32	Surgery improves survival in elderly with breast cancer. A study of 465 patients in a single institution. <i>European Journal of Surgical Oncology</i> , 2015 , 41, 635-40	3.6	13
31	Breast conservative surgery after neoadjuvant chemotherapy in breast cancer patients: comparison of two tumor localization methods. <i>European Journal of Surgical Oncology</i> , 2011 , 37, 1038-43	3.6	13

30	SARS-CoV-2 infection and venous thromboembolism after surgery: an international prospective cohort study. <i>Anaesthesia</i> , 2022 , 77, 28-39	6.6	12
29	Social inequalities in quality of life in a cohort of women diagnosed with breast cancer in Barcelona (DAMA Cohort). <i>Cancer Epidemiology</i> , 2018 , 54, 38-47	2.8	11
28	Intraoperative assessment of sentinel lymph node by one-step nucleic acid amplification in breast cancer patients after neoadjuvant treatment reduces the need for a second surgery for axillary lymph node dissection. <i>Breast</i> , 2017 , 31, 40-45	3.6	11
27	Lymphatic mapping could not be impaired in the presence of breast carcinoma and coexisting small lymphocytic lymphoma. <i>American Journal of Case Reports</i> , 2013 , 14, 322-5	1.3	11
26	Surgery improves breast cancer-specific survival in octogenarians with early-stage breast cancer. <i>International Journal of Surgery</i> , 2013 , 11, 554-7	7.5	10
25	Detection of sentinel lymph node in breast cancer recurrence may change adjuvant treatment decision in patients with breast cancer recurrence and previous axillary surgery. <i>Breast</i> , 2014 , 23, 460-5	3.6	9
24	A randomized study comparing different doses of superparamagnetic iron oxide tracer for sentinel lymph node biopsy in breast cancer: The SUNRISE study. <i>European Journal of Surgical Oncology</i> , 2020 , 46, 2195-2201	3.6	7
23	Effects of pre-operative isolation on postoperative pulmonary complications after elective surgery: an international prospective cohort study. <i>Anaesthesia</i> , 2021 , 76, 1454-1464	6.6	7
22	Learning curves in intraoperative ultrasound guided surgery in breast cancer based on complete breast cancer excision and no need for second surgeries. <i>European Journal of Surgical Oncology</i> , 2019 , 45, 578-583	3.6	6
21	Percutaneous ultrasound-guided vacuum-assisted excision of benign breast lesions: A learning curve to assess outcomes. <i>British Journal of Radiology</i> , 2019 , 92, 20180626	3.4	6
20	SOLTI-1805 TOT-HER3 Study Concept: A Window-of-Opportunity Trial of Patritumab Deruxtecan, a HER3 Directed Antibody Drug Conjugate, in Patients With Early Breast Cancer. <i>Frontiers in Oncology</i> , 2021 , 11, 638482	5.3	5
19	Intraoperative ultrasound guided breast surgery: paving the way for personalized surgery. <i>Gland Surgery</i> , 2016 , 5, 366-8	2.2	5
18	Preclinical In Vivo Validation of the RAD51 Test for Identification of Homologous Recombination-Deficient Tumors and Patient Stratification.. <i>Cancer Research</i> , 2022 , 82, 1646-1657	10.1	4
17	Cancer-related fatigue and its determinants in a cohort of women with breast cancer: the DAMA Cohort. <i>Supportive Care in Cancer</i> , 2020 , 28, 5213-5221	3.9	3
16	Influence of Social Determinants, Lifestyle, Emotional Well-Being and the Use of Unconventional Therapies in Breast Cancer Progression in a Cohort of Women in Barcelona: Protocol for the DAMA Cohort. <i>JMIR Research Protocols</i> , 2017 , 6, e249	2	3
15	20 ERBB3 mRNA expression in breast cancer (BC): A SOLTI biomarker discovery analysis. <i>Annals of Oncology</i> , 2020 , 31, S15-S16	10.3	3
14	Leveraging the increased rates of pathologic complete response after neoadjuvant treatment in breast cancer to de-escalate surgical treatments. <i>Journal of Surgical Oncology</i> , 2021 , 123, 71-79	2.8	3
13	Her2 positive and triple negative breast cancer patients with clinically negative nodes at diagnosis and breast pathologic complete response may spare axillary surgery after neoadjuvant treatment. <i>European Journal of Cancer</i> , 2018 , 92, S21	7.5	2

12	Determinants of concordance in clinically relevant genes (CRG) from synchronously acquired tumor biopsies (tBx) and ctDNA in metastatic breast cancer (MBC).. <i>Journal of Clinical Oncology</i> , 2019 , 37, 1075-1075	2.2	2
11	Axillary staging based on molecular analysis: Results of the B-CLOSER-II study. <i>Pathology Research and Practice</i> , 2020 , 216, 153197	3.4	2
10	Prognostic value of ctDNA detection in patients with early breast cancer undergoing neoadjuvant therapy: A systematic review and meta-analysis.. <i>Cancer Treatment Reviews</i> , 2022 , 104, 102362	14.4	2
9	Breast cancer during pregnancy: matched study of diagnostic approach, tumor characteristics, and prognostic factors. <i>Tumori</i> , 2020 , 106, 378-387	1.7	1
8	Increased detection of sentinel nodes in breast cancer patients with the use of superparamagnetic iron oxide tracer.. <i>Journal of Clinical Oncology</i> , 2014 , 32, 100-100	2.2	1
7	Five microRNAs in Serum Are Able to Differentiate Breast Cancer Patients From Healthy Individuals. <i>Frontiers in Oncology</i> , 2020 , 10, 586268	5.3	1
6	Dietary changes and food habits: social and clinical determinants in a cohort of women diagnosed with breast cancer in Barcelona (DAMA cohort). <i>Cancer Causes and Control</i> , 2021 , 32, 1355-1364	2.8	0
5	Use of non-conventional therapies in a cohort of women with breast cancer (DAMA cohort) in Barcelona (Spain). A mixed-methods study. <i>European Journal of Integrative Medicine</i> , 2020 , 37, 101148	1.7	
4	Positive sentinel lymph node: the evolution of axillary surgery and intraoperative assessment of sentinel lymph nodes. <i>Breast Cancer Management</i> , 2014 , 3, 369-376	0.7	
3	Health-Related Quality of Life After Nipple-Sparing Mastectomy: Results From the INSPIRE Registry. <i>Annals of Surgical Oncology</i> , 2021 , 1	3.1	
2	Analysis of total tumor load of sentinel lymph node as a prognostic factor in patients with early breast cancer.. <i>Journal of Clinical Oncology</i> , 2016 , 34, 1042-1042	2.2	
1	Abstract PD7-09: Safety of conservative surgery with accelerated partial breast re-irradiation for isolated ipsilateral breast cancer recurrence regardless of immunohistochemical subtype. A multicentric prospective study. <i>Cancer Research</i> , 2022 , 82, PD7-09-PD7-09	10.1	