

Byung-Ho Kang

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4454176/byung-ho-kang-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

77
papers

3,164
citations

32
h-index

55
g-index

84
ext. papers

3,913
ext. citations

7.8
avg, IF

5.29
L-index

#	Paper	IF	Citations
77	Accelerated remodeling of the mesophyll-bundle sheath interface in the maize C4 cycle mutant leaves.. <i>Scientific Reports</i> , 2022 , 12, 5057	4.9	
76	A glossary of plant cell structures: Current insights and future questions. <i>Plant Cell</i> , 2021 ,	11.6	3
75	Chloroplast thylakoid ascorbate peroxidase PtotAPX plays a key role in chloroplast development by decreasing hydrogen peroxide in <i>Populus tomentosa</i> . <i>Journal of Experimental Botany</i> , 2021 , 72, 4333-4334	7.4	4
74	Friendly mediates membrane depolarization-induced mitophagy in Arabidopsis. <i>Current Biology</i> , 2021 , 31, 1931-1944.e4	6.3	13
73	The disassembly of lipid droplets in <i>Chlamydomonas</i> . <i>New Phytologist</i> , 2021 , 231, 1359-1364	9.8	4
72	Arabidopsis seedling establishment under waterlogging requires ABCG5-mediated formation of a dense cuticle layer. <i>New Phytologist</i> , 2021 , 229, 156-172	9.8	13
71	Three-dimensional reconstruction and comparison of vacuolar membranes in response to viral infection. <i>Journal of Integrative Plant Biology</i> , 2021 , 63, 353-364	8.3	5
70	Plant and animal chromatin three-dimensional organization: similar structures but different functions. <i>Journal of Experimental Botany</i> , 2020 , 71, 5119-5128	7	15
69	CrABCA2 Facilitates Triacylglycerol Accumulation in under Nitrogen Starvation. <i>Molecules and Cells</i> , 2020 , 43, 48-57	3.5	2
68	The Arabidopsis Protein Disulfide Isomerase Subfamily M Isoform, PDI9, Localizes to the Endoplasmic Reticulum and Influences Pollen Viability and Proper Formation of the Pollen Exine During Heat Stress. <i>Frontiers in Plant Science</i> , 2020 , 11, 610052	6.2	2
67	Electron Microscopy Views of Dimorphic Chloroplasts in C4 Plants. <i>Frontiers in Plant Science</i> , 2020 , 11, 1020	6.2	3
66	The phosphatidylethanolamine-binding protein DTH1 mediates degradation of lipid droplets in. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020 , 117, 23131-23139	11.5	8
65	Correlative Light and Electron Microscopy Imaging of the Plant trans-Golgi Network. <i>Methods in Molecular Biology</i> , 2020 , 2177, 59-67	1.4	3
64	Modular enzyme assembly for enhanced cascade biocatalysis and metabolic flux. <i>Nature Communications</i> , 2019 , 10, 4248	17.4	78
63	Electron tomography of plant organelles and the outlook for correlative microscopic approaches. <i>New Phytologist</i> , 2019 , 223, 1756-1761	9.8	10
62	Identification of Long Noncoding RNAs in the Developing Endosperm of Maize. <i>Methods in Molecular Biology</i> , 2019 , 1933, 49-65	1.4	1
61	Dietary fatty acids promote lipid droplet diversity through seipin enrichment in an ER subdomain. <i>Nature Communications</i> , 2019 , 10, 2902	17.4	32

60	Electron Tomography Analysis of Thylakoid Assembly and Fission in Chloroplasts of a Single-Cell C4 plant, <i>Bienertia sinuspersici</i> . <i>Scientific Reports</i> , 2019 , 9, 19640	4.9	4
59	A whole-cell electron tomography model of vacuole biogenesis in <i>Arabidopsis</i> root cells. <i>Nature Plants</i> , 2019 , 5, 95-105	11.5	49
58	Three-Dimensional Analysis of Chloroplast Structures Associated with Virus Infection. <i>Plant Physiology</i> , 2018 , 176, 282-294	6.6	33
57	The trans-Golgi sorting and the exocytosis of xylogalacturonan from the root border/border-like cell are conserved among monocot and dicot plant species. <i>Plant Signaling and Behavior</i> , 2018 , 13, e1469352	3.5	2
56	Thylakoid-Bound Polysomes and a Dynamin-Related Protein, FZL, Mediate Critical Stages of the Linear Chloroplast Biogenesis Program in Greening <i>Arabidopsis</i> Cotyledons. <i>Plant Cell</i> , 2018 , 30, 1476-1495	11.6	26
55	A Non-Classical Member of the Protein Disulfide Isomerase Family, PDI7 of <i>Arabidopsis thaliana</i> , Localizes to the cis-Golgi and Endoplasmic Reticulum Membranes. <i>Plant and Cell Physiology</i> , 2017 , 58, 1103-1117	4.9	6
54	Spatio-temporal analysis of coding and long noncoding transcripts during maize endosperm development. <i>Scientific Reports</i> , 2017 , 7, 3838	4.9	12
53	SH3 Domain-Containing Protein 2 Plays a Crucial Role at the Step of Membrane Tubulation during Cell Plate Formation. <i>Plant Cell</i> , 2017 , 29, 1388-1405	11.6	30
52	ATG9 regulates autophagosome progression from the endoplasmic reticulum in <i>Arabidopsis</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017 , 114, E426-E435	11.5	133
51	Semiautomatic Segmentation of Plant Golgi Stacks in Electron Tomograms Using 3dmod. <i>Methods in Molecular Biology</i> , 2017 , 1662, 97-104	1.4	8
50	3D Printing of Plant Golgi Stacks from Their Electron Tomographic Models. <i>Methods in Molecular Biology</i> , 2017 , 1662, 105-113	1.4	
49	A distinct class of vesicles derived from the trans-Golgi mediates secretion of xylogalacturonan in the root border cell. <i>Plant Journal</i> , 2017 , 92, 596-610	6.9	36
48	Nuclear Pore Permeabilization Is a Convergent Signaling Event in Effector-Triggered Immunity. <i>Cell</i> , 2016 , 166, 1526-1538.e11	56.2	77
47	STEM Tomography Imaging of Hypertrophied Golgi Stacks in Mucilage-Secreting Cells. <i>Methods in Molecular Biology</i> , 2016 , 1496, 55-62	1.4	7
46	Kinetics and specificity of paternal mitochondrial elimination in <i>Caenorhabditis elegans</i> . <i>Nature Communications</i> , 2016 , 7, 12569	17.4	36
45	Postmeiotic development of pollen surface layers requires two <i>Arabidopsis</i> ABCG-type transporters. <i>Plant Cell Reports</i> , 2016 , 35, 1863-73	5.1	22
44	Mitochondrial endonuclease G mediates breakdown of paternal mitochondria upon fertilization. <i>Science</i> , 2016 , 353, 394-9	33.3	107
43	<i>Arabidopsis</i> protein disulfide isomerase-8 is a type I endoplasmic reticulum transmembrane protein with thiol-disulfide oxidase activity. <i>BMC Plant Biology</i> , 2016 , 16, 181	5.3	5

42	Acute heart failure with cardiomyocyte atrophy induced in adult mice by ablation of cardiac myosin light chain kinase. <i>Cardiovascular Research</i> , 2016 , 111, 34-43	9.9	23
41	Unconventional Protein Secretion in Plants. <i>Methods in Molecular Biology</i> , 2016 , 1459, 47-63	1.4	14
40	Characterization of a <i>Chlamydomonas reinhardtii</i> mutant defective in a maltose transporter 2015 , 58, 344-351		5
39	Conserved Functions of the MATE Transporter BIG EMBRYO1 in Regulation of Lateral Organ Size and Initiation Rate. <i>Plant Cell</i> , 2015 , 27, 2288-300	11.6	43
38	C2-O-02 Dimorphic secretory vesicles produced from the Golgi stacks of mucilage secreting root cap cells. <i>Microscopy (Oxford, England)</i> , 2015 , 64, i65.1-i65	1.3	
37	Shared elements of host-targeting pathways among apicomplexan parasites of differing lifestyles. <i>Cellular Microbiology</i> , 2015 , 17, 1618-39	3.9	25
36	AtPGL3 is an Arabidopsis BURP domain protein that is localized to the cell wall and promotes cell enlargement. <i>Frontiers in Plant Science</i> , 2015 , 6, 412	6.2	16
35	Auxin-callose-mediated plasmodesmal gating is essential for tropic auxin gradient formation and signaling. <i>Developmental Cell</i> , 2014 , 28, 132-46	10.2	113
34	Adaptive expansion of the maize maternally expressed gene (Meg) family involves changes in expression patterns and protein secondary structures of its members. <i>BMC Plant Biology</i> , 2014 , 14, 204	5.3	12
33	Retention mechanisms for ER and Golgi membrane proteins. <i>Trends in Plant Science</i> , 2014 , 19, 508-15	13.1	65
32	High-pressure freezing and low-temperature processing of plant tissue samples for electron microscopy. <i>Methods in Molecular Biology</i> , 2014 , 1080, 147-57	1.4	16
31	Reconstructing plant cells in 3D by serial section electron tomography. <i>Methods in Molecular Biology</i> , 2014 , 1080, 159-70	1.4	18
30	Defective chloroplast development inhibits maintenance of normal levels of abscisic acid in a mutant of the Arabidopsis RH3 DEAD-box protein during early post-germination growth. <i>Plant Journal</i> , 2013 , 73, 720-32	6.9	39
29	Cis-Golgi cisternal assembly and biosynthetic activation occur sequentially in plants and algae. <i>Traffic</i> , 2013 , 14, 551-67	5.7	64
28	Vaccinia virions deficient in transcription enzymes lack a nucleocapsid. <i>Virology</i> , 2012 , 434, 50-8	3.6	10
27	CED-1, CED-7, and TTR-52 regulate surface phosphatidylserine expression on apoptotic and phagocytic cells. <i>Current Biology</i> , 2012 , 22, 1267-75	6.3	60
26	Callose deposition in the phloem plasmodesmata and inhibition of phloem transport in citrus leaves infected with "Candidatus Liberibacter asiaticus". <i>Protoplasma</i> , 2012 , 249, 687-97	3.4	107
25	Overexpression of Arabidopsis plasmodesmata germin-like proteins disrupts root growth and development. <i>Plant Cell</i> , 2012 , 24, 3630-48	11.6	59

24	Functional characterization of Arabidopsis thaliana isopropylmalate dehydrogenases reveals their important roles in gametophyte development. <i>New Phytologist</i> , 2011 , 189, 160-75	9.8	26
23	Electron tomography of RabA4b- and PI-4K α -labeled trans Golgi network compartments in Arabidopsis. <i>Traffic</i> , 2011 , 12, 313-29	5.7	196
22	Discovery of Genes Expressed in Basal Endosperm Transfer Cells in Maize Using 454 Transcriptome Sequencing. <i>Plant Molecular Biology Reporter</i> , 2011 , 29, 835-847	1.7	20
21	Protein disulfide isomerase-2 of Arabidopsis mediates protein folding and localizes to both the secretory pathway and nucleus, where it interacts with maternal effect embryo arrest factor. <i>Molecules and Cells</i> , 2011 , 32, 459-75	3.5	31
20	<i>Ta. Liberibacter asiaticus</i> carries an excision plasmid prophage and a chromosomally integrated prophage that becomes lytic in plant infections. <i>Molecular Plant-Microbe Interactions</i> , 2011 , 24, 458-68	3.6	87
19	Shrinkage and fragmentation of the trans-Golgi network in non-meristematic plant cells. <i>Plant Signaling and Behavior</i> , 2011 , 6, 884-6	2.5	11
18	Leishmania parasitophorous vacuoles interact continuously with the host cell's endoplasmic reticulum; parasitophorous vacuoles are hybrid compartments. <i>Cellular Microbiology</i> , 2010 , 12, 1480-94	3.9	51
17	Auxin-mediated ribosomal biogenesis regulates vacuolar trafficking in Arabidopsis. <i>Plant Cell</i> , 2010 , 22, 143-58	11.6	71
16	Electron microscopy and high-pressure freezing of Arabidopsis. <i>Methods in Cell Biology</i> , 2010 , 96, 259-83	1.8	55
15	Miniature1-encoded cell wall invertase is essential for assembly and function of wall-in-growth in the maize endosperm transfer cell. <i>Plant Physiology</i> , 2009 , 151, 1366-76	6.6	66
14	Statolith sedimentation kinetics and force transduction to the cortical endoplasmic reticulum in gravity-sensing Arabidopsis columella cells. <i>Plant Cell</i> , 2009 , 21, 843-60	11.6	115
13	Bcl-2 proteins EGL-1 and CED-9 do not regulate mitochondrial fission or fusion in Caenorhabditis elegans. <i>Current Biology</i> , 2009 , 19, 768-73	6.3	23
12	Caenorhabditis elegans drp-1 and fis-2 regulate distinct cell-death execution pathways downstream of ced-3 and independent of ced-9. <i>Molecular Cell</i> , 2008 , 31, 586-597	17.6	113
11	Nanoscale architecture of endoplasmic reticulum export sites and of Golgi membranes as determined by electron tomography. <i>Plant Physiology</i> , 2008 , 147, 1454-68	6.6	142
10	Electron microscopy analysis of maize basal endosperm transfer cells processed by high-pressure freezing and freeze-substitution. <i>Microscopy and Microanalysis</i> , 2008 , 14, 1502-1503	0.5	
9	ER-to-Golgi transport by COPII vesicles in Arabidopsis involves a ribosome-excluding scaffold that is transferred with the vesicles to the Golgi matrix. <i>Protoplasma</i> , 2008 , 234, 51-64	3.4	80
8	The cyclic nucleotide gated cation channel AtCNGC10 traffics from the ER via Golgi vesicles to the plasma membrane of Arabidopsis root and leaf cells. <i>BMC Plant Biology</i> , 2007 , 7, 48	5.3	48
7	Identification and characterization of COPIa- and COPIb-type vesicle classes associated with plant and algal Golgi. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 163-8	11.5	106

6	Members of the Arabidopsis dynamin-like gene family, ADL1, are essential for plant cytokinesis and polarized cell growth. <i>Plant Cell</i> , 2003 , 15, 899-913	11.6	139
5	The dynamin-like protein ADL1C is essential for plasma membrane maintenance during pollen maturation. <i>Plant Journal</i> , 2003 , 35, 1-15	6.9	78
4	Three-Dimensional Analysis of Syncytial-Type Cell Plates during Endosperm Cellularization Visualized by High Resolution Electron Tomography. <i>Plant Cell</i> , 2001 , 13, 2033	11.6	
3	The arabidopsis cell plate-associated dynamin-like protein, ADL1Ap, is required for multiple stages of plant growth and development. <i>Plant Physiology</i> , 2001 , 126, 47-68	6.6	96
2	Three-Dimensional Analysis of Syncytial-Type Cell Plates during Endosperm Cellularization Visualized by High Resolution Electron Tomography. <i>Plant Cell</i> , 2001 , 13, 2033-2051	11.6	131
1	Friendly regulates membrane depolarization induced mitophagy in Arabidopsis		1