

Giovanni Targher

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4466641/giovanni-targher-publications-by-year.pdf>

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

488
papers

30,756
citations

85
h-index

163
g-index

543
ext. papers

37,774
ext. citations

6.6
avg, IF

7.82
L-index

#	Paper	IF	Citations
488	Non-alcoholic fatty liver disease and risk of incident chronic kidney disease: an updated meta-analysis. <i>Gut</i> , 2022 , 71, 156-162	19.2	56
487	Association of metabolic dysfunction-associated fatty liver disease with kidney disease.. <i>Nature Reviews Nephrology</i> , 2022 ,	14.9	5
486	Efficacy of peroxisome proliferator-activated receptor agonists, glucagon-like peptide-1 receptor agonists, or sodium-glucose cotransporter-2 inhibitors for treatment of non-alcoholic fatty liver disease: a systematic review.. <i>The Lancet Gastroenterology and Hepatology</i> , 2022 ,	18.8	6
485	J-shaped relationship between serum zinc levels and the severity of hepatic necro-inflammation in patients with MAFLD.. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2022 ,	4.5	2
484	Risk of non-alcoholic fatty liver disease in patients with chronic plaque psoriasis: an updated systematic review and meta-analysis of observational studies.. <i>Journal of Endocrinological Investigation</i> , 2022 , 1	5.2	3
483	Association between higher serum uric acid levels and plasma N-terminal pro-B-type natriuretic peptide concentrations in patients with coronary artery disease and without overt heart failure.. <i>International Journal of Cardiology</i> , 2022 ,	3.2	2
482	Metabolic Dysfunction-associated Fatty Liver Disease is Associated with Greater Impairment of Lung Function than Nonalcoholic Fatty Liver Disease.. <i>Journal of Clinical and Translational Hepatology</i> , 2022 , 10, 230-237	5.2	2
481	Risk of Heart Failure in Patients With Nonalcoholic Fatty Liver Disease: JACC Review Topic of the Week.. <i>Journal of the American College of Cardiology</i> , 2022 , 79, 180-191	15.1	3
480	A novel radiomics signature based on T2-weighted imaging accurately predicts hepatic inflammation in individuals with biopsy-proven nonalcoholic fatty liver disease: a derivation and independent validation study.. <i>Hepatobiliary Surgery and Nutrition</i> , 2022 , 11, 212-226	2.1	0
479	Interaction of , and Increases Susceptibility to Nonalcoholic Steatohepatitis.. <i>Journal of Clinical and Translational Hepatology</i> , 2022 , 10, 219-229	5.2	0
478	Effects of pioglitazone on cardiovascular events and all-cause mortality in patients with type 2 diabetes: A meta-analysis of randomized controlled trials.. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2022 , 32, 529-536	4.5	1
477	Italian guidelines for the treatment of type 2 diabetes.. <i>Acta Diabetologica</i> , 2022 , 59, 579	3.9	0
476	Potential Blood DNA Methylation Biomarker Genes for Diagnosis of Liver Fibrosis in Patients With Biopsy-Proven Non-alcoholic Fatty Liver Disease.. <i>Frontiers in Medicine</i> , 2022 , 9, 864570	4.9	0
475	Global multi-stakeholder endorsement of the MAFLD definition.. <i>The Lancet Gastroenterology and Hepatology</i> , 2022 ,	18.8	18
474	Lifestyle Interventions for Non-Obese Patients Both with, and at Risk, of Non-Alcoholic Fatty Liver Disease. <i>Diabetes and Metabolism Journal</i> , 2022 , 46, 391-401	5	0
473	Elastographic parameters of liver steatosis and fibrosis predict independently the risk of incident chronic kidney disease and acute myocardial infarction in patients with type 2 diabetes mellitus. <i>Journal of Diabetes and Its Complications</i> , 2022 , 108226	3.2	0
472	Associations of liver volume and other markers of hepatic steatosis with all-cause mortality in the general population. <i>Liver International</i> , 2021 ,	7.9	2

471	The EASL-Lancet Liver Commission: protecting the next generation of Europeans against liver disease complications and premature mortality. <i>Lancet, The</i> , 2021 ,	4.0	12
470	Among simple non-invasive scores, Pro-C3 and ADAPT best exclude advanced fibrosis in Asian patients with MAFLD.. <i>Metabolism: Clinical and Experimental</i> , 2021 , 128, 154958	12.7	0
469	Growth differentiation factor-15 and the association between type 2 diabetes and liver fibrosis in NAFLD. <i>Nutrition and Diabetes</i> , 2021 , 11, 32	4.7	3
468	PNPLA3 rs738409 C>G Variant Influences the Association Between Visceral Fat and Significant Fibrosis in Biopsy-proven Nonalcoholic Fatty Liver Disease. <i>Journal of Clinical and Translational Hepatology</i> , 2021 , 000, 000-000	5.2	
467	Prognostic Role of Pericardial Fat on the Incidence of Heart Failure. <i>Journal of the American College of Cardiology</i> , 2021 , 78, e111	15.1	
466	Optimal thresholds for ultrasound attenuation parameter in the evaluation of hepatic steatosis severity: evidence from a cohort of patients with biopsy-proven fatty liver disease. <i>European Journal of Gastroenterology and Hepatology</i> , 2021 , 33, 430-435	2.2	3
465	polymorphism influences the association between sarcopenia and liver fibrosis in adults with biopsy-proven non-alcoholic fatty liver disease. <i>British Journal of Nutrition</i> , 2021 , 126, 813-824	3.6	5
464	Association and Interaction Between Serum Interleukin-6 Levels and Metabolic Dysfunction-Associated Fatty Liver Disease in Patients With Severe Coronavirus Disease 2019. <i>Frontiers in Endocrinology</i> , 2021 , 12, 604100	5.7	10
463	Non-alcoholic fatty liver disease and increased risk of incident extrahepatic cancers: a meta-analysis of observational cohort studies. <i>Gut</i> , 2021 ,	19.2	19
462	Individualized Polygenic Risk Score Identifies NASH in the Eastern Asia Region: A Derivation and Validation Study. <i>Clinical and Translational Gastroenterology</i> , 2021 , 12, e00321	4.2	0
461	TA allele of rs2070673 in the CYP2E1 gene is associated with lobular inflammation and nonalcoholic steatohepatitis in patients with biopsy-proven nonalcoholic fatty liver disease. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2021 , 36, 2925-2934	4	1
460	Machine learning algorithm outperforms fibrosis markers in predicting significant fibrosis in biopsy-confirmed NAFLD. <i>Journal of Hepato-Biliary-Pancreatic Sciences</i> , 2021 , 28, 593-603	2.8	4
459	Association between MBOAT7 rs641738 polymorphism and non-alcoholic fatty liver in overweight or obese children. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021 , 31, 1548-1555	4.5	3
458	Non-alcoholic fatty liver disease in obese children and adolescents: a role for nutrition?. <i>European Journal of Clinical Nutrition</i> , 2021 ,	5.2	7
457	The complex link between NAFLD and type 2 diabetes mellitus - mechanisms and treatments. <i>Nature Reviews Gastroenterology and Hepatology</i> , 2021 , 18, 599-612	24.2	55
456	Sex influences the association between appendicular skeletal muscle mass to visceral fat area ratio and non-alcoholic steatohepatitis in patients with biopsy-proven non-alcoholic fatty liver disease. <i>British Journal of Nutrition</i> , 2021 , 1-8	3.6	4
455	Biological disease-modifying antirheumatic drugs may mitigate the risk of psoriatic arthritis in patients with chronic plaque psoriasis. <i>Annals of the Rheumatic Diseases</i> , 2021 ,	2.4	8
454	The HSD17B13 rs72613567 variant is associated with lower levels of albuminuria in patients with biopsy-proven nonalcoholic fatty liver disease. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021 , 31, 1822-1831	4.5	2

453	Significant liver fibrosis, as assessed by fibroscan, is independently associated with chronic vascular complications of type 2 diabetes: A multicenter study. <i>Diabetes Research and Clinical Practice</i> , 2021 , 177, 108884	7.4	3
452	Plasma Bile Acid Profile in Patients with and without Type 2 Diabetes. <i>Metabolites</i> , 2021 , 11,	5.6	3
451	Non-alcoholic fatty liver disease: a multisystem disease requiring a multidisciplinary and holistic approach. <i>The Lancet Gastroenterology and Hepatology</i> , 2021 , 6, 578-588	18.8	37
450	Association between increased plasma ceramides and chronic kidney disease in patients with and without ischemic heart disease. <i>Diabetes and Metabolism</i> , 2021 , 47, 101152	5.4	9
449	Extrapulmonary complications of COVID-19: A multisystem disease?. <i>Journal of Medical Virology</i> , 2021 , 93, 323-335	19.7	63
448	Association between lower plasma adiponectin levels and higher plasma thrombin generation parameters in men with type 2 diabetes: role of plasma triglycerides. <i>Journal of Endocrinological Investigation</i> , 2021 , 44, 547-555	5.2	2
447	NAFLD-related mortality: simple hepatic steatosis is not as 'benign' as thought. <i>Gut</i> , 2021 , 70, 1212-1213	19.2	8
446	Association between positivity of serum autoantibodies and liver disease severity in patients with biopsy-proven NAFLD. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021 , 31, 552-560	4.5	2
445	Diffuse Idiopathic Skeletal Hyperostosis (DISH) in Type 2 Diabetes: A New Imaging Possibility and a New Biomarker. <i>Calcified Tissue International</i> , 2021 , 108, 231-239	3.9	6
444	MAFLD and risk of CKD. <i>Metabolism: Clinical and Experimental</i> , 2021 , 115, 154433	12.7	55
443	NAFLD, and cardiovascular and cardiac diseases: Factors influencing risk, prediction and treatment. <i>Diabetes and Metabolism</i> , 2021 , 47, 101215	5.4	29
442	Effect of metformin on all-cause mortality and major adverse cardiovascular events: An updated meta-analysis of randomized controlled trials. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021 , 31, 699-704	4.5	11
441	Non-alcoholic fatty liver disease and risk of incident diabetes mellitus: an updated meta-analysis of 501 022 adult individuals. <i>Gut</i> , 2021 , 70, 962-969	19.2	80
440	Glucagon-Like Peptide-1 Receptor Agonists for Treatment of Nonalcoholic Fatty Liver Disease and Nonalcoholic Steatohepatitis: An Updated Meta-Analysis of Randomized Controlled Trials. <i>Metabolites</i> , 2021 , 11,	5.6	50
439	Liver Fibrosis Biomarkers Accurately Exclude Advanced Fibrosis and Are Associated with Higher Cardiovascular Risk Scores in Patients with NAFLD or Viral Chronic Liver Disease. <i>Diagnostics</i> , 2021 , 11,	3.8	22
438	Prevalence of hepatic steatosis in patients with type 2 diabetes and response to glucose-lowering treatments. A multicenter retrospective study in Italian specialist care. <i>Journal of Endocrinological Investigation</i> , 2021 , 44, 1879-1889	5.2	5
437	Nonalcoholic Fatty Liver Disease and Cardiovascular Disease. <i>Clinical Liver Disease</i> , 2021 , 17, 19-22	2.2	13
436	Associations of Hydroxysteroid 17-beta Dehydrogenase 13 Variants with Liver Histology in Chinese Patients with Metabolic-associated Fatty Liver Disease. <i>Journal of Clinical and Translational Hepatology</i> , 2021 , 9, 194-202	5.2	1

435	Is COVID-19 lockdown associated with vitamin D deficiency?. <i>European Journal of Public Health</i> , 2021 , 31, 278-279	2.1	6
434	Links between metabolic syndrome and metabolic dysfunction-associated fatty liver disease. <i>Trends in Endocrinology and Metabolism</i> , 2021 , 32, 500-514	8.8	24
433	Beneficial effects of glucagon-like peptide 1 receptor agonists on glucose control, cardiovascular risk profile, and non-alcoholic fatty liver disease. An expert opinion of the Italian diabetes society. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021 , 31, 3257-3270	4.5	4
432	Non-alcoholic fatty liver disease-related risk of cardiovascular disease and other cardiac complications. <i>Diabetes, Obesity and Metabolism</i> , 2021 ,	6.7	7
431	Improvement of glycemic control in type 2 diabetes: A systematic review and meta-analysis of randomized controlled trials. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021 , 31, 2539-2546	4.5	4
430	Association between non-alcoholic fatty liver disease and impaired cardiac sympathetic/parasympathetic balance in subjects with and without type 2 diabetes-The Cooperative Health Research in South Tyrol (CHRIS)-NAFLD sub-study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021 , 31, 3414-3420	4.5	3
429	Incorporating fatty liver disease in multidisciplinary care and novel clinical trial designs for patients with metabolic diseases. <i>The Lancet Gastroenterology and Hepatology</i> , 2021 , 6, 743-753	18.8	14
428	Coffee, Atrial Fibrillation, and Circulating Ceramides in Patients with Chronic Heart Failure. <i>Journal of Agricultural and Food Chemistry</i> , 2021 , 69, 11236-11245	5.7	1
427	Non-alcoholic fatty liver disease is a risk factor for cardiovascular and cardiac diseases: further evidence that a holistic approach to treatment is needed. <i>Gut</i> , 2021 ,	19.2	1
426	Non-alcoholic fatty liver disease and risk of fatal and non-fatal cardiovascular events: an updated systematic review and meta-analysis. <i>The Lancet Gastroenterology and Hepatology</i> , 2021 , 6, 903-913	18.8	28
425	Changes in markers of hepatic steatosis and fibrosis in patients with type 2 diabetes during treatment with glucagon-like peptide-1 receptor agonists. A multicenter retrospective longitudinal study. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2021 , 31, 3474-3483	4.5	3
424	acNASH index to diagnose nonalcoholic steatohepatitis: a prospective derivation and global validation study. <i>EClinicalMedicine</i> , 2021 , 41, 101145	11.3	2
423	Non-Alcoholic Fatty Liver Disease Is Associated With Reduced Glomerular Filtration Rate in Patients With Chronic Plaque Psoriasis.. <i>Journal of Cutaneous Medicine and Surgery</i> , 2021 , 12034754211066906	1.6	0
422	Pre-existing type 2 diabetes is associated with increased all-cause death independently of echocardiographic predictors of poor prognosis only in ischemic heart disease. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020 , 30, 2036-2040	4.5	1
421	Risk of severe illness from COVID-19 in patients with metabolic dysfunction-associated fatty liver disease and increased fibrosis scores. <i>Gut</i> , 2020 , 69, 1545-1547	19.2	93
420	Patients with diabetes are at higher risk for severe illness from COVID-19. <i>Diabetes and Metabolism</i> , 2020 , 46, 335-337	5.4	84
419	Obesity Is a Risk Factor for Greater COVID-19 Severity. <i>Diabetes Care</i> , 2020 , 43, e72-e74	14.6	199
418	Diabetes as a risk factor for greater COVID-19 severity and in-hospital death: A meta-analysis of observational studies. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020 , 30, 1236-1248	4.5	126

417	Effect of insulin secretagogues on major cardiovascular events and all-cause mortality: A meta-analysis of randomized controlled trials. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020 , 30, 1601-1608	4.5	12
416	Detrimental effects of metabolic dysfunction-associated fatty liver disease and increased neutrophil-to-lymphocyte ratio on severity of COVID-19. <i>Diabetes and Metabolism</i> , 2020 , 46, 505-507	5.4	19
415	PNPLA3 I148M gene variant and chronic kidney disease in type 2 diabetic patients with NAFLD: Clinical and experimental findings. <i>Liver International</i> , 2020 , 40, 1130-1141	7.9	16
414	Global epidemiology of lean non-alcoholic fatty liver disease: A systematic review and meta-analysis. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2020 , 35, 2041-2050	4	24
413	Nonalcoholic Fatty Liver Disease and Implications for Older Adults with Diabetes. <i>Clinics in Geriatric Medicine</i> , 2020 , 36, 527-547	3.8	3
412	Lower levels of plasma NT-proBNP are associated with higher prevalence of NASH in patients with biopsy-proven NAFLD. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020 , 30, 1820-1825	4.5	5
411	Relation between plasma ceramides and cardiovascular death in chronic heart failure: A subset analysis of the GISSI-HF trial. <i>ESC Heart Failure</i> , 2020 , 7, 3288	3.7	6
410	PNPLA3 polymorphism influences the association between high-normal TSH level and NASH in euthyroid adults with biopsy-proven NAFLD. <i>Diabetes and Metabolism</i> , 2020 , 46, 496-503	5.4	3
409	Further advices on measuring lipoprotein(a) for reducing the residual cardiovascular risk on statin therapy. <i>Clinical Chemistry and Laboratory Medicine</i> , 2020 , 58, e144-e147	5.9	1
408	Synbiotics Alter Fecal Microbiomes, But Not Liver Fat or Fibrosis, in a Randomized Trial of Patients With Nonalcoholic Fatty Liver Disease. <i>Gastroenterology</i> , 2020 , 158, 1597-1610.e7	13.3	59
407	Is Nonalcoholic Fatty Liver Disease Not a Risk Factor for Cardiovascular Disease: Not Yet Time for a Change of Heart. <i>Hepatology</i> , 2020 , 71, 1867-1869	11.2	11
406	Complications, morbidity and mortality of nonalcoholic fatty liver disease. <i>Metabolism: Clinical and Experimental</i> , 2020 , 111S, 154170	12.7	113
405	Epidemiology and pathophysiology of the association between NAFLD and metabolically healthy or metabolically unhealthy obesity. <i>Annals of Hepatology</i> , 2020 , 19, 359-366	3.1	33
404	A new definition for metabolic dysfunction-associated fatty liver disease: An international expert consensus statement. <i>Journal of Hepatology</i> , 2020 , 73, 202-209	13.4	764
403	Development and validation of a novel non-invasive test for diagnosing fibrotic non-alcoholic steatohepatitis in patients with biopsy-proven non-alcoholic fatty liver disease. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2020 , 35, 1804-1812	4	4
402	Combined and sequential non-invasive approach to diagnosing non-alcoholic steatohepatitis in patients with non-alcoholic fatty liver disease and persistently normal alanine aminotransferase levels. <i>BMJ Open Diabetes Research and Care</i> , 2020 , 8,	4.5	11
401	COVID-19 and Liver Dysfunction: Current Insights and Emergent Therapeutic Strategies. <i>Journal of Clinical and Translational Hepatology</i> , 2020 , 8, 18-24	5.2	231
400	NAFLD, Diabetes, and Other Endocrine Diseases: Clinical Implications 2020 , 147-168		

399	Sodium-Glucose Cotransporter-2 Inhibitors for Treatment of Nonalcoholic Fatty Liver Disease: A Meta-Analysis of Randomized Controlled Trials. <i>Metabolites</i> , 2020 , 11,	5.6	22
398	Diabetes and NAFLD. <i>Endocrinology</i> , 2020 , 495-521	0.1	
397	GLP-1 receptor agonists for NAFLD treatment in patients with and without type 2 diabetes: an updated meta-analysis. <i>Exploration of Medicine</i> , 2020 , 1, 108-123	1.1	2
396	Echocardiographic parameters according to insulin dose in young patients affected by type 1 diabetes. <i>PLoS ONE</i> , 2020 , 15, e0244483	3.7	
395	NAFLD as a driver of chronic kidney disease. <i>Journal of Hepatology</i> , 2020 , 72, 785-801	13.4	104
394	Effect of PNPLA3 polymorphism on diagnostic performance of various noninvasive markers for diagnosing and staging nonalcoholic fatty liver disease. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2020 , 35, 1057-1064	4	18
393	PNPLA3 rs738409 is associated with renal glomerular and tubular injury in NAFLD patients with persistently normal ALT levels. <i>Liver International</i> , 2020 , 40, 107-119	7.9	46
392	Liver fibrosis by FibroScan independently of established cardiovascular risk parameters associates with macrovascular and microvascular complications in patients with type 2 diabetes. <i>Liver International</i> , 2020 , 40, 347-354	7.9	27
391	Efficacy and safety of anti-hyperglycaemic drugs in patients with non-alcoholic fatty liver disease with or without diabetes: An updated systematic review of randomized controlled trials. <i>Diabetes and Metabolism</i> , 2020 , 46, 427-441	5.4	42
390	Prospective evaluation of non-alcoholic fatty liver disease by elastographic methods of liver steatosis and fibrosis; controlled attenuation parameter and liver stiffness measurements. <i>Journal of Diabetes and Its Complications</i> , 2020 , 34, 107512	3.2	8
389	What's new in NAFLD pathogenesis, biomarkers and treatment?. <i>Nature Reviews Gastroenterology and Hepatology</i> , 2020 , 17, 70-71	24.2	20
388	High-Dose Vitamin D Supplementation and Bone Health. <i>JAMA - Journal of the American Medical Association</i> , 2020 , 323, 92	27.4	1
387	Treatment algorithm in patients with type 2 diabetes and atherosclerotic cardiovascular disease or high/very high cardiovascular risk. <i>European Heart Journal</i> , 2020 , 41, 331	9.5	3
386	Higher liver stiffness scores are associated with early kidney dysfunction in patients with histologically proven non-cirrhotic NAFLD. <i>Diabetes and Metabolism</i> , 2020 , 46, 288-295	5.4	9
385	Screening for non-alcoholic fatty liver disease using liver stiffness measurement and its association with chronic kidney disease and cardiovascular complications in patients with type 2 diabetes. <i>Diabetes and Metabolism</i> , 2020 , 46, 296-303	5.4	21
384	NAFLD fibrosis score (NFS) can be used in outpatient services to identify chronic vascular complications besides advanced liver fibrosis in type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2020 , 34, 107684	3.2	2
383	Management of type 2 diabetes for prevention of cardiovascular disease. An expert opinion of the Italian Diabetes Society. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2020 , 30, 1926-1936	4.5	4
382	Concordance between MAFLD and NAFLD diagnostic criteria in 'real-world' data. <i>Liver International</i> , 2020 , 40, 2879-2880	7.9	14

381	What's Past Is Prologue: History of Nonalcoholic Fatty Liver Disease. <i>Metabolites</i> , 2020 , 10,	5.6	3
380	Abnormal liver enzymes in children and infants with COVID-19: A narrative review of case-series studies. <i>Pediatric Obesity</i> , 2020 , 15, e12723	4.6	8
379	Nonalcoholic Fatty Liver Disease and Estimated Insulin Resistance in Obese Youth: A Mendelian Randomization Analysis. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020 , 105,	5.6	15
378	ACE2: A Linkage for the Interplay Between COVID-19 and Decompensated Cirrhosis. <i>American Journal of Gastroenterology</i> , 2020 , 115, 1544	0.7	7
377	Human and molecular genetics shed lights on fatty liver disease and diabetes conundrum. <i>Endocrinology, Diabetes and Metabolism</i> , 2020 , 3, e00179	2.7	5
376	Association between specific plasma ceramides and high-sensitivity C-reactive protein levels in postmenopausal women with type 2 diabetes. <i>Diabetes and Metabolism</i> , 2020 , 46, 326-330	5.4	4
375	Associations between specific plasma ceramides and severity of coronary-artery stenosis assessed by coronary angiography. <i>Diabetes and Metabolism</i> , 2020 , 46, 150-157	5.4	16
374	NAFLD and increased risk of cardiovascular disease: clinical associations, pathophysiological mechanisms and pharmacological implications. <i>Gut</i> , 2020 , 69, 1691-1705	19.2	118
373	Association Between Nonalcoholic Fatty Liver Disease and Reduced Bone Mineral Density in Children: A Meta-Analysis. <i>Hepatology</i> , 2019 , 70, 812-823	11.2	16
372	Association between non-alcoholic fatty liver disease and risk of atrial fibrillation in adult individuals: An updated meta-analysis. <i>Liver International</i> , 2019 , 39, 758-769	7.9	43
371	NAFLD in Some Common Endocrine Diseases: Prevalence, Pathophysiology, and Principles of Diagnosis and Management. <i>International Journal of Molecular Sciences</i> , 2019 , 20,	6.3	43
370	Association between non-alcoholic fatty liver disease and decreased lung function in adults: A systematic review and meta-analysis. <i>Diabetes and Metabolism</i> , 2019 , 45, 536-544	5.4	17
369	Association between Helicobacter pylori infection and risk of nonalcoholic fatty liver disease: An updated meta-analysis. <i>Metabolism: Clinical and Experimental</i> , 2019 , 96, 56-65	12.7	24
368	Relationship Between PNPLA3 rs738409 Polymorphism and Decreased Kidney Function in Children With NAFLD. <i>Hepatology</i> , 2019 , 70, 142-153	11.2	23
367	Increased aortic stiffness index in patients with type 1 diabetes without cardiovascular disease compared to controls. <i>Journal of Endocrinological Investigation</i> , 2019 , 42, 1109-1115	5.2	3
366	Letter: non-alcoholic fatty liver disease is associated with a history of osteoporotic fractures but not with low bone mineral density-authors' reply. <i>Alimentary Pharmacology and Therapeutics</i> , 2019 , 49, 961-962	6.1	
365	Influence of hypertriglyceridemia, hyperbilirubinemia and hemolysis on thrombin generation in human plasma. <i>Clinical Chemistry and Laboratory Medicine</i> , 2019 , 57, 1784-1789	5.9	8
364	Pathogenesis of hypothyroidism-induced NAFLD: Evidence for a distinct disease entity?. <i>Digestive and Liver Disease</i> , 2019 , 51, 462-470	3.3	26

363	Plasma N-terminal propeptide of type III procollagen accurately predicts liver fibrosis severity in children with non-alcoholic fatty liver disease. <i>Liver International</i> , 2019 , 39, 2317-2329	7.9	14
362	Increased aortic stiffness in adults with chronic indeterminate Chagas disease. <i>PLoS ONE</i> , 2019 , 14, e0220689	1	
361	Prevalence of prediabetes and diabetes in children and adolescents with biopsy-proven non-alcoholic fatty liver disease. <i>Journal of Hepatology</i> , 2019 , 71, 802-810	13.4	18
360	Contribution of a genetic risk score to clinical prediction of hepatic steatosis in obese children and adolescents. <i>Digestive and Liver Disease</i> , 2019 , 51, 1586-1592	3.3	22
359	Diabetes and NAFLD. <i>Endocrinology</i> , 2019 , 1-27	0.1	
358	Increased red blood cell distribution width and platelet-to-lymphocyte ratio for predicting all-cause mortality in patients with type 2 diabetes and advanced heart failure: a causal association or epiphenomenon?. <i>Kardiologia Polska</i> , 2019 , 77, 587-588	0.9	
357	Association between PNPLA3rs738409 polymorphism decreased kidney function in postmenopausal type 2 diabetic women with or without non-alcoholic fatty liver disease. <i>Diabetes and Metabolism</i> , 2019 , 45, 480-487	5.4	22
356	Systematic review with meta-analysis: non-alcoholic fatty liver disease is associated with a history of osteoporotic fractures but not with low bone mineral density. <i>Alimentary Pharmacology and Therapeutics</i> , 2019 , 49, 375-388	6.1	22
355	Does high LDL-cholesterol cause cardiovascular disease?. <i>Expert Review of Clinical Pharmacology</i> , 2019 , 12, 91	3.8	2
354	Association between non-alcoholic fatty liver disease and bone turnover biomarkers in post-menopausal women with type 2 diabetes. <i>Diabetes and Metabolism</i> , 2019 , 45, 347-355	5.4	17
353	Estimating the real burden of cardiovascular mortality in diabetes. <i>European Review for Medical and Pharmacological Sciences</i> , 2019 , 23, 6700-6706	2.9	2
352	Long-Acting GLP-1 Receptor Agonist Exenatide Influence on the Autonomic Cardiac Sympatho-Vagal Balance. <i>Journal of the Endocrine Society</i> , 2018 , 2, 53-62	0.4	4
351	Prognostic impact of elevated serum uric acid levels on long-term outcomes in patients with chronic heart failure: A post-hoc analysis of the GISSI-HF (Gruppo Italiano per lo Studio della Sopravvivenza nella Insufficienza Cardiaca-Heart Failure) trial. <i>Metabolism: Clinical and Experimental</i> , 2018 , 83, 205-215	12.7	20
350	Mortality from infectious diseases in diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2018 , 28, 444-450	4.5	28
349	Nonalcoholic Fatty Liver Disease and Risk of Incident Type 2 Diabetes: A Meta-analysis. <i>Diabetes Care</i> , 2018 , 41, 372-382	14.6	262
348	Nonalcoholic fatty liver disease and chronic vascular complications of diabetes mellitus. <i>Nature Reviews Endocrinology</i> , 2018 , 14, 99-114	15.2	170
347	Non-alcoholic fatty liver disease and increased risk of all-cause mortality in elderly patients admitted for acute heart failure. <i>International Journal of Cardiology</i> , 2018 , 265, 162-168	3.2	20
346	Liver fat content, non-alcoholic fatty liver disease, and risk of ischaemic heart disease. <i>European Heart Journal</i> , 2018 , 39, 3398	9.5	2

345	Risk of cardiomyopathy and cardiac arrhythmias in patients with nonalcoholic fatty liver disease. <i>Nature Reviews Gastroenterology and Hepatology</i> , 2018 , 15, 425-439	24.2	114
344	Diabetes and NAFLD. <i>Endocrinology</i> , 2018 , 1-27	0.1	
343	Hyperuricemia is associated with an increased prevalence of paroxysmal atrial fibrillation in patients with type 2 diabetes referred for clinically indicated 24-h Holter monitoring. <i>Journal of Endocrinological Investigation</i> , 2018 , 41, 223-231	5.2	14
342	Psoriasis and the metabolic syndrome. <i>Clinics in Dermatology</i> , 2018 , 36, 21-28	3	120
341	Stereotactic body radiotherapy for lung oligometastases impacts on systemic treatment-free survival: a cohort study. <i>Medical Oncology</i> , 2018 , 35, 121	3.7	19
340	Left ventricular chamber dilation and filling pressure may help to categorise patients with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2018 , 6, e000529	4.5	2
339	Tests for diagnosing and monitoring non-alcoholic fatty liver disease in adults. <i>BMJ, The</i> , 2018 , 362, k27349	3.9	54
338	Association Between Primary Hypothyroidism and Nonalcoholic Fatty Liver Disease: A Systematic Review and Meta-Analysis. <i>Thyroid</i> , 2018 , 28, 1270-1284	6.2	50
337	Diabetes and NAFLD. <i>Endocrinology</i> , 2018 , 495-521	0.1	
336	Clinical relevance of liver histopathology and different histological classifications of NASH in adults. <i>Expert Review of Gastroenterology and Hepatology</i> , 2018 , 12, 351-367	4.2	34
335	Hypertension, diabetes, atherosclerosis and NASH: Cause or consequence?. <i>Journal of Hepatology</i> , 2018 , 68, 335-352	13.4	298
334	Nonalcoholic fatty liver disease increases risk of incident chronic kidney disease: A systematic review and meta-analysis. <i>Metabolism: Clinical and Experimental</i> , 2018 , 79, 64-76	12.7	171
333	NAFLD, diabete tipo 2, rischio cardiovascolare?. <i>L Endocrinologo</i> , 2018 , 19, 255-259	0	
332	Association of Plasma Ceramides With Myocardial Perfusion in Patients With Coronary Artery Disease Undergoing Stress Myocardial Perfusion Scintigraphy. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2018 , 38, 2854-2861	9.4	21
331	The E/e' ratio difference between subjects with type 2 diabetes and controls. A meta-analysis of clinical studies. <i>PLoS ONE</i> , 2018 , 13, e0209794	3.7	4
330	Association between decreasing estimated glomerular filtration rate and risk of cardiac conduction defects in patients with type 2 diabetes. <i>Diabetes and Metabolism</i> , 2018 , 44, 473-481	5.4	1
329	Association between plasma ceramides and inducible myocardial ischemia in patients with established or suspected coronary artery disease undergoing myocardial perfusion scintigraphy. <i>Metabolism: Clinical and Experimental</i> , 2018 , 85, 305-312	12.7	11
328	Association between nonalcoholic fatty liver disease and colorectal tumours in asymptomatic adults undergoing screening colonoscopy: a systematic review and meta-analysis. <i>Metabolism: Clinical and Experimental</i> , 2018 , 87, 1-12	12.7	48

327	Early impairment in left ventricular longitudinal systolic function is associated with an increased risk of incident atrial fibrillation in patients with type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 413-418	3.2	19
326	Psychological distress, self-efficacy and glycemic control in type 2 diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2017 , 27, 300-306	4.5	30
325	AISF position paper on nonalcoholic fatty liver disease (NAFLD): Updates and future directions. <i>Digestive and Liver Disease</i> , 2017 , 49, 471-483	3.3	179
324	Non-alcoholic fatty liver disease: an emerging driving force in chronic kidney disease. <i>Nature Reviews Nephrology</i> , 2017 , 13, 297-310	14.9	146
323	Nonalcoholic fatty liver disease is associated with an increased prevalence of distal symmetric polyneuropathy in adult patients with type 1 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 1021-1026	3.2	30
322	Association Between Diabetes and 1-Year Adverse Clinical Outcomes in a Multinational Cohort of Ambulatory Patients With Chronic Heart Failure: Results From the ESC-HFA Heart Failure Long-Term Registry. <i>Diabetes Care</i> , 2017 , 40, 671-678	14.6	65
321	Association between subclinical left ventricular systolic dysfunction and glycemic control in asymptomatic type 2 diabetic patients with preserved left ventricular function. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 1035-1040	3.2	6
320	Non-alcoholic fatty liver disease as driving force in coronary heart disease?. <i>Gut</i> , 2017 , 66, 213-214	19.2	9
319	Effect of aspirin on renal disease progression in patients with type 2 diabetes: A multicenter, double-blind, placebo-controlled, randomized trial. The renal disease progression by aspirin in diabetic patients (LEDA) trial. Rationale and study design. <i>American Heart Journal</i> , 2017 , 189, 120-127	4.9	6
318	Ultrasonographic fatty liver indicator detects mild steatosis and correlates with metabolic/histological parameters in various liver diseases. <i>Metabolism: Clinical and Experimental</i> , 2017 , 72, 57-65	12.7	80
317	Low-grade endotoxemia, gut permeability and platelet activation in patients with impaired fasting glucose. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2017 , 27, 890-895	4.5	19
316	Non-alcoholic fatty liver disease and its relationship with cardiovascular disease and other extrahepatic diseases. <i>Gut</i> , 2017 , 66, 1138-1153	19.2	508
315	Relation of elevated serum uric acid levels to first-degree heart block and other cardiac conduction defects in hospitalized patients with type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 1691-1697	3.2	5
314	Prognostic Impact of Diabetes on Long-term Survival Outcomes in Patients With Heart Failure: A Meta-analysis. <i>Diabetes Care</i> , 2017 , 40, 1597-1605	14.6	52
313	Prognostic Impact of Diabetes and Prediabetes on Survival Outcomes in Patients With Chronic Heart Failure: A Post-Hoc Analysis of the GISSI-HF (Gruppo Italiano per lo Studio della Sopravvivenza nella Insufficienza Cardiaca-Heart Failure) Trial. <i>Journal of the American Heart Association</i> , 2017 , 6,	6	39
312	In-hospital and 1-year mortality associated with diabetes in patients with acute heart failure: results from the ESC-HFA Heart Failure Long-Term Registry. <i>European Journal of Heart Failure</i> , 2017 , 19, 54-65	12.3	92
311	Plasma Leptin in Patients at Intermediate to High Cardiovascular Risk With and Without Type 2 Diabetes Mellitus. <i>Journal of Clinical Laboratory Analysis</i> , 2017 , 31,	3	2
310	Mitral Regurgitation and Increased Risk of All-Cause and Cardiovascular Mortality in Patients with Type 2 Diabetes. <i>American Journal of Medicine</i> , 2017 , 130, 70-76.e1	2.4	15

309	Type 2 diabetes mellitus and risk of hepatocellular carcinoma: spotlight on nonalcoholic fatty liver disease. <i>Annals of Translational Medicine</i> , 2017 , 5, 270	3.2	68
308	Nonalcoholic fatty liver disease and increased risk of 1-year all-cause and cardiac hospital readmissions in elderly patients admitted for acute heart failure. <i>PLoS ONE</i> , 2017 , 12, e0173398	3.7	26
307	Nonalcoholic fatty liver disease is associated with an increased risk of heart block in hospitalized patients with type 2 diabetes mellitus. <i>PLoS ONE</i> , 2017 , 12, e0185459	3.7	29
306	Non-alcoholic fatty liver disease and risk of cardiovascular disease. <i>Metabolism: Clinical and Experimental</i> , 2016 , 65, 1136-50	12.7	152
305	Hyperuricemia is associated with an increased prevalence of atrial fibrillation in hospitalized patients with type 2 diabetes. <i>Journal of Endocrinological Investigation</i> , 2016 , 39, 159-67	5.2	25
304	Nonalcoholic fatty liver disease: cause or consequence of type 2 diabetes?. <i>Liver International</i> , 2016 , 36, 1563-1579	7.9	96
303	Nonalcoholic fatty liver disease is independently associated with an increased incidence of cardiovascular disease in adult patients with type 1 diabetes. <i>International Journal of Cardiology</i> , 2016 , 225, 387-391	3.2	39
302	The aspartate aminotransferase-to-alanine aminotransferase ratio predicts all-cause and cardiovascular mortality in patients with type 2 diabetes. <i>Medicine (United States)</i> , 2016 , 95, e4821	1.8	19
301	Evidence of left atrial remodeling and left ventricular diastolic dysfunction in type 2 diabetes mellitus with preserved systolic function. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2016 , 26, 1026-1032	4.5	11
300	Obesity: Metabolically healthy obesity and NAFLD. <i>Nature Reviews Gastroenterology and Hepatology</i> , 2016 , 13, 442-4	24.2	28
299	Evidence that non-alcoholic fatty liver disease and polycystic ovary syndrome are associated by necessity rather than chance: a novel hepato-ovarian axis?. <i>Endocrine</i> , 2016 , 51, 211-21	4	53
298	Cardiovascular Disease and Myocardial Abnormalities in Nonalcoholic Fatty Liver Disease. <i>Digestive Diseases and Sciences</i> , 2016 , 61, 1246-67	4	75
297	Nonalcoholic Fatty Liver Disease Is Associated With Higher 1-year All-Cause Rehospitalization Rates in Patients Admitted for Acute Heart Failure. <i>Medicine (United States)</i> , 2016 , 95, e2760	1.8	15
296	A "systems medicine" approach to the study of non-alcoholic fatty liver disease. <i>Digestive and Liver Disease</i> , 2016 , 48, 333-42	3.3	42
295	Association between hepatic steatosis and serum liver enzyme levels with atrial fibrillation in the general population: The Study of Health in Pomerania (SHIP). <i>Atherosclerosis</i> , 2016 , 245, 123-31	3.1	33
294	Prognostic impact of in-hospital hyperglycemia in hospitalized patients with acute heart failure: Results of the IN-HF (Italian Network on Heart Failure) Outcome registry. <i>International Journal of Cardiology</i> , 2016 , 203, 587-93	3.2	27
293	Black esophagus syndrome associated with diabetic ketoacidosis. <i>World Journal of Clinical Cases</i> , 2016 , 4, 56-9	1.6	17
292	Relationship between Non-Alcoholic Fatty Liver Disease and Psoriasis: A Novel Hepato-Dermal Axis?. <i>International Journal of Molecular Sciences</i> , 2016 , 17, 217	6.3	47

291	Type 2 Diabetes in Non-Alcoholic Fatty Liver Disease and Hepatitis C Virus Infection--Liver: The "Musketeer" in the Spotlight. <i>International Journal of Molecular Sciences</i> , 2016 , 17, 355	6.3	30
290	Fatty liver is associated with an increased risk of diabetes and cardiovascular disease - Evidence from three different disease models: NAFLD, HCV and HIV. <i>World Journal of Gastroenterology</i> , 2016 , 22, 9674-9693	5.6	69
289	Echocardiographically Derived Pulse Wave Velocity and Diastolic Dysfunction Are Associated with an Increased Incidence of Atrial Fibrillation in Patients with Systolic Heart Failure. <i>Echocardiography</i> , 2016 , 33, 1024-31	1.5	7
288	Nonalcoholic fatty liver disease, cardiovascular outcomes, and mortality in patients undergoing a coronary angiogram. <i>Hepatology</i> , 2016 , 64, 684-5	11.2	3
287	Global epidemiology of nonalcoholic fatty liver disease: Meta-analytic assessment of prevalence, incidence, and outcomes. <i>Hepatology</i> , 2016 , 64, 1388-9	11.2	60
286	Nonalcoholic fatty liver disease is associated with an almost twofold increased risk of incident type 2 diabetes and metabolic syndrome. Evidence from a systematic review and meta-analysis. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2016 , 31, 936-44	4	378
285	Nonalcoholic Fatty Liver Disease Is Associated With Ventricular Arrhythmias in Patients With Type 2 Diabetes Referred for Clinically Indicated 24-Hour Holter Monitoring. <i>Diabetes Care</i> , 2016 , 39, 1416-23	14.6	66
284	Factors associated with significant liver steatosis and fibrosis as assessed by transient elastography in patients with one or more components of the metabolic syndrome. <i>Journal of Diabetes and Its Complications</i> , 2016 , 30, 1347-53	3.2	25
283	Non-alcoholic fatty liver disease and risk of incident cardiovascular disease: A meta-analysis. <i>Journal of Hepatology</i> , 2016 , 65, 589-600	13.4	640
282	EASL-EASD-EASO Clinical Practice Guidelines for the management of non-alcoholic fatty liver disease: is universal screening appropriate?. <i>Diabetologia</i> , 2016 , 59, 1141-4	10.3	59
281	Risk of type 2 diabetes in patients with non-alcoholic fatty liver disease: Causal association or epiphenomenon?. <i>Diabetes and Metabolism</i> , 2016 , 42, 142-56	5.4	59
280	Severe hypoglycemia in patients with known diabetes requiring emergency department care: A report from an Italian multicenter study. <i>Journal of Clinical and Translational Endocrinology</i> , 2016 , 5, 46-52 ⁴		4
279	The mean platelet volume is significantly associated with higher glycosylated hemoglobin in a large population of unselected outpatients. <i>Primary Care Diabetes</i> , 2015 , 9, 226-30	2.4	15
278	Prevalence of diabetes across different immigrant groups in North-eastern Italy. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2015 , 25, 924-30	4.5	13
277	Prevalence of neuropathy in type 2 diabetic patients and its association with other diabetes complications: The Verona Diabetic Foot Screening Program. <i>Journal of Diabetes and Its Complications</i> , 2015 , 29, 1066-70	3.2	53
276	Relationship between increased left atrial volume and microvascular complications in patients with type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2015 , 29, 822-8	3.2	9
275	Prevalence of Cardiovascular Autonomic Neuropathy in a Cohort of Patients With Newly Diagnosed Type 2 Diabetes: The Verona Newly Diagnosed Type 2 Diabetes Study (VNDS). <i>Diabetes Care</i> , 2015 , 38, 1487-93	14.6	42
274	NAFLD: a multisystem disease. <i>Journal of Hepatology</i> , 2015 , 62, S47-64	13.4	1286

273	Circulating Markers of Liver Function and Cardiovascular Disease Risk. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2015 , 35, 2290-6	9.4	36
272	A Perspective on Metabolic Syndrome and Nonalcoholic Fatty Liver Disease. <i>Metabolic Syndrome and Related Disorders</i> , 2015 , 13, 235-8	2.6	21
271	Epidemiological modifiers of non-alcoholic fatty liver disease: Focus on high-risk groups. <i>Digestive and Liver Disease</i> , 2015 , 47, 997-1006	3.3	279
270	Diagnosis and management of cardiovascular risk in nonalcoholic fatty liver disease. <i>Expert Review of Gastroenterology and Hepatology</i> , 2015 , 9, 629-50	4.2	57
269	Nonalcoholic Fatty Liver Disease Is Independently Associated with Early Left Ventricular Diastolic Dysfunction in Patients with Type 2 Diabetes. <i>PLoS ONE</i> , 2015 , 10, e0135329	3.7	61
268	Nonalcoholic fatty liver disease and decreased bone mineral density: is there a link?. <i>Journal of Endocrinological Investigation</i> , 2015 , 38, 817-25	5.2	52
267	Heart valve calcification in patients with type 2 diabetes and nonalcoholic fatty liver disease. <i>Metabolism: Clinical and Experimental</i> , 2015 , 64, 879-87	12.7	55
266	Lower levels of 25-hydroxyvitamin D3 are associated with a higher prevalence of microvascular complications in patients with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2015 , 3, e000058	4.5	32
265	Usefulness of subclinical left ventricular midwall dysfunction to predict cardiovascular mortality in patients with type 2 diabetes mellitus. <i>American Journal of Cardiology</i> , 2014 , 113, 1409-14	3	20
264	Nonalcoholic fatty liver disease is independently associated with an increased incidence of chronic kidney disease in patients with type 1 diabetes. <i>Diabetes Care</i> , 2014 , 37, 1729-36	14.6	98
263	Mortality from chronic liver diseases in diabetes. <i>American Journal of Gastroenterology</i> , 2014 , 109, 1020-5.7		83
262	Adiponectin and migraine: systematic review of clinical evidence. <i>Neurological Sciences</i> , 2014 , 35, 1167-73.5		10
261	CKD and nonalcoholic fatty liver disease. <i>American Journal of Kidney Diseases</i> , 2014 , 64, 638-52	7.4	120
260	Ectopic fat, insulin resistance, and nonalcoholic fatty liver disease: implications for cardiovascular disease. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2014 , 34, 1155-61	9.4	99
259	Risk of cardiovascular, cardiac and arrhythmic complications in patients with non-alcoholic fatty liver disease. <i>World Journal of Gastroenterology</i> , 2014 , 20, 1724-45	5.6	160
258	Reply: To PMID 23504926. <i>Hepatology</i> , 2014 , 59, 352	11.2	
257	Hemostatic and fibrinolytic abnormalities in polycystic ovary syndrome. <i>Seminars in Thrombosis and Hemostasis</i> , 2014 , 40, 600-18	5.3	13
256	Inter-atrial shunt inversion by the sitting position in a patient with a patent foramen ovale and acute pulmonary embolism. <i>European Heart Journal</i> , 2014 , 35, 1032	9.5	

255	Hyperuricemia in patients with chronic plaque psoriasis. <i>Journal of the American Academy of Dermatology</i> , 2014 , 70, 127-30	4.5	38
254	Association of nonalcoholic fatty liver disease with QTc interval in patients with type 2 diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2014 , 24, 663-9	4.5	55
253	Nonalcoholic fatty liver disease is associated with aortic valve sclerosis in patients with type 2 diabetes mellitus. <i>PLoS ONE</i> , 2014 , 9, e88371	3.7	39
252	Increased red blood cell distribution width (RDW) is associated with higher glycosylated hemoglobin (HbA1c) in the elderly. <i>Clinical Laboratory</i> , 2014 , 60, 2095-8	2	22
251	Increased aortic pulse wave velocity as measured by echocardiography is strongly associated with poor prognosis in patients with heart failure. <i>Journal of the American Society of Echocardiography</i> , 2013 , 26, 714-20	5.8	26
250	Both resistance training and aerobic training reduce hepatic fat content in type 2 diabetic subjects with nonalcoholic fatty liver disease (the RAED2 Randomized Trial). <i>Hepatology</i> , 2013 , 58, 1287-95	11.2	211
249	Elevated serum uric acid levels are associated with non-alcoholic fatty liver disease independently of metabolic syndrome features in the United States: Liver ultrasound data from the National Health and Nutrition Examination Survey. <i>Metabolism: Clinical and Experimental</i> , 2013 , 62, 392-9	12.7	106
248	Low 25-hydroxyvitamin D level is independently associated with non-alcoholic fatty liver disease. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2013 , 23, 792-8	4.5	51
247	Relation of elevated serum uric acid levels to incidence of atrial fibrillation in patients with type 2 diabetes mellitus. <i>American Journal of Cardiology</i> , 2013 , 112, 499-504	3	53
246	Inappropriate left ventricular mass independently predicts cardiovascular mortality in patients with type 2 diabetes. <i>International Journal of Cardiology</i> , 2013 , 168, 4953-6	3.2	13
245	Progression of NAFLD to diabetes mellitus, cardiovascular disease or cirrhosis. <i>Nature Reviews Gastroenterology and Hepatology</i> , 2013 , 10, 330-44	24.2	1022
244	Clinical Review: Nonalcoholic fatty liver disease: a novel cardiometabolic risk factor for type 2 diabetes and its complications. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, 483-95	5.6	203
243	Nonalcoholic fatty liver disease and reduced serum vitamin D(3) levels. <i>Metabolic Syndrome and Related Disorders</i> , 2013 , 11, 217-28	2.6	24
242	Uric acid as a target of therapy in CKD. <i>American Journal of Kidney Diseases</i> , 2013 , 61, 134-46	7.4	165
241	Non-alcoholic fatty liver disease is associated with an increased prevalence of atrial fibrillation in hospitalized patients with type 2 diabetes. <i>Clinical Science</i> , 2013 , 125, 301-9	6.5	80
240	Diagnosis and management of nonalcoholic fatty liver disease and its hemostatic/thrombotic and vascular complications. <i>Seminars in Thrombosis and Hemostasis</i> , 2013 , 39, 214-28	5.3	41
239	Non-alcoholic fatty liver disease is associated with an increased incidence of atrial fibrillation in patients with type 2 diabetes. <i>PLoS ONE</i> , 2013 , 8, e57183	3.7	119
238	Cardiovascular and Systemic Risk in Nonalcoholic Fatty Liver Disease - Atherosclerosis as a Major Player in the Natural Course of NAFLD. <i>Current Pharmaceutical Design</i> , 2013 , 19, 5177-5192	3.3	80

237	Glycated haemoglobin is inversely related to serum vitamin D levels in type 2 diabetic patients. <i>PLoS ONE</i> , 2013 , 8, e82733	3.7	36
236	Cardiovascular and Systemic Risk in Nonalcoholic Fatty Liver Disease - Atherosclerosis as a Major Player in the Natural Course of NAFLD. <i>Current Pharmaceutical Design</i> , 2013 , 19, 5177-5192	3.3	70
235	Cardiovascular and systemic risk in nonalcoholic fatty liver disease - atherosclerosis as a major player in the natural course of NAFLD. <i>Current Pharmaceutical Design</i> , 2013 , 19, 5177-92	3.3	45
234	Serum uric acid levels and incident chronic kidney disease in patients with type 2 diabetes and preserved kidney function. <i>Diabetes Care</i> , 2012 , 35, 99-104	14.6	167
233	Non-alcoholic fatty liver disease is independently associated with left ventricular hypertrophy in hypertensive Type 2 diabetic individuals. <i>Journal of Endocrinological Investigation</i> , 2012 , 35, 215-8	5.2	44
232	ABO blood group, hypercoagulability, and cardiovascular and cancer risk. <i>Critical Reviews in Clinical Laboratory Sciences</i> , 2012 , 49, 137-49	9.4	76
231	Random plasma glucose measurement may improve the diagnostic specificity of highly sensitive troponin in the emergency department. <i>International Journal of Cardiology</i> , 2012 , 155, 172-3	3.2	4
230	Erythrocyte mechanical fragility is increased in patients with type 2 diabetes. <i>European Journal of Internal Medicine</i> , 2012 , 23, 150-3	3.9	43
229	Increased risk of cardiovascular disease and chronic kidney disease in NAFLD. <i>Nature Reviews Gastroenterology and Hepatology</i> , 2012 , 9, 372-81	24.2	90
228	Aortic and mitral annular calcifications are predictive of all-cause and cardiovascular mortality in patients with type 2 diabetes. <i>Diabetes Care</i> , 2012 , 35, 1781-6	14.6	52
227	Diagnostic significance of haematological testing in patients presenting at the Emergency Department. <i>Emergency Care Journal</i> , 2012 , 8, 7	1.2	1
226	Predictable impact of the routine implementation of the CKD-EPI equation for estimating glomerular filtration rate by a simulation study. <i>Rivista Italiana Della Medicina Di Laboratorio</i> , 2012 , 8, 107-113	1.1	
225	Increased prevalence of chronic kidney disease in patients with Type 1 diabetes and non-alcoholic fatty liver. <i>Diabetic Medicine</i> , 2012 , 29, 220-6	3.5	48
224	Optimal therapy for reduction of lipoprotein(a). <i>Journal of Clinical Pharmacy and Therapeutics</i> , 2012 , 37, 1-3	2.2	23
223	Higher random plasma glucose level is associated with increased plasma cardiac troponin in emergency department patients with suspected acute coronary syndrome. <i>American Journal of Cardiology</i> , 2012 , 109, 775-6	3	1
222	Haemoglobin A1c and diagnosis of diabetes. Not ready for the prime time?. <i>Annals of Clinical Biochemistry</i> , 2012 , 49, 508	2.2	3
221	Increased pulse pressure independently predicts incident atrial fibrillation in patients with type 2 diabetes. <i>Diabetes Care</i> , 2012 , 35, 2337-9	14.6	18
220	Predictors of estimated GFR decline in patients with type 2 diabetes and preserved kidney function. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2012 , 7, 401-8	6.9	138

219	Nonalcoholic fatty liver disease is associated with left ventricular diastolic dysfunction in patients with type 2 diabetes. <i>Diabetes Care</i> , 2012 , 35, 389-95	14.6	134
218	Pasireotide in Cushing's disease. <i>New England Journal of Medicine</i> , 2012 , 366, 2134; author reply 2134-5	59.2	12
217	Vitamin D, thrombosis, and hemostasis: more than skin deep. <i>Seminars in Thrombosis and Hemostasis</i> , 2012 , 38, 114-24	5.3	49
216	Further insights on the relationship between bilirubin and C-reactive protein. <i>Clinical Chemistry and Laboratory Medicine</i> , 2012 , 50, 2229-30	5.9	4
215	Comparison of two creatinine-based estimating equations in predicting all-cause and cardiovascular mortality in patients with type 2 diabetes. <i>Diabetes Care</i> , 2012 , 35, 2347-53	14.6	20
214	The effect of combined calcium and cholecalciferol supplementation on bone mineral density in elderly women with moderate chronic kidney disease. <i>Clinical Nephrology</i> , 2012 , 77, 358-65	2.1	7
213	Association between nonalcoholic liver disease and chronic kidney disease: an ultrasound analysis from NHANES 1988-1994. <i>American Journal of Nephrology</i> , 2012 , 36, 466-71	4.6	54
212	Increased prevalence of cardiovascular disease in Type 1 diabetic patients with non-alcoholic fatty liver disease. <i>Journal of Endocrinological Investigation</i> , 2012 , 35, 535-40	5.2	19
211	Pancreatic fat accumulation and its relationship with liver fat content and other fat depots in obese individuals. <i>Journal of Endocrinological Investigation</i> , 2012 , 35, 748-53	5.2	12
210	Arterial thrombus formation in cardiovascular disease. <i>Nature Reviews Cardiology</i> , 2011 , 8, 502-12	14.8	180
209	Hyperthyroidism and venous thrombosis: a casual or causal association? A systematic literature review. <i>Clinical and Applied Thrombosis/Hemostasis</i> , 2011 , 17, 387-92	3.3	39
208	Screening and therapeutic management of lipoprotein(a) excess: review of the epidemiological evidence, guidelines and recommendations. <i>Clinica Chimica Acta</i> , 2011 , 412, 797-801	6.2	20
207	Detection of chronic kidney disease in hospitalized patients: is one estimating glomerular filtration rate equation better than another?. <i>European Journal of Internal Medicine</i> , 2011 , 22, 119-20; author reply 120-1	3.9	7
206	Risk of chronic kidney disease in patients with non-alcoholic fatty liver disease: is there a link?. <i>Journal of Hepatology</i> , 2011 , 54, 1020-9	13.4	131
205	A laboratory standpoint on the role of hemoglobin A1c for the diagnosis of diabetes in childhood: more doubts than certainties?. <i>Pediatric Diabetes</i> , 2011 , 12, 183-6	3.6	19
204	Tomatoes, lycopene-containing foods and cancer risk. <i>British Journal of Cancer</i> , 2011 , 104, 1234-5	8.7	10
203	Liver enzymes, nonalcoholic fatty liver disease, and incident cardiovascular disease. <i>Hepatology</i> , 2011 , 53, 375	11.2	11
202	Glomerular filtration rate, albuminuria and risk of cardiovascular and all-cause mortality in type 2 diabetic individuals. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2011 , 21, 294-301	4.5	23

201	The role of serum uric acid in cardiovascular disease in type 2 diabetic and non-diabetic subjects: a narrative review. <i>Journal of Endocrinological Investigation</i> , 2011 , 34, 881-6	5.2	22
200	Extra-skeletal effects of vitamin D deficiency in chronic kidney disease. <i>Annals of Medicine</i> , 2011 , 43, 273-82	1.5	24
199	Safety of recombinant activated factor VII in randomized clinical trials. <i>New England Journal of Medicine</i> , 2011 , 364, 575; author reply 575-6	59.2	5
198	Venous thromboembolism in chronic liver disease. <i>Seminars in Thrombosis and Hemostasis</i> , 2011 , 37, 66-76; 3		6
197	Hemostatic disorders in type 1 diabetes mellitus. <i>Seminars in Thrombosis and Hemostasis</i> , 2011 , 37, 58-65; 3		18
196	Impact of reference category and number of traits in the cluster on risk of coronary heart disease in metabolic syndrome: prospective data from the Bruneck study. <i>Metabolic Syndrome and Related Disorders</i> , 2011 , 9, 313-8	2.6	1
195	Risk of all-cause and cardiovascular mortality in patients with chronic liver disease. <i>Gut</i> , 2011 , 60, 1602-3; author reply 1603-4	19.2	10
194	Relationship between early diastolic dysfunction and abnormal microvolt T-wave alternans in patients with type 2 diabetes. <i>Circulation: Cardiovascular Imaging</i> , 2011 , 4, 408-14	3.9	11
193	High-normal HbA1c is a strong predictor of type 2 diabetes in the general population. <i>Diabetes Care</i> , 2011 , 34, 1038-40	14.6	39
192	High-sensitivity C-reactive protein, obesity, and subclinical atherosclerosis: implications of JUPITER from the MESA study. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2011 , 31, 1251-2	9.4	7
191	Relationship of hepatic steatosis and alanine aminotransferase with coronary calcification. <i>Clinical Chemistry and Laboratory Medicine</i> , 2011 , 49, 741	5.9	
190	Commentary: Liver enzymes and the risk of adverse cardiovascular outcomes--the lower, the better?. <i>International Journal of Epidemiology</i> , 2011 , 40, 1539-41	7.8	8
189	Risk of cardiovascular disease and chronic kidney disease in diabetic patients with non-alcoholic fatty liver disease: just a coincidence?. <i>Journal of Endocrinological Investigation</i> , 2011 , 34, 544-51	5.2	13
188	Relationship between kidney function and liver histology in subjects with nonalcoholic steatohepatitis. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2010 , 5, 2166-71	6.9	156
187	Glycated hemoglobin (HbA1c): old dogmas, a new perspective?. <i>Clinical Chemistry and Laboratory Medicine</i> , 2010 , 48, 609-14	5.9	48
186	Serum bilirubin levels and cardiovascular disease risk: a Janus Bifrons?. <i>Advances in Clinical Chemistry</i> , 2010 , 50, 47-63	5.8	46
185	Moderate red wine consumption and cardiovascular disease risk: beyond the "French paradox". <i>Seminars in Thrombosis and Hemostasis</i> , 2010 , 36, 59-70	5.3	122
184	Disorders of coagulation and hemostasis in abdominal obesity: emerging role of fatty liver. <i>Seminars in Thrombosis and Hemostasis</i> , 2010 , 36, 41-8	5.3	39

183	Disorders of hemostasis associated with chronic kidney disease. <i>Seminars in Thrombosis and Hemostasis</i> , 2010 , 36, 34-40	5.3	142
182	Elevated serum gamma-glutamyltransferase activity is associated with increased risk of mortality, incident type 2 diabetes, cardiovascular events, chronic kidney disease and cancer - a narrative review. <i>Clinical Chemistry and Laboratory Medicine</i> , 2010 , 48, 147-57	5.9	83
181	Prevention and treatment of nonalcoholic fatty liver disease. <i>Digestive and Liver Disease</i> , 2010 , 42, 331-40	4.3	15
180	Risk of cardiovascular disease in patients with nonalcoholic fatty liver disease. <i>New England Journal of Medicine</i> , 2010 , 363, 1341-50	59.2	1353
179	Prevalence of non-alcoholic fatty liver disease and its association with cardiovascular disease in patients with type 1 diabetes. <i>Journal of Hepatology</i> , 2010 , 53, 713-8	13.4	171
178	Vaccination, squalene and anti-squalene antibodies: facts or fiction?. <i>European Journal of Internal Medicine</i> , 2010 , 21, 70-3	3.9	42
177	Relationship between serum gamma-glutamyltransferase and chronic kidney disease in the United States adult population. Findings from the National Health and Nutrition Examination Survey 2001-2006. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2010 , 20, 583-90	4.5	47
176	Anaemia, independent of chronic kidney disease, predicts all-cause and cardiovascular mortality in type 2 diabetic patients. <i>Atherosclerosis</i> , 2010 , 210, 575-80	3.1	26
175	Usefulness of the triglyceride to high-density lipoprotein cholesterol ratio for predicting mortality risk in type 2 diabetes: role of kidney dysfunction. <i>Atherosclerosis</i> , 2010 , 212, 287-91	3.1	14
174	Hemostatic abnormalities in endocrine and metabolic disorders. <i>European Journal of Endocrinology</i> , 2010 , 162, 439-51	6.5	44
173	The use of recombinant activated FVII in postpartum hemorrhage. <i>Clinical Obstetrics and Gynecology</i> , 2010 , 53, 219-27	1.7	106
172	Epidemiological association between uric acid concentration in plasma, lipoprotein(a), and the traditional lipid profile. <i>Clinical Cardiology</i> , 2010 , 33, E76-80	3.3	49
171	Non-alcoholic fatty liver disease is independently associated with an increased prevalence of chronic kidney disease and retinopathy in type 1 diabetic patients. <i>Diabetologia</i> , 2010 , 53, 1341-8	10.3	115
170	Non-alcoholic Fatty Liver Disease and Cardiovascular Disease Risk. <i>Current Cardiovascular Risk Reports</i> , 2010 , 4, 32-39	0.9	4
169	Effect of lovastatin on primary prevention of cardiovascular events in mild CKD and kidney function loss: a post hoc analysis of the Air Force/Texas Coronary Atherosclerosis Prevention Study. <i>American Journal of Kidney Diseases</i> , 2010 , 55, 42-9	7.4	43
168	Relation of serum phosphorus levels to ankle brachial pressure index (from the Third National Health and Nutrition Examination Survey). <i>American Journal of Cardiology</i> , 2010 , 106, 564-8	3	36
167	Determinants of anaemia in the very elderly: a major contribution from impaired renal function?. <i>Blood Transfusion</i> , 2010 , 8, 44-8	3.6	9
166	Elevated serum uric acid concentrations independently predict cardiovascular mortality in type 2 diabetic patients. <i>Diabetes Care</i> , 2009 , 32, 1716-20	14.6	94

165	Prevalence of thyroid autoimmunity and subclinical hypothyroidism in persons with chronic kidney disease not requiring chronic dialysis. <i>Clinical Chemistry and Laboratory Medicine</i> , 2009 , 47, 1367-71	5.9	31
164	Clinical usefulness of measuring red blood cell distribution width on admission in patients with acute coronary syndromes. <i>Clinical Chemistry and Laboratory Medicine</i> , 2009 , 47, 353-7	5.9	70
163	Relationship of serum bilirubin concentrations to kidney function and albuminuria in the United States adult population. Findings from the National Health and Nutrition Examination Survey 2001-2006. <i>Clinical Chemistry and Laboratory Medicine</i> , 2009 , 47, 1055-62	5.9	35
162	Vitamin B12, folate, and anemia in old age. <i>Archives of Internal Medicine</i> , 2009 , 169, 716		3
161	Genetic and biochemical heterogeneity of cardiac troponins: clinical and laboratory implications. <i>Clinical Chemistry and Laboratory Medicine</i> , 2009 , 47, 1183-94	5.9	31
160	Procalcitonin values after dialysis is closely related to type of dialysis membrane. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2009 , 69, 703-7	2	14
159	Extracorporeal immunoabsorption for the treatment of coagulation inhibitors. <i>Seminars in Thrombosis and Hemostasis</i> , 2009 , 35, 76-80	5.3	17
158	Hemostatic and fibrinolytic abnormalities in endocrine diseases: a narrative review. <i>Seminars in Thrombosis and Hemostasis</i> , 2009 , 35, 605-12	5.3	10
157	Interpatient phenotypic inconsistency in severe congenital hemophilia: a systematic review of the role of inherited thrombophilia. <i>Seminars in Thrombosis and Hemostasis</i> , 2009 , 35, 307-12	5.3	36
156	Nonalcoholic fatty liver disease as a contributor to hypercoagulation and thrombophilia in the metabolic syndrome. <i>Seminars in Thrombosis and Hemostasis</i> , 2009 , 35, 277-87	5.3	106
155	25-Hydroxyvitamin D deficiency and inflammation and their association with hemoglobin levels in chronic kidney disease. <i>American Journal of Nephrology</i> , 2009 , 30, 64-72	4.6	52
154	Biochemical correlates of lipoprotein(a) in a general adult population. Possible implications for cardiovascular risk assessment. <i>Journal of Thrombosis and Thrombolysis</i> , 2009 , 27, 44-7	5.1	7
153	Inherited and acquired risk factors for arterial ischemic stroke in childhood. <i>Journal of Thrombosis and Thrombolysis</i> , 2009 , 27, 239-48	5.1	5
152	Eosinophilia and first-line coagulation testing. <i>Journal of Thrombosis and Thrombolysis</i> , 2009 , 28, 90-3	5.1	8
151	Antithrombotic prophylaxis in patients with von Willebrand disease undergoing major surgery: when is it necessary?. <i>Journal of Thrombosis and Thrombolysis</i> , 2009 , 28, 215-9	5.1	12
150	Hyperthyroidism is associated with shortened APTT and increased fibrinogen values in a general population of unselected outpatients. <i>Journal of Thrombosis and Thrombolysis</i> , 2009 , 28, 362-5	5.1	26
149	Dark chocolate: consumption for pleasure or therapy?. <i>Journal of Thrombosis and Thrombolysis</i> , 2009 , 28, 482-8	5.1	17
148	The red blood cell distribution width is associated with serum levels of thyroid stimulating hormone in the general population. <i>International Journal of Laboratory Hematology</i> , 2009 , 31, 581-2	2.5	14

147	Relationship of serum gamma-glutamyltransferase to atherogenic dyslipidemia and glycemic control in type 2 diabetes. <i>Obesity</i> , 2009 , 17, 370-4	8	14
146	Response to NASH Predicts Plasma Inflammatory Biomarkers Independently of Visceral Fat in Men <i>Obesity</i> , 2009 , 17, 627-627	8	
145	Bilirubin concentration and cardiovascular risk profile. <i>Liver International</i> , 2009 , 29, 315-6	7.9	7
144	Epidemiological association between fasting plasma glucose and shortened APTT. <i>Clinical Biochemistry</i> , 2009 , 42, 118-20	3.5	33
143	Abnormal serum alanine aminotransferase levels are associated with impaired insulin sensitivity in young women with polycystic ovary syndrome. <i>Journal of Endocrinological Investigation</i> , 2009 , 32, 695-700	5.2	27
142	Hemophilia and cancer: a new challenge for hemophilia centers. <i>Cancer Treatment Reviews</i> , 2009 , 35, 374-7	14.4	38
141	Relationship between serum phosphate and cardiovascular risk factors in a large cohort of adult outpatients. <i>Diabetes Research and Clinical Practice</i> , 2009 , 84, e3-5	7.4	14
140	Relationship between albuminuria and hemoglobin level. <i>Diabetes Research and Clinical Practice</i> , 2009 , 86, e62-3	7.4	1
139	Is fasting glucose variability a risk factor for retinopathy in people with type 2 diabetes?. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2009 , 19, 334-9	4.5	33
138	Higher HDL cholesterol levels are associated with a lower incidence of chronic kidney disease in patients with type 2 diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2009 , 19, 580-6	4.5	41
137	Migraine, valproic acid, and lipoprotein(a). <i>Pediatric Neurology</i> , 2009 , 41, 78; author reply 78-9	2.9	
136	Non-alcoholic fatty liver disease in patients with chronic plaque psoriasis. <i>Journal of Hepatology</i> , 2009 , 51, 758-64	13.4	179
135	Relationship between serum bilirubin and kidney function in non-diabetic and diabetic individuals. <i>Kidney International</i> , 2009 , 75, 863	9.9	18
134	25-Hydroxyvitamin D deficiency is independently associated with cardiovascular disease in the Third National Health and Nutrition Examination Survey. <i>Atherosclerosis</i> , 2009 , 205, 255-60	3.1	315
133	Effect of simvastatin on kidney function loss in patients with coronary heart disease: findings from the Scandinavian Simvastatin Survival Study (4S). <i>Atherosclerosis</i> , 2009 , 205, 202-6	3.1	34
132	Anaphylaxis in patients with congenital bleeding disorders and inhibitors. <i>Blood Coagulation and Fibrinolysis</i> , 2009 , 20, 225-9	1	24
131	Risks and benefits of replacing conventional plasma lipids with apolipoprotein measurement. <i>Acta Cardiologica</i> , 2009 , 64, 413-4; author reply 414	0.9	
130	NASH predicts plasma inflammatory biomarkers independently of visceral fat in men. <i>Obesity</i> , 2008 , 16, 1394-9	8	152

129	Diabetic retinopathy is associated with an increased incidence of cardiovascular events in Type 2 diabetic patients. <i>Diabetic Medicine</i> , 2008 , 25, 45-50	3.5	64
128	Relationship between soluble CD40 ligand and gamma-glutamyltransferase concentrations in non-drinking, young Type 1 diabetic individuals. <i>Diabetic Medicine</i> , 2008 , 25, 1283-8	3.5	1
127	Association between serum TSH, free T4 and serum liver enzyme activities in a large cohort of unselected outpatients. <i>Clinical Endocrinology</i> , 2008 , 68, 481-4	3.4	44
126	Immune tolerance with rituximab in congenital haemophilia with inhibitors: a systematic literature review based on individual patients' analysis. <i>Haemophilia</i> , 2008 , 14, 903-12	3.3	61
125	Help me, Doctor! My D-dimer is raised. <i>Annals of Medicine</i> , 2008 , 40, 594-605	1.5	66
124	Arm blood pressure index and lipoprotein(a) in renal transplant recipients. <i>Transplantation Proceedings</i> , 2008 , 40, 3499	1.1	
123	The metabolic syndrome and the risk of arterial and venous thrombosis. <i>Thrombosis Research</i> , 2008 , 122, 727-35	8.2	45
122	Inherited platelet disorders. <i>Clinica Chimica Acta</i> , 2008 , 387, 1-8	6.2	17
121	The paradoxical relationship between serum uric acid and cardiovascular disease. <i>Clinica Chimica Acta</i> , 2008 , 392, 1-7	6.2	162
120	Laboratory, clinical and therapeutic aspects of acquired hemophilia A. <i>Clinica Chimica Acta</i> , 2008 , 395, 14-8	6.2	51
119	Relationship between red blood cell distribution width and kidney function tests in a large cohort of unselected outpatients. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2008 , 68, 745-8	2	112
118	Sudden cardiac death: prevalence, pathogenesis, and prevention. <i>Annals of Medicine</i> , 2008 , 40, 360-75	1.5	24
117	Plasma D-dimer in the diagnosis of acute aortic dissection. <i>European Heart Journal</i> , 2008 , 29, 1207; author reply 1207-8	9.5	5
116	Association of inflammation with anaemia in patients with chronic kidney disease not requiring chronic dialysis. <i>Nephrology Dialysis Transplantation</i> , 2008 , 23, 2879-83	4.3	28
115	Detecting of chronic kidney disease in older people by the MDRD and MCQ formulas. <i>Age and Ageing</i> , 2008 , 37, 722	3	4
114	Prevalence of subclinical hypothyroidism in patients with chronic kidney disease. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2008 , 3, 1296-300	6.9	138
113	Increased risk of CKD among type 2 diabetics with nonalcoholic fatty liver disease. <i>Journal of the American Society of Nephrology: JASN</i> , 2008 , 19, 1564-70	12.7	143
112	Prevalence of folic Acid and vitamin B12 deficiencies in patients with thyroid disorders. <i>American Journal of the Medical Sciences</i> , 2008 , 336, 50-2	2.2	11

111	Plasma gamma-glutamyl transferase activity predicts homocysteine concentration in a large cohort of unselected outpatients. <i>Internal Medicine</i> , 2008 , 47, 705-7	1.1	6
110	Is liver fat detrimental to vessels?: intersections in the pathogenesis of NAFLD and atherosclerosis. <i>Clinical Science</i> , 2008 , 115, 1-12	6.5	57
109	Glucose challenge test does not predict gestational diabetes mellitus. <i>Internal Medicine</i> , 2008 , 47, 1171-4.1		1
108	Natriuretic peptides for assessing the prognosis of acute pulmonary embolism. <i>Chest</i> , 2008 , 133, 1531	5.3	1
107	The use of recombinant factor VIIa in liver diseases. <i>Blood Coagulation and Fibrinolysis</i> , 2008 , 19, 341-8	1	19
106	Relationship between thyroid status and renal function in a general population of unselected outpatients. <i>Clinical Biochemistry</i> , 2008 , 41, 625-7	3.5	24
105	Acquired factor VIII inhibitors in oncohematology: a systematic review. <i>Critical Reviews in Oncology/Hematology</i> , 2008 , 66, 194-9	7	44
104	Iron and thrombosis. <i>Annals of Hematology</i> , 2008 , 87, 167-73	3	84
103	Non-alcoholic fatty liver disease is independently associated with an increased prevalence of chronic kidney disease and proliferative/laser-treated retinopathy in type 2 diabetic patients. <i>Diabetologia</i> , 2008 , 51, 444-50	10.3	242
102	Increased risk of cardiovascular disease in non-alcoholic fatty liver disease: causal effect or epiphenomenon?. <i>Diabetologia</i> , 2008 , 51, 1947-53	10.3	342
101	Uric acid concentration in patient with acute coronary syndrome. <i>Internal and Emergency Medicine</i> , 2008 , 3, 409-11	3.7	3
100	The effect of iron depletion on chronic hepatitis C virus infection. <i>Hepatology International</i> , 2008 , 2, 335-8.1	4.6	44
99	Insulin effect on serum potassium and auto-inhibition of insulin secretion is intact in a patient with leprechaunism despite severe impairment of substrates metabolism. <i>Diabetes/Metabolism Research and Reviews</i> , 2008 , 24, 205-10	7.5	0
98	Variability of body weight, pulse pressure and glycaemia strongly predict total mortality in elderly type 2 diabetic patients. The Verona Diabetes Study. <i>Diabetes/Metabolism Research and Reviews</i> , 2008 , 24, 624-8	7.5	50
97	Effect of hemodialysis on traditional and innovative cardiac markers. <i>Journal of Clinical Laboratory Analysis</i> , 2008 , 22, 59-65	3	14
96	Relationship between serum vitamin D and inflammatory markers in the general population: comment on the article by Patel et al. <i>Arthritis and Rheumatism</i> , 2008 , 58, 913-4		3
95	RESPONSE: LETTER TO THE EDITOR. <i>PACE - Pacing and Clinical Electrophysiology</i> , 2008 , 31, 261-261	1.6	
94	Association between 25-hydroxyvitamin D deficiency and cardiovascular disease in type 2 diabetic patients with mild kidney dysfunction. <i>Nephrology Dialysis Transplantation</i> , 2008 , 23, 269-74	4.3	33

93	Relationship between mean platelet volume and biochemical components of the metabolic syndrome. <i>Clinical Drug Investigation</i> , 2007 , 27, 731-2; author reply 732-3	3.2	6
92	"Buffalo" hump in nonalcoholic fatty liver disease. <i>Hepatology</i> , 2007 , 46, 1311-2; author reply 1312	11.2	
91	Relationship between ABO blood group and von Willebrand factor levels: from biology to clinical implications. <i>Thrombosis Journal</i> , 2007 , 5, 14	5.6	109
90	Relationship between abnormal microvolt T-wave alternans and poor glycemic control in type 2 diabetic patients. <i>PACE - Pacing and Clinical Electrophysiology</i> , 2007 , 30, 1267-72	1.6	20
89	Non-alcoholic fatty liver disease, the metabolic syndrome and the risk of cardiovascular disease: the plot thickens. <i>Diabetic Medicine</i> , 2007 , 24, 1-6	3.5	170
88	Effect of moderate aerobic exercise on sympatho-vagal balance in Type 2 diabetic patients. <i>Diabetic Medicine</i> , 2007 , 24, 370-6	3.5	42
87	Relationship between lipoprotein(a) and fasting plasma glucose in the general population. <i>European Journal of Clinical Investigation</i> , 2007 , 37, 826-7	4.6	1
86	Relationship between lipoprotein(a) and thyroid function status in the general population. <i>Archives of Medical Research</i> , 2007 , 38, 905-6; author reply 907-8	6.6	5
85	Prophylaxis in von Willebrand disease. <i>Annals of Hematology</i> , 2007 , 86, 699-704	3	10
84	Pathogenesis, clinical and laboratory aspects of thrombosis in cancer. <i>Journal of Thrombosis and Thrombolysis</i> , 2007 , 24, 29-38	5.1	43
83	Alanine aminotransferase as an independent predictor of incident nonalcoholic fatty liver disease. <i>Clinical Chemistry</i> , 2007 , 53, 1159; author reply 1159-61	5.5	3
82	The role of iron in diabetes and its complications: response to Swaminathan et al. <i>Diabetes Care</i> , 2007 , 30, e132	14.6	3
81	Nonalcoholic fatty liver disease is independently associated with an increased incidence of cardiovascular events in type 2 diabetic patients. <i>Diabetes Care</i> , 2007 , 30, 2119-21	14.6	395
80	Prevalence of Nonalcoholic Fatty Liver Disease and Its Association With Cardiovascular Disease Among Type 2 Diabetic Patients: Response to Schindhelm, Heine, and Diamant. <i>Diabetes Care</i> , 2007 , 30, e95-e95	14.6	1
79	Retinopathy predicts cardiovascular mortality in type 2 diabetic men and women: response to Juutilainen et al. <i>Diabetes Care</i> , 2007 , 30, e51; author reply e52	14.6	3
78	Effect of serum gamma-glutamyltransferase and obesity on the risk of dyslipidemia and poor glycemic control in type 2 diabetic patients: cross-sectional findings from the Verona Diabetes Study. <i>Clinical Chemistry</i> , 2007 , 53, 1867-9; author reply 1869-70	5.5	5
77	Vitamin D deficiency among Italian children. <i>Cmaj</i> , 2007 , 177, 1529-30; author reply 1530	3.5	19
76	Reduced von Willebrand factor-cleaving protease levels in secondary thrombotic microangiopathies and other diseases. <i>Seminars in Thrombosis and Hemostasis</i> , 2007 , 33, 787-97	5.3	40

75	Pioglitazone in nonalcoholic steatohepatitis. <i>New England Journal of Medicine</i> , 2007 , 356, 1067; author reply 1068-9	59.2	18
74	Relationship between gamma-glutamyltransferase, fasting plasma glucose, and triglycerides in the general population. <i>Clinical Chemistry</i> , 2007 , 53, 1866-7; author reply 1869-70	5.5	9
73	Plasma PAI-1 levels are increased in patients with nonalcoholic steatohepatitis. <i>Diabetes Care</i> , 2007 , 30, e31-2	14.6	51
72	Has homocysteine shrunk?. <i>Clinical Chemistry and Laboratory Medicine</i> , 2007 , 45, 1419-20	5.9	1
71	Monitoring glycaemic control: is there evidence for appropriate use of routine measurement of glycated haemoglobin?. <i>Clinical Chemistry and Laboratory Medicine</i> , 2007 , 45, 1065-7	5.9	15
70	Lipoprotein(a), thrombophilia and venous thrombosis. <i>Acta Haematologica</i> , 2007 , 117, 246-7	2.7	6
69	Associations between serum 25-hydroxyvitamin D3 concentrations and liver histology in patients with non-alcoholic fatty liver disease. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2007 , 17, 517-24	4.5	294
68	The significance of evaluating conventional inflammatory markers in Von Willebrand factor measurement. <i>Clinica Chimica Acta</i> , 2007 , 381, 167-70	6.2	17
67	Relationship between gamma-glutamyltransferase, lipids and lipoprotein(a) in the general population. <i>Clinica Chimica Acta</i> , 2007 , 384, 163-6	6.2	20
66	Differences and similarities in early atherosclerosis between patients with non-alcoholic steatohepatitis and chronic hepatitis B and C. <i>Journal of Hepatology</i> , 2007 , 46, 1126-32	13.4	133
65	Non-alcoholic fatty liver disease as a determinant of cardiovascular disease. <i>Atherosclerosis</i> , 2007 , 190, 18-9; author reply 20-1	3.1	12
64	Non-alcoholic fatty liver disease and increased risk of cardiovascular disease. <i>Atherosclerosis</i> , 2007 , 191, 235-40	3.1	431
63	Ginkgo biloba, inflammation and lipoprotein(a). <i>Atherosclerosis</i> , 2007 , 195, 417-8; author reply 419-22	3.1	15
62	Prevalence of nonalcoholic fatty liver disease and its association with cardiovascular disease among type 2 diabetic patients. <i>Diabetes Care</i> , 2007 , 30, 1212-8	14.6	709
61	Platelets and lipoprotein(a) in retinal vein occlusion: Mutual targets for aspirin therapy. <i>Thrombosis and Haemostasis</i> , 2007 , 97, 1059-1060	7	4
60	Serum 25-hydroxyvitamin D3 concentrations and prevalence of cardiovascular disease among type 2 diabetic patients. <i>Diabetes Care</i> , 2006 , 29, 722-4	14.6	200
59	Relations between carotid artery wall thickness and liver histology in subjects with nonalcoholic fatty liver disease. <i>Diabetes Care</i> , 2006 , 29, 1325-30	14.6	312
58	Non-alcoholic fatty liver disease is associated with carotid artery wall thickness in diet-controlled type 2 diabetic patients. <i>Journal of Endocrinological Investigation</i> , 2006 , 29, 55-60	5.2	74

57	Visceral adipose tissue may mediate the link between non-alcoholic fatty liver disease and endocrine abnormalities. <i>Journal of Hepatology</i> , 2006 , 45, 454; author reply 454-6	13.4	3
56	Relationship between high-sensitivity C-reactive protein levels and liver histology in subjects with non-alcoholic fatty liver disease. <i>Journal of Hepatology</i> , 2006 , 45, 879-81; author reply 881-2	13.4	65
55	Effects of moderate-intensity exercise training on plasma biomarkers of inflammation and endothelial dysfunction in older patients with type 2 diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2006 , 16, 543-9	4.5	112
54	Inflammatory variables may mediate the link between low plasma vitamin B6 concentrations and cardiovascular disease in type 2 diabetes. <i>Nutrition, Metabolism and Cardiovascular Diseases</i> , 2006 , 16, e9-e10	4.5	1
53	Carotid atherosclerosis and rheumatoid arthritis. <i>Annals of Internal Medicine</i> , 2006 , 145, 231; author reply 231	8	
52	Measurement of microvolt T-wave alternans, a new arrhythmic risk stratification test, in Type 2 diabetic patients without clinical cardiovascular disease. <i>Diabetic Medicine</i> , 2006 , 23, 207-10	3.5	10
51	Increased prevalence of cardiovascular disease in Type 2 diabetic patients with non-alcoholic fatty liver disease. <i>Diabetic Medicine</i> , 2006 , 23, 403-9	3.5	125
50	The International Diabetes Federation definition of the metabolic syndrome independently predicts future cardiovascular events in Type 2 diabetic patients. The Valpolicella Heart Diabetes Study. <i>Diabetic Medicine</i> , 2006 , 23, 1270-1	3.5	9
49	Associations between liver histology and cortisol secretion in subjects with nonalcoholic fatty liver disease. <i>Clinical Endocrinology</i> , 2006 , 64, 337-41	3.4	67
48	Hypovitaminosis D among unselected medical inpatients and outpatients in Northern Italy. <i>Clinical Endocrinology</i> , 2006 , 64, 475	3.4	3
47	Associations between plasma adiponectin concentrations and liver histology in patients with nonalcoholic fatty liver disease. <i>Clinical Endocrinology</i> , 2006 , 64, 679-83	3.4	141
46	Serum 25-hydroxyvitamin D3 concentrations and carotid artery intima-media thickness among type 2 diabetic patients. <i>Clinical Endocrinology</i> , 2006 , 65, 593-7	3.4	198
45	Nonalcoholic fatty liver disease and risk of future cardiovascular events among type 2 diabetic patients. <i>Diabetes</i> , 2005 , 54, 3541-6	0.9	449
44	Increased plasma markers of inflammation and endothelial dysfunction and their association with microvascular complications in Type 1 diabetic patients without clinically manifest macroangiopathy. <i>Diabetic Medicine</i> , 2005 , 22, 999-1004	3.5	92
43	Relationship of non-alcoholic hepatic steatosis to cortisol secretion in diet-controlled Type 2 diabetic patients. <i>Diabetic Medicine</i> , 2005 , 22, 1146-50	3.5	18
42	Non-alcoholic hepatic steatosis and its relation to increased plasma biomarkers of inflammation and endothelial dysfunction in non-diabetic men. Role of visceral adipose tissue. <i>Diabetic Medicine</i> , 2005 , 22, 1354-8	3.5	134
41	Associations between liver histology and early carotid atherosclerosis in subjects with nonalcoholic fatty liver disease. <i>Hepatology</i> , 2005 , 42, 974-5; discussion 975	11.2	21
40	Nonalcoholic fatty liver disease and atherosclerosis. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2005 , 25, e117; author reply e117-8	9.4	2

39	Associations between liver histology and carotid intima-media thickness in patients with nonalcoholic fatty liver disease. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2005 , 25, 2687-8	9.4	25
38	Relation of nonalcoholic hepatic steatosis to early carotid atherosclerosis in healthy men: role of visceral fat accumulation. <i>Diabetes Care</i> , 2004 , 27, 2498-500	14.6	150
37	Hypoadiponectinemia is closely associated with nonalcoholic hepatic steatosis in obese subjects. <i>Diabetes Care</i> , 2004 , 27, 2085-6	14.6	27
36	Soluble CD40L in young type 1 diabetic individuals without clinical microvascular and macrovascular complications. <i>Diabetes Care</i> , 2004 , 27, 1236-7	14.6	8
35	Decreased plasma adiponectin concentrations are closely associated with nonalcoholic hepatic steatosis in obese individuals. <i>Clinical Endocrinology</i> , 2004 , 61, 700-3	3.4	93
34	Relationship of nonalcoholic hepatic steatosis to overnight low-dose dexamethasone suppression test in obese individuals. <i>Clinical Endocrinology</i> , 2004 , 61, 711-5	3.4	22
33	The Metabolic Syndrome is an independent predictor of cardiovascular disease in Type 2 diabetic subjects. Prospective data from the Verona Diabetes Complications Study. <i>Diabetic Medicine</i> , 2004 , 21, 52-8	3.5	213
32	Predictors of insulin sensitivity in Type 2 diabetes mellitus. <i>Diabetic Medicine</i> , 2002 , 19, 535-42	3.5	42
31	HOMA-estimated insulin resistance is an independent predictor of cardiovascular disease in type 2 diabetic subjects: prospective data from the Verona Diabetes Complications Study. <i>Diabetes Care</i> , 2002 , 25, 1135-41	14.6	390
30	Plasma total homocysteine levels are associated with von Willebrand factor, soluble intercellular adhesion molecule-1, and soluble tumor necrosis factor-alpha receptors in young type 1 diabetic patients without clinical evidence of macrovascular complications. <i>Diabetes Care</i> , 2001 , 24, 1496-7	14.6	8
29	Elevated plasma levels of soluble receptors of TNF-alpha and their association with smoking and microvascular complications in young adults with type 1 diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001 , 86, 3805-8	5.6	32
28	Elevated levels of interleukin-6 in young adults with type 1 diabetes without clinical evidence of microvascular and macrovascular complications. <i>Diabetes Care</i> , 2001 , 24, 956-7	14.6	55
27	Serum leptin concentrations in young smokers with type 1 diabetes. <i>Diabetes Care</i> , 2001 , 24, 793	14.6	7
26	Relation between soluble adhesion molecules and insulin sensitivity in type 2 diabetic individuals: role of adipose tissue. <i>Diabetes Care</i> , 2001 , 24, 1961-6	14.6	46
25	Intracellular partition of plasma glucose disposal in hypertensive and normotensive subjects with type 2 diabetes mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001 , 86, 2073-9	5.6	8
24	Homeostasis model assessment closely mirrors the glucose clamp technique in the assessment of insulin sensitivity: studies in subjects with various degrees of glucose tolerance and insulin sensitivity. <i>Diabetes Care</i> , 2000 , 23, 57-63	14.6	1906
23	Cigarette smoking and plasma total homocysteine levels in young adults with type 1 diabetes. <i>Diabetes Care</i> , 2000 , 23, 524-8	14.6	34
22	Effects of glucosamine infusion on insulin secretion and insulin action in humans. <i>Diabetes</i> , 2000 , 49, 926-35	0.9	122

21	Effect of chronic treatment with lacidipine or lisinopril on intracellular partitioning of glucose metabolism in type 2 diabetes mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1999 , 84, 1544-50	5.6	11
20	Chronic cigarette smoking is associated with increased plasma circulating intercellular adhesion molecule 1 levels in young type 1 diabetic patients. <i>Diabetes Care</i> , 1999 , 22, 1871-4	14.6	15
19	Prevalence of insulin resistance in metabolic disorders: the Bruneck Study. <i>Diabetes</i> , 1998 , 47, 1643-9	0.9	647
18	Intimal-medial thickness of the carotid artery in nondiabetic and NIDDM patients. Relationship with insulin resistance. <i>Diabetes Care</i> , 1997 , 20, 627-31	14.6	128
17	Cigarette smoking and insulin resistance in patients with noninsulin-dependent diabetes mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1997 , 82, 3619-24	5.6	134
16	Obesity worsens cardiovascular risk profiles independently of hyperinsulinaemia. <i>Journal of Internal Medicine</i> , 1997 , 241, 463-70	10.8	4
15	Relationship between fasting insulin and cardiovascular risk factors is already present in young men: the Verona Young Men Atherosclerosis Risk Factors Study. <i>European Journal of Clinical Investigation</i> , 1997 , 27, 248-54	4.6	10
14	Cigarette Smoking and Insulin Resistance in Patients with Noninsulin-Dependent Diabetes Mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1997 , 82, 3619-3624	5.6	107
13	Serum uric acid and related factors in 500 hospitalized subjects. <i>Metabolism: Clinical and Experimental</i> , 1996 , 45, 1557-61	12.7	62
12	Visceral fat accumulation and its relation to plasma hemostatic factors in healthy men. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 1996 , 16, 368-74	9.4	120
11	The white blood cell count: its relationship to plasma insulin and other cardiovascular risk factors in healthy male individuals. <i>Journal of Internal Medicine</i> , 1996 , 239, 435-41	10.8	59
10	Plasma factor VII and its relation to adipose tissue fatty acids and other atherogenic risk factors in healthy men. <i>European Journal of Clinical Investigation</i> , 1996 , 26, 247-53	4.6	6
9	Increase in circulating products of lipid peroxidation in smokers with IDDM. <i>Diabetes Care</i> , 1996 , 19, 1233-6	14.6	12
8	Liver Steatosis and Its Relation to Plasma Haemostatic Factors in Apparently Healthy Men - Role of the Metabolic Syndrome. <i>Thrombosis and Haemostasis</i> , 1996 , 76, 069-073	7	63
7	Ultrasonographic intra-abdominal depth and its relation to haemostatic factors in healthy males 1996 , 20, 882-5		1
6	Fasting serum insulin in relation to components of the metabolic syndrome in European healthy men: the European Fat Distribution Study. <i>Metabolism: Clinical and Experimental</i> , 1995 , 44, 35-40	12.7	39
5	Relationships of blood pressure to fibrinolysis: influence of anthropometry, metabolic profile and behavioural variables. <i>Journal of Hypertension</i> , 1995 , 13, 659-66	1.9	19
4	Plasma fibrinogen in relation to serum insulin, smoking habits and adipose tissue fatty acids in healthy men. <i>European Journal of Clinical Investigation</i> , 1994 , 24, 126-30	4.6	23

3	Relationships of plasminogen activator inhibitor-1 to anthropometry, serum insulin, triglycerides and adipose tissue fatty acids in healthy men. <i>Atherosclerosis</i> , 1994 , 106, 139-47	3.1	32
2	From nonalcoholic fatty liver disease to metabolic dysfunction-associated fatty liver disease: is it time for a change of terminology?. <i>Hepatoma Research</i> , 2020,	4.3	3
1	NAFLD and Cardiovascular Risk Factors: Implications for Vascular Disease 71-83		