

Nazareno Suardi

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4469497/nazareno-suardi-publications-by-year.pdf>

Version: 2024-04-17

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

267
papers

10,277
citations

56
h-index

89
g-index

434
ext. papers

12,080
ext. citations

3.7
avg. IF

5.52
L-index

#	Paper	IF	Citations
267	Radiation therapy after radical prostatectomy is associated with higher other-cause mortality.. <i>Cancer Causes and Control</i> , 2022 , 33, 769	2.8	
266	Comment on: "The surgical learning curve for salvage robot-assisted radical prostatectomy: a prospective single-surgeon study". <i>Minerva Urology and Nephrology</i> , 2021 , 73, 680-682	2.3	
265	Response to Re: External beam radiotherapy and radical prostatectomy are associated with better survival in Asian prostate cancer patients. <i>International Journal of Urology</i> , 2021 ,	2.3	2
264	Improving the stratification of intermediate risk prostate cancer. <i>Minerva Urology and Nephrology</i> , 2021 ,	2.3	3
263	Re: Sophie Knipper, Luigi Ascalone, Benjamin Ziegler, et al. Salvage Surgery in Patients with Local Recurrence After Radical Prostatectomy. <i>Eur Urol</i> 2021;79:537-44: Surgical Treatment of Local Recurrence Following Radical Prostatectomy: Reality or Illusion?. <i>European Urology</i> , 2021 , 79, e132-e133	10.2	
262	Presence of biopsy Gleason pattern 5 + 3 is associated with higher mortality after radical prostatectomy but not after external beam radiotherapy compared to other Gleason Grade Group IV patterns. <i>Prostate</i> , 2021 , 81, 778-784	4.2	1
261	External beam radiotherapy and radical prostatectomy are associated with better survival in Asian prostate cancer patients. <i>International Journal of Urology</i> , 2021 ,	2.3	4
260	Bladder cancer histological variants: which parameters could predict the concordance between transurethral resection of bladder tumor and radical cystectomy specimens?. <i>Central European Journal of Urology</i> , 2021 , 74, 355-361	0.9	0
259	Sexuality during COVID lockdown: a cross-sectional Italian study among hospital workers and their relatives. <i>International Journal of Impotence Research</i> , 2021 , 33, 131-136	2.3	13
258	Re: Ming-Chun Chan, Sharon E.K. Yeo, Yew-Lam Chong, Yee-Mun Lee. Stepping Forward: Urologists' Efforts During the COVID-19 Outbreak in Singapore. <i>Eur Urol</i> 2020;78:e38-9. <i>European Urology</i> , 2020 , 78, e42	10.2	7
257	Hospital care in Departments defined as COVID-free: A proposal for a safe hospitalization protecting healthcare professionals and patients not affected by COVID-19. <i>Archivio Italiano Di Urologia Andrologia</i> , 2020 , 92,	1.6	8
256	ABLATE: a score to predict complications and recurrence rate in percutaneous treatments of renal lesions. <i>Medical Oncology</i> , 2020 , 37, 26	3.7	7
255	Port-site metastasis and atypical recurrences after robotic-assisted radical cystectomy (RARC): an updated comprehensive and systematic review of current evidences. <i>Journal of Robotic Surgery</i> , 2020 , 14, 805-812	2.9	6
254	Long-term Outcomes of Salvage Lymph Node Dissection for Nodal Recurrence of Prostate Cancer After Radical Prostatectomy: Not as Good as Previously Thought. <i>European Urology</i> , 2020 , 78, 661-669	10.2	30
253	Assessing the Best Surgical Template at Salvage Pelvic Lymph Node Dissection for Nodal Recurrence of Prostate Cancer After Radical Prostatectomy: When Can Bilateral Dissection be Omitted? Results from a Multi-institutional Series. <i>European Urology</i> , 2020 , 78, 779-782	10.2	6
252	Re: Reconsidering Prostate Cancer Mortality - The Future of PSA Screening: The future of PSA screening by reconsidering prostate cancer mortality. <i>European Urology</i> , 2020 , 78, 929	10.2	
251	Minimally invasive strategies for the treatment of prostate cancer recurrence after radiation therapy: a systematic review. <i>Minerva Urologica E Nefrologica = the Italian Journal of Urology and Nephrology</i> , 2020 , 72, 563-578	4.4	4

250	Urology apps: overview of current types and use. <i>Central European Journal of Urology</i> , 2020 , 73, 369-372.	0.9	1
249	Reply by Authors. <i>Journal of Urology</i> , 2020 , 204, 302	2.5	
248	Underestimation of Positron Emission Tomography/Computerized Tomography in Assessing Tumor Burden in Prostate Cancer Nodal Recurrence: Head-to-Head Comparison of Ga-PSMA and C-Choline in a Large, Multi-Institutional Series of Extended Salvage Lymph Node Dissections. <i>Journal of Urology</i> , 2020 , 204, 296-302	2.5	17
247	Re: Giuseppe Simone, Umberto Anceschi, Gabriele Tuderti, et al. Robot-assisted Partial Adrenalectomy for the Treatment of Conn Syndrome: Surgical Technique, and Perioperative and Functional Outcomes. <i>Eur Urol</i> 2019;75:811-6. <i>European Urology</i> , 2019 , 76, e142-e143	10.2	
246	Robotic surgery in patients with achondroplastic dwarfism: evaluation of risks and issues in an anatomical challenging bilateral partial nephrectomy. <i>Journal of Robotic Surgery</i> , 2019 , 13, 783-786	2.9	1
245	Re: Active Surveillance of Prostate Cancer is a Viable Option in Men Younger than 60 Years. <i>European Urology</i> , 2019 , 76, 404	10.2	
244	Development of a New Comorbidity Assessment Tool for Specific Prediction of Perioperative Mortality in Contemporary Patients Treated with Radical Cystectomy. <i>Annals of Surgical Oncology</i> , 2019 , 26, 1942-1949	3.1	7
243	En Bloc HoLEP with early apical release in men with benign prostatic hyperplasia. <i>World Journal of Urology</i> , 2019 , 37, 2451-2458	4	29
242	Impact of multiparametric MRI and MRI-targeted biopsy on pre-therapeutic risk assessment in prostate cancer patients candidate for radical prostatectomy. <i>World Journal of Urology</i> , 2019 , 37, 221-234	4	19
241	Robotic laparoendoscopic single-site radical prostatectomy (R-LESS-RP) with daVinci Single-Site platform. Concept and evolution of the technique following an IDEAL phase 1. <i>Journal of Robotic Surgery</i> , 2019 , 13, 215-226	2.9	10
240	Robotic-assisted laparoscopic pyeloplasty with the use of the Contour3 stent: description of the technique and analysis of outcomes after the first 30 cases. <i>Central European Journal of Urology</i> , 2019 , 72, 51-53	0.9	4
239	The Impact of Experience on the Risk of Surgical Margins and Biochemical Recurrence after Robot-Assisted Radical Prostatectomy: A Learning Curve Study. <i>Journal of Urology</i> , 2019 , 202, 108-113	2.5	40
238	Technologies for image-guided surgery for managing lymphatic metastases in prostate cancer. <i>Nature Reviews Urology</i> , 2019 , 16, 159-171	5.5	34
237	Identifying the Optimal Candidate for Salvage Lymph Node Dissection for Nodal Recurrence of Prostate Cancer: Results from a Large, Multi-institutional Analysis. <i>European Urology</i> , 2019 , 75, 176-183	10.2	60
236	Increasing Rate of Noninterventional Treatment Management in Localized Prostate Cancer Candidates for Active Surveillance: A North American Population-Based Study. <i>Clinical Genitourinary Cancer</i> , 2019 , 17, 72-78.e4	3.3	5
235	Neoadjuvant and adjuvant treatment in high-risk prostate cancer. <i>Expert Review of Clinical Pharmacology</i> , 2018 , 11, 425-438	3.8	8
234	Identifying candidates for super-extended staging pelvic lymph node dissection among patients with high-risk prostate cancer. <i>BJU International</i> , 2018 , 121, 421-427	5.6	17
233	Perioperative and Oncologic Outcomes of Nephrectomy and Caval Thrombectomy Using Extracorporeal Circulation and Deep Hypothermic Circulatory Arrest for Renal Cell Carcinoma Invading the Supradiaphragmatic Inferior Vena Cava and/or Right Atrium. <i>European Urology</i> , 2018 , 73, 783-789	10.2	19

232 Surgical Treatment for LUTS/BPH: Laser Devices **2018**, 257-288

231 Cytoreductive Radical Prostatectomy in Men with Prostate Cancer and Skeletal Metastases. *European Urology Oncology*, **2018**, 1, 46-53 6.7 26

230 Association Between Prostate Imaging Reporting and Data System (PI-RADS) Score for the Index Lesion and Multifocal, Clinically Significant Prostate Cancer. *European Urology Oncology*, **2018**, 1, 29-36 6.7 26

229 A proposal of a new nomogram for predicting upstaging in contemporary DAmico low-risk prostate cancer patients. *World Journal of Urology*, **2017**, 35, 189-197 4 12

228 Robot-assisted Radical Prostatectomy and Extended Pelvic Lymph Node Dissection in Patients with Locally-advanced Prostate Cancer. *European Urology*, **2017**, 71, 249-256 10.2 50

227 Impact of Postoperative Radiotherapy in Men with Persistently Elevated Prostate-specific Antigen After Radical Prostatectomy for Prostate Cancer: A Long-term Survival Analysis. *European Urology*, **2017**, 72, 910-917 10.2 17

226 Evaluating the effect of time from prostate cancer diagnosis to radical prostatectomy on cancer control: Can surgery be postponed safely?. *Urologic Oncology: Seminars and Original Investigations*, **2017**, 35, 150.e9-150.e15 2.8 29

225 New surgical approaches for clinically high-risk or metastatic prostate cancer. *Expert Review of Anticancer Therapy*, **2017**, 17, 1013-1031 3.5 5

224 Assessing the Impact of Surgeon Experience on Urinary Continence Recovery After Robot-Assisted Radical Prostatectomy: Results of Four High-Volume Surgeons. *Journal of Endourology*, **2017**, 31, 872-877 7.7 30

223 The New Prostate Cancer Grading System Does Not Improve Prediction of Clinical Recurrence After Radical Prostatectomy: Results of a Large, Two-Center Validation Study. *Prostate*, **2017**, 77, 263-273 4.2 18

222 When to perform preoperative chest computed tomography for renal cancer staging. *BJU International*, **2017**, 120, 490-496 5.6 4

221 Robot-assisted Salvage Lymph Node Dissection for Clinically Recurrent Prostate Cancer. *European Urology*, **2017**, 72, 432-438 10.2 32

220 Early Postoperative Radiotherapy is Associated with Worse Functional Outcomes in Patients with Prostate Cancer. *Journal of Urology*, **2017**, 197, 669-675 2.5 39

219 Reply to Marc A. Bjurlin, Lee C. Zhao, and Michael D. Stifelman Letter to the Editor Re: Nicolò Maria Buffi, Giovanni Lughezzani, Rodolfo Hurle, et al. Robot-assisted Surgery for Benign Ureteral Strictures: Experience and Outcomes from Four Tertiary Care Institutions. *Eur Urol*. In press. <http://dx.doi.org/10.1016/j.eururo.2016.07.022>. *European Urology*, **2017**, 71, e92-e93

218 Incidence and Predictors of 30-Day Readmission After Robot-Assisted Radical Prostatectomy. *Clinical Genitourinary Cancer*, **2017**, 15, 67-71 3.3 12

217 Robot-assisted Surgery for Benign Ureteral Strictures: Experience and Outcomes from Four Tertiary Care Institutions. *European Urology*, **2017**, 71, 945-951 10.2 41

216 A robotic needle driver to facilitate vesico-urethral anastomosis during laparoscopic radical prostatectomy. *Urologia*, **2017**, 0 1.2 1

215 Pelvic lymph node dissection in prostate cancer: indications, extent and tailored approaches. *Urologia*, **2017**, 84, 9-19 1.2 20

214	Robotic Radical Cystectomy and Urinary Diversions: Step-by-Step Technique 2017 , 683-693		
213	The role of cytoreductive radical prostatectomy (cRP) in men with hormone-sensitive, metastatic prostate cancer (mPCA).. <i>Journal of Clinical Oncology</i> , 2017 , 35, 241-241	2.2	
212	Robot assisted lymphadenectomy in urology: pelvic, retroperitoneal and inguinal. <i>Minerva Urology and Nephrology</i> , 2017 , 69, 38-55	2.3	10
211	Evaluation of positive surgical margins in patients undergoing robot-assisted and open radical prostatectomy according to preoperative risk groups. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2016 , 34, 57.e1-7	2.8	20
210	Rationale for local treatment in the management of metastatic prostate cancer. <i>Current Opinion in Supportive and Palliative Care</i> , 2016 , 10, 266-72	2.6	5
209	Re: Acute Toxicity and Quality of Life After Dose-intensified Salvage Radiation Therapy for Biochemically Recurrent Prostate Cancer After Prostatectomy: First Results of the Randomized Trial SAKK 09/10. <i>European Urology</i> , 2016 , 69, 966	10.2	1
208	A Multi-institutional Analysis of Perioperative Outcomes in 106 Men Who Underwent Radical Prostatectomy for Distant Metastatic Prostate Cancer at Presentation. <i>European Urology</i> , 2016 , 69, 788-94	10.2	99
207	The Role of Prostate-specific Antigen Persistence After Radical Prostatectomy for the Prediction of Clinical Progression and Cancer-specific Mortality in Node-positive Prostate Cancer Patients. <i>European Urology</i> , 2016 , 69, 1142-8	10.2	38
206	Potential Effect of Antiplatelet and Anticoagulant Therapy on the Timing of the Diagnosis of Bladder Cancer. <i>Clinical Genitourinary Cancer</i> , 2016 , 14, e245-50	3.3	4
205	Predicting survival of men with recurrent prostate cancer after radical prostatectomy. <i>European Journal of Cancer</i> , 2016 , 54, 27-34	7.5	25
204	A nomogram predicting the cancer-specific mortality in patients eligible for radical cystectomy evaluating clinical data and neoadjuvant cisplatinum-based chemotherapy. <i>World Journal of Urology</i> , 2016 , 34, 207-13	4	10
203	Non-surgically related causes of erectile dysfunction after bilateral nerve-sparing radical prostatectomy. <i>Prostate Cancer and Prostatic Diseases</i> , 2016 , 19, 185-90	6.2	12
202	Current Standard Technique for Modern Flexible Ureteroscopy: Tips and Tricks. <i>European Urology</i> , 2016 , 70, 188-194	10.2	72
201	STAMPEDE trial and patients with non-metastatic prostate cancer. <i>Lancet, The</i> , 2016 , 388, 234-5	40	3
200	What is the optimal definition of misclassification in patients with very low-risk prostate cancer eligible for active surveillance? Results from a multi-institutional series. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015 , 33, 164.e1-9	2.8	26
199	Patterns of Clinical Recurrence of Node-positive Prostate Cancer and Impact on Long-term Survival. <i>European Urology</i> , 2015 , 68, 777-84	10.2	41
198	A Detailed Analysis of the Association Between Postoperative Phosphodiesterase Type 5 Inhibitor Use and the Risk of Biochemical Recurrence After Radical Prostatectomy. <i>European Urology</i> , 2015 , 68, 750-3	10.2	28
197	Predicting Life Expectancy in Men Diagnosed with Prostate Cancer. <i>European Urology</i> , 2015 , 68, 756-65	10.2	43

196	Long-term oncologic outcomes of laparoscopic renal cryoablation as primary treatment for small renal masses. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015 , 33, 22.e1-22.e9	2.8	38
195	Long-term outcomes of salvage lymph node dissection for clinically recurrent prostate cancer: results of a single-institution series with a minimum follow-up of 5 years. <i>European Urology</i> , 2015 , 67, 299-309	10.2	158
194	More extensive pelvic lymph node dissection improves survival in patients with node-positive prostate cancer. <i>European Urology</i> , 2015 , 67, 212-9	10.2	134
193	Importance of prostate volume in the stratification of patients with intermediate-risk prostate cancer. <i>International Journal of Urology</i> , 2015 , 22, 555-61	2.3	11
192	Identification of pathologically favorable disease in intermediate-risk prostate cancer patients: Implications for active surveillance candidates selection. <i>Prostate</i> , 2015 , 75, 1484-91	4.2	19
191	Reply to Berardino De Bari, Stefano Arcangeli, and Filippo Alongi letter to the editor re: Nazareno Suardi, Andrea Gallina, Giuliana Lista, et al. impact of adjuvant radiation therapy on urinary continence recovery after radical prostatectomy. <i>Eur Urol</i> 2014;65:546-51. <i>European Urology</i> , 2015 , 67, 676-678	10.2	
190	Oncological predictive value of the 2004 World Health Organisation grading classification in primary T1 non-muscle-invasive bladder cancer. A step forward or back?. <i>BJU International</i> , 2015 , 115, 267-73	5.6	18
189	Penile rehabilitation after radical prostatectomy: does it work?. <i>Translational Andrology and Urology</i> , 2015 , 4, 110-23	2.3	24
188	Impact of adjuvant radiation therapy on urinary continence recovery after radical prostatectomy. <i>European Urology</i> , 2014 , 65, 546-51	10.2	68
187	Head-to-head comparison of lymph node density and number of positive lymph nodes in stratifying the outcome of patients with lymph node-positive prostate cancer submitted to radical prostatectomy and extended lymph node dissection. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 29-34	2.8	20
186	Higher-than-expected severe (Grade 3-4) late urinary toxicity after postprostatectomy hypofractionated radiotherapy: a single-institution analysis of 1176 patients. <i>European Urology</i> , 2014 , 66, 1024-30	10.2	70
185	The number of cores at first biopsy may suggest the need for a confirmatory biopsy in patients eligible for active surveillance-implication for clinical decision making in the real-life setting. <i>Urology</i> , 2014 , 84, 634-41	1.6	6
184	How to optimize patient selection for robot-assisted radical prostatectomy: functional outcome analyses from a tertiary referral center. <i>Journal of Endourology</i> , 2014 , 28, 792-800	2.7	19
183	The key role of time in predicting progression-free survival in patients with renal cell carcinoma treated with partial or radical nephrectomy: conditional survival analysis. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 43.e9-16	2.8	10
182	MP70-15 LONG TERM OUTCOMES OF SALVAGE LYMPH NODE DISSECTION FOR CLINICALLY RECURRENT PROSTATE CANCER: RESULTS OF A SINGLE INSTITUTION SERIES WITH A MINIMUM FOLLOW-UP OF 5 YEARS. <i>Journal of Urology</i> , 2014 , 191,	2.5	6
181	External validation of the European association of urology recommendations for pelvic lymph node dissection in patients treated with robot-assisted radical prostatectomy. <i>Journal of Endourology</i> , 2014 , 28, 416-23	2.7	29
180	Preoperative erectile function is the only predictor of the use of a high number of phosphodiesterase type-5 inhibitors after bilateral nerve-sparing radical prostatectomy. <i>International Journal of Impotence Research</i> , 2014 , 26, 201-4	2.3	8
179	Extent of lymph node dissection at nephrectomy affects cancer-specific survival and metastatic progression in specific sub-categories of patients with renal cell carcinoma (RCC). <i>BJU International</i> , 2014 , 114, 210-5	5.6	51

178	Impact of adjuvant radiotherapy on survival of patients with node-positive prostate cancer. <i>Journal of Clinical Oncology</i> , 2014 , 32, 3939-47	2.2	184
177	Prevalence and predictors of concomitant low sexual desire/interest and new-onset erectile dysfunction - a picture from the everyday clinical practice. <i>Andrology</i> , 2014 , 2, 702-8	4.2	9
176	Predicting the risk of bone metastasis in prostate cancer. <i>Cancer Treatment Reviews</i> , 2014 , 40, 3-11	14.4	43
175	Utility of [11C]choline PET/CT in guiding lesion-targeted salvage therapies in patients with prostate cancer recurrence localized to a single lymph node at imaging: results from a pathologically validated series. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2014 , 32, 38.e9-16	2.8	54
174	Salvage therapy of small volume prostate cancer nodal failures: a review of the literature. <i>Critical Reviews in Oncology/Hematology</i> , 2014 , 90, 24-35	7	22
173	Indication for and extension of pelvic lymph node dissection during robot-assisted radical prostatectomy: an analysis of five European institutions. <i>European Urology</i> , 2014 , 66, 635-43	10.2	38
172	Predicting survival of patients with node-positive prostate cancer following multimodal treatment. <i>European Urology</i> , 2014 , 65, 554-62	10.2	61
171	Fascial Layers in Nerve Sparing Robot-Assisted Radical Prostatectomy. <i>Urology Practice</i> , 2014 , 1, 86-91	0.8	1
170	Re: Prasanna Sooriakumaran, Abhishek Srivastava, Shahrokh F. Shariat, et al. A multinational, multi-institutional study comparing positive surgical margin rates among 22,393 open, laparoscopic, and robot-assisted radical prostatectomy patients. <i>Eur Urol</i> . In press. http://dx.doi.org/10.1016/j.eururo.2013.11.018 . <i>European Urology</i> , 2014 , 65, e89-90	10.2	0
169	Impact of preoperative thrombocytosis on pathological outcomes and survival in patients treated with radical cystectomy for bladder carcinoma. <i>Anticancer Research</i> , 2014 , 34, 3225-30	2.3	31
168	Radical prostatectomy represents an effective treatment in patients with specimen-confined high pathological Gleason score prostate cancer. <i>BJU International</i> , 2013 , 111, 723-30	5.6	20
167	Staging lymphadenectomy in renal cell carcinoma must be extended: a sensitivity curve analysis. <i>BJU International</i> , 2013 , 111, 412-8	5.6	15
166	Nerve-sparing approach during radical prostatectomy is strongly associated with the rate of postoperative urinary continence recovery. <i>BJU International</i> , 2013 , 111, 717-22	5.6	73
165	When to perform lymph node dissection in patients with renal cell carcinoma: a novel approach to the preoperative assessment of risk of lymph node invasion at surgery and of lymph node progression during follow-up. <i>BJU International</i> , 2013 , 112, E59-66	5.6	35
164	Preoperative sex steroids are significant predictors of early biochemical recurrence after radical prostatectomy. <i>World Journal of Urology</i> , 2013 , 31, 275-80	4	25
163	Long-term evaluation of survival, continence and potency (SCP) outcomes after robot-assisted radical prostatectomy (RARP). <i>BJU International</i> , 2013 , 112, 338-45	5.6	32
162	Assessing the most accurate formula to predict the risk of lymph node metastases from prostate cancer in contemporary patients treated with radical prostatectomy and extended pelvic lymph node dissection. <i>Radiotherapy and Oncology</i> , 2013 , 109, 211-6	5.3	15
161	Reply to Manish Garg, Apul Goel and Jai Prakash letter to the editor re: Renzo Colombo, Lorenzo Rocchini, Nazareno Suardi, et al. Neoadjuvant short-term intensive intravesical mitomycin C regimen compared with weekly schedule for low-grade recurrent non-muscle-invasive bladder cancer: preliminary results of a randomised phase 2 study. <i>Eur Urol</i> 2012;62:797-802. <i>European Urology</i> , 2013 , 63, e7-8	10.2	

160	Postoperative phosphodiesterase type 5 inhibitor administration increases the rate of urinary continence recovery after bilateral nerve-sparing radical prostatectomy. <i>International Journal of Urology</i> , 2013 , 20, 413-9	2.3	17
159	774 HEAD-TO-HEAD COMPARISON OF LYMPH NODE DENSITY AND NUMBER OF POSITIVE LYMPH NODES IN STRATIFYING THE OUTCOME OF PATIENTS WITH LYMPH NODE POSITIVE PROSTATE CANCER SUBMITTED TO RADICAL PROSTATECTOMY AND EXTENDED PELVIC LYMPH NODE DISSECTION. <i>Journal of Urology</i> , 2013 , 189,	2.5	1
158	The number of cores taken in patients diagnosed with a single microfocus at initial biopsy is a major predictor of insignificant prostate cancer. <i>Journal of Urology</i> , 2013 , 189, 854-9	2.5	11
157	772 SALVAGE LYMPH NODE DISSECTION FOR PATIENTS TREATED WITH RADICAL PROSTATECTOMY WITH BIOCHEMICAL RECURRENCE AND IMAGING-DETECTED NODAL METASTASES. <i>Journal of Urology</i> , 2013 , 189,	2.5	1
156	Survival following biochemical recurrence after radical prostatectomy and adjuvant radiotherapy in patients with prostate cancer: the impact of competing causes of mortality and patient stratification. <i>European Urology</i> , 2013 , 64, 557-64	10.2	33
155	1838 WHEN TO PERFORM LYMPH NODE DISSECTION IN RENAL CELL CARCINOMA PATIENTS: A NOVEL APPROACH TO PREOPERATIVELY ASSESS THE RISK OF LYMPH NODE INVASION AT SURGERY AND NODAL PROGRESSION DURING FOLLOW UP. <i>Journal of Urology</i> , 2013 , 189,	2.5	1
154	1831 THE NUMBER OF LYMPH NODES REMOVED IN RENAL CELL CARCINOMA DOES AFFECT CANCER SPECIFIC SURVIVAL IN SPECIFIC SUBGROUPS OF PATIENTS: RESULTS FROM A SYSTEMATIC ANALYSIS. <i>Journal of Urology</i> , 2013 , 189,	2.5	2
153	Presence of positive surgical margin in patients with organ-confined prostate cancer equals to extracapsular extension negative surgical margin. A plea for TNM staging system reclassification. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2013 , 31, 1497-503	2.8	11
152	963 DEVELOPMENT AND INTERNAL VALIDATION OF A NOMOGRAM PREDICTING BIOCHEMICAL RECURRENCE AFTER EARLY SALVAGE RADIOTHERAPY IN PROSTATE CANCER PATIENTS TREATED WITH RADICAL PROSTATECTOMY. <i>Journal of Urology</i> , 2013 , 189,	2.5	2
151	Selecting the optimal candidate for adjuvant radiotherapy after radical prostatectomy for prostate cancer: a long-term survival analysis. <i>European Urology</i> , 2013 , 63, 998-1008	10.2	88
150	Circulating sex steroids and prostate cancer: introducing the time-dependency theory. <i>World Journal of Urology</i> , 2013 , 31, 267-73	4	5
149	The European Association of Urology Robotic Urology Section (ERUS) survey of robot-assisted radical prostatectomy (RARP). <i>BJU International</i> , 2013 , 111, 596-603	5.6	26
148	Impact of the introduction of a robotic training programme on prostate cancer stage migration at a single tertiary referral centre. <i>BJU International</i> , 2013 , 111, 1222-30	5.6	28
147	A novel tool to assess the risk of urinary incontinence after nerve-sparing radical prostatectomy. <i>BJU International</i> , 2013 , 111, 905-13	5.6	13
146	Extended pelvic lymph node dissection in prostate cancer: a 20-year audit in a single center. <i>Annals of Oncology</i> , 2013 , 24, 1459-66	10.3	65
145	Effect of number and location of distant metastases on renal cell carcinoma mortality in candidates for cytoreductive nephrectomy: implications for multimodal therapy. <i>International Journal of Urology</i> , 2013 , 20, 572-9	2.3	19
144	One patient out of four with newly diagnosed erectile dysfunction is a young man--worrisome picture from the everyday clinical practice. <i>Journal of Sexual Medicine</i> , 2013 , 10, 1833-41	1.1	79
143	Spatial distribution of positive cores improves the selection of patients with low-risk prostate cancer as candidates for active surveillance. <i>BJU International</i> , 2013 , 112, E234-42	5.6	1

142	Updated nomogram predicting lymph node invasion in patients with prostate cancer undergoing extended pelvic lymph node dissection: the essential importance of percentage of positive cores. <i>European Urology</i> , 2012 , 61, 480-7	10.2	425
141	Performance characteristics of computed tomography in detecting lymph node metastases in contemporary patients with prostate cancer treated with extended pelvic lymph node dissection. <i>European Urology</i> , 2012 , 61, 1132-8	10.2	95
140	Prediction of functional outcomes after nerve-sparing radical prostatectomy: results of conditional survival analyses. <i>European Urology</i> , 2012 , 62, 42-52	10.2	54
139	Choosing the best candidates for penile rehabilitation after bilateral nerve-sparing radical prostatectomy. <i>Journal of Sexual Medicine</i> , 2012 , 9, 608-17	1.1	29
138	Erectile function outcome after bilateral nerve sparing radical prostatectomy: which patients may be left untreated?. <i>Journal of Sexual Medicine</i> , 2012 , 9, 903-8	1.1	29
137	Extended pelvic lymph node dissection does not affect erectile function recovery in patients treated with bilateral nerve-sparing radical prostatectomy. <i>Journal of Sexual Medicine</i> , 2012 , 9, 2187-94	1.1	17
136	Influence of obesity on tumour volume in patients with prostate cancer. <i>BJU International</i> , 2012 , 109, 678-84	5.6	22
135	Unilateral positive biopsies in low risk prostate cancer patients diagnosed with extended transrectal ultrasound-guided biopsy schemes do not predict unilateral prostate cancer at radical prostatectomy. <i>BJU International</i> , 2012 , 110, E64-8	5.6	27
134	Age-adjusted validation of the most stringent criteria for active surveillance in low-risk prostate cancer patients. <i>Cancer</i> , 2012 , 118, 973-80	6.4	22
133	Preoperative circulating sex hormones are not predictors of positive surgical margins at open radical prostatectomy. <i>World Journal of Urology</i> , 2012 , 30, 533-9	4	6
132	Neoadjuvant short-term intensive intravesical mitomycin C regimen compared with weekly schedule for low-grade recurrent non-muscle-invasive bladder cancer: preliminary results of a randomised phase 2 study. <i>European Urology</i> , 2012 , 62, 797-802	10.2	26
131	Indications for pelvic nodal treatment in prostate cancer should change. Validation of the Roach formula in a large extended nodal dissection series. <i>International Journal of Radiation Oncology Biology Physics</i> , 2012 , 83, 624-9	4	20
130	Preoperative erectile function represents a significant predictor of postoperative urinary continence recovery in patients treated with bilateral nerve sparing radical prostatectomy. <i>Journal of Urology</i> , 2012 , 187, 569-74	2.5	30
129	982 STAGING LYMPHADENECTOMY IN RENAL CELL CARCINOMA MUST BE EXTENDED: A SENSITIVITY CURVE ANALYSES. <i>Journal of Urology</i> , 2012 , 187,	2.5	1
128	771 SHOULD AN EXTENDED NODAL TEMPLATE FOR HIGH RISK PROSTATE CANCER ALWAYS INCLUDE REMOVAL OF COMMON ILIAC LYMPH NODES?. <i>Journal of Urology</i> , 2012 , 187,	2.5	1
127	Long-term biochemical recurrence rates after robot-assisted radical prostatectomy: analysis of a single-center series of patients with a minimum follow-up of 5 years. <i>Urology</i> , 2012 , 79, 133-8	1.6	60
126	National Comprehensive Cancer Network practice guidelines 2011: Need for more accurate recommendations for pelvic lymph node dissection in prostate cancer. <i>Journal of Urology</i> , 2012 , 188, 423-8	2.5	19
125	The extent of lymphadenectomy does affect cancer specific survival in pathologically confirmed T4 renal cell carcinoma. <i>Urologia</i> , 2012 , 79, 109-15	1.2	17

124	Lymphatic spread of nodal metastases in high-risk prostate cancer: The ascending pathway from the pelvis to the retroperitoneum. <i>Prostate</i> , 2012 , 72, 186-92	4.2	58
123	Assessing the risk of lymph node invasion in patients with intermediate risk prostate cancer treated with extended pelvic lymph node dissection. A novel prediction tool. <i>Prostate</i> , 2012 , 72, 499-506	4.2	10
122	There is no way to identify patients who will harbor small volume, unilateral prostate cancer at final pathology. implications for focal therapies. <i>Prostate</i> , 2012 , 72, 925-30	4.2	11
121	Optimizing postoperative sexual function after radical prostatectomy. <i>Therapeutic Advances in Urology</i> , 2012 , 4, 347-65	3.2	12
120	Serum sex steroids depict a nonlinear u-shaped association with high-risk prostate cancer at radical prostatectomy. <i>Clinical Cancer Research</i> , 2012 , 18, 3648-57	12.9	53
119	Prostate saturation biopsy following a first negative biopsy: state of the art. <i>Urologia Internationalis</i> , 2012 , 89, 126-35	1.9	20
118	Prostate cancer in 2010: GSU: misclassification or biological progression?. <i>Nature Reviews Urology</i> , 2011 , 8, 65-6	5.5	2
117	1408 A NEW MULTIMODAL ANESTHESIOLOGICAL AND NUTRITIONAL APPROACH IN RADICAL CYSTECTOMY WITH URINARY DIVERSION BASED ON ILEAL SEGMENT: A SINGLE-CENTRE, PROSPECTIVE, RANDOMIZED STUDY. <i>Journal of Urology</i> , 2011 , 185,	2.5	1
116	Trans-rectal versus trans-perineal saturation rebiopsy of the prostate: is there a difference in cancer detection rate?. <i>Urology</i> , 2011 , 77, 921-5	1.6	73
115	Robot-Assisted Cystectomy: Strengths and Weaknesses. <i>European Urology Supplements</i> , 2011 , 10, e12-e169		1
114	Metabolic syndrome and benign prostatic hyperplasia: evidence of a potential relationship, hypothesized etiology, and prevention. <i>Korean Journal of Urology</i> , 2011 , 52, 507-16		38
113	Salvage lymphadenectomy in postprostatectomy patients with prostate-specific antigen recurrence. <i>Current Opinion in Urology</i> , 2011 , 21, 237-40	2.8	11
112	How can we predict lymphorrhoea and clinically significant lymphocoeles after radical prostatectomy and pelvic lymphadenectomy? Clinical implications. <i>BJU International</i> , 2011 , 107, 1095-101	5.6	42
111	Sex hormone-binding globulin is a significant predictor of extracapsular extension in men undergoing radical prostatectomy. <i>BJU International</i> , 2011 , 107, 1243-9	5.6	11
110	The role of transrectal saturation biopsy in tumour localization: pathological correlation after retropubic radical prostatectomy and implication for focal ablative therapy. <i>BJU International</i> , 2011 , 108, 366-71	5.6	29
109	Critical assessment of the European Association of Urology guideline indications for pelvic lymph node dissection at radical prostatectomy. <i>BJU International</i> , 2011 , 108, 1769-75	5.6	18
108	When should we expect no residual tumor (pT0) once we submit incidental T1a-b prostate cancers to radical prostatectomy?. <i>International Journal of Urology</i> , 2011 , 18, 148-53	2.3	17
107	Does diabetes mellitus increase the risk of high-grade prostate cancer in patients undergoing radical prostatectomy?. <i>Prostate Cancer and Prostatic Diseases</i> , 2011 , 14, 74-8	6.2	31

106	What is the definition of a satisfactory erectile function after bilateral nerve sparing radical prostatectomy?. <i>Journal of Sexual Medicine</i> , 2011 , 8, 1210-7	1.1	32
105	Combination of adjuvant hormonal and radiation therapy significantly prolongs survival of patients with pT2-4 pN+ prostate cancer: results of a matched analysis. <i>European Urology</i> , 2011 , 59, 832-40	10.2	141
104	Pelvic/retroperitoneal salvage lymph node dissection for patients treated with radical prostatectomy with biochemical recurrence and nodal recurrence detected by [11C]choline positron emission tomography/computed tomography. <i>European Urology</i> , 2011 , 60, 935-43	10.2	163
103	Preoperative hypogonadism is not an independent predictor of high-risk disease in patients undergoing radical prostatectomy. <i>Cancer</i> , 2011 , 117, 3953-62	6.4	44
102	Circulating estradiol, but not testosterone, is a significant predictor of high-grade prostate cancer in patients undergoing radical prostatectomy. <i>Cancer</i> , 2011 , 117, 5029-38	6.4	31
101	The prognostic significance of capsular incision into tumor during radical prostatectomy. <i>International Braz J Urol: Official Journal of the Brazilian Society of Urology</i> , 2011 , 37, 549-550	2	1
100	Penile Rehabilitation After Robotic Radical Prostatectomy: The Best Strategy 2011 , 361-370		
99	Mid-term complications after placement of the male adjustable suburethral sling: a single center experience. <i>International Braz J Urol: Official Journal of the Brazilian Society of Urology</i> , 2011 , 37, 552-553 ²		
98	Testing the most stringent criteria for selection of candidates for active surveillance in patients with low-risk prostate cancer. <i>BJU International</i> , 2010 , 105, 1548-52	5.6	41
97	A nomogram predicting prostate cancer-specific mortality after radical prostatectomy. <i>Urologia Internationalis</i> , 2010 , 84, 132-40	1.9	22
96	The rate of secondary malignancies after radical prostatectomy versus external beam radiation therapy for localized prostate cancer: a population-based study on 17,845 patients. <i>International Journal of Radiation Oncology Biology Physics</i> , 2010 , 76, 342-8	4	72
95	When to perform bone scan in patients with newly diagnosed prostate cancer: external validation of the currently available guidelines and proposal of a novel risk stratification tool. <i>European Urology</i> , 2010 , 57, 551-8	10.2	112
94	Predicting erectile function recovery after bilateral nerve sparing radical prostatectomy: a proposal of a novel preoperative risk stratification. <i>Journal of Sexual Medicine</i> , 2010 , 7, 2521-31	1.1	85
93	Predictive models before and after radical prostatectomy. <i>Prostate</i> , 2010 , 70, 1371-8	4.2	24
92	Surgery and erectile dysfunction. <i>Archivos Espanoles De Urologia</i> , 2010 , 63, 640-8	0.4	3
91	Prediction of delayed graft function after renal transplantation. <i>Canadian Urological Association Journal</i> , 2009 , 3, 377-82	1.2	31
90	Lymphovascular invasion predicts clinical outcomes in patients with node-negative upper tract urothelial carcinoma. <i>Journal of Clinical Oncology</i> , 2009 , 27, 612-8	2.2	209
89	Unsuccessful investigation of preoperative sexual health issues in the prostate cancer "couple": results of a real-life psychometric survey at a major tertiary academic center. <i>Journal of Sexual Medicine</i> , 2009 , 6, 3347-55	1.1	6

88	Assessment of pathological prostate cancer characteristics in men with favorable biopsy features on predominantly sextant biopsy. <i>European Urology</i> , 2009 , 55, 617-28-6	10.2	18
87	A preoperative prognostic model for patients treated with nephrectomy for renal cell carcinoma. <i>European Urology</i> , 2009 , 55, 287-95	10.2	96
86	Two positive nodes represent a significant cut-off value for cancer specific survival in patients with node positive prostate cancer. A new proposal based on a two-institution experience on 703 consecutive N+ patients treated with radical prostatectomy, extended pelvic lymph node dissection and adjuvant therapy. <i>European Urology</i> , 2009 , 55, 261-70	10.2	213
85	Long-term follow-up of patients with prostate cancer and nodal metastases treated by pelvic lymphadenectomy and radical prostatectomy: the positive impact of adjuvant radiotherapy. <i>European Urology</i> , 2009 , 55, 1003-11	10.2	138
84	Are infertile men less healthy than fertile men? Results of a prospective case-control survey. <i>European Urology</i> , 2009 , 56, 1025-31	10.2	94
83	Pre-treatment biomarker levels improve the accuracy of post-prostatectomy nomogram for prediction of biochemical recurrence. <i>Prostate</i> , 2009 , 69, 886-94	4.2	34
82	Effect of body mass index on prostate-specific antigen and percentage free prostate-specific antigen: results from a prostate cancer screening cohort of 1490 men. <i>International Journal of Urology</i> , 2009 , 16, 91-5	2.3	5
81	The probability of Gleason score upgrading between biopsy and radical prostatectomy can be accurately predicted. <i>International Journal of Urology</i> , 2009 , 16, 526-9	2.3	15
80	Obesity does not increase the risk of lymph node metastases in patients with clinically localized prostate cancer undergoing radical prostatectomy and extended pelvic lymph node dissection. <i>International Journal of Urology</i> , 2009 , 16, 676-81	2.3	8
79	Tumour architecture is an independent predictor of outcomes after nephroureterectomy: a multi-institutional analysis of 1363 patients. <i>BJU International</i> , 2009 , 103, 307-11	5.6	131
78	Leydig cell tumour of the testis: presentation, therapy, long-term follow-up and the role of organ-sparing surgery in a single-institution experience. <i>BJU International</i> , 2009 , 103, 197-200	5.6	57
77	Assessing the minimum number of lymph nodes needed at radical cystectomy in patients with bladder cancer. <i>BJU International</i> , 2009 , 103, 1359-62	5.6	58
76	Baseline renal function, ischaemia time and blood loss predict the rate of renal failure after partial nephrectomy. <i>BJU International</i> , 2009 , 103, 1632-5	5.6	36
75	External validation of the updated partin tables in a cohort of French and Italian men. <i>International Journal of Radiation Oncology Biology Physics</i> , 2009 , 73, 347-52	4	42
74	Partin Tables cannot accurately predict the pathological stage at radical prostatectomy. <i>European Journal of Surgical Oncology</i> , 2009 , 35, 123-8	3.6	27
73	Evidence-based sex-related outcomes after radical nephroureterectomy for upper tract urothelial carcinoma: results of large multicenter study. <i>Urology</i> , 2009 , 73, 142-6	1.6	62
72	Partial versus radical nephrectomy in patients with adverse clinical or pathologic characteristics. <i>Urology</i> , 2009 , 73, 1300-5	1.6	55
71	Population-based analysis of normal Total PSA and percentage of free/Total PSA values: results from screening cohort. <i>Urology</i> , 2009 , 73, 1323-7	1.6	10

70	Biopsy core number represents one of foremost predictors of clinically significant gleason sum upgrading in patients with low-risk prostate cancer. <i>Urology</i> , 2009 , 73, 1087-91	1.6	93
69	Assessment of the minimum number of lymph nodes needed to detect lymph node invasion at radical nephroureterectomy in patients with upper tract urothelial cancer. <i>Urology</i> , 2009 , 74, 1070-4	1.6	48
68	Prevention and Management of Postprostatectomy Erectile Dysfunction. <i>European Urology Supplements</i> , 2009 , 8, 80-87	0.9	3
67	Benign Prostatic Hyperplasia and Its Aetiologies. <i>European Urology Supplements</i> , 2009 , 8, 865-871	0.9	75
66	Is sperm banking of interest to patients with nongerm cell urological cancer before potentially fertility damaging treatments?. <i>Journal of Urology</i> , 2009 , 182, 1101-7	2.5	14
65	MORTALITY PREDICTIONS IN PATIENTS WITH ADRENOCORTICAL CARCINOMA. <i>Journal of Urology</i> , 2009 , 181, 9-10	2.5	3
64	PATIENTS WITH ORGAN CONFINED PROSTATE CANCER AND POSTITIVE SURGICAL MARGINS HAVE SIMILAR RECURRENCE RATES COMPARED TO PATIENTS WITH EXTRA-CAPSULAR EXTENSION AND NEGATIVE SURGICAL MARGINS. A PLEA FOR STAGE RE-CLASSIFICATION. <i>Journal of Urology</i> , 2009 , 181, 290-290	2.5	44
63	GENDER IS AN IMPORTANT PREDICTOR OF CANCER-SPECIFIC SURVIVAL IN PATIENT WITH TRANSITIONAL CELL CARCINOMA AFTER RADICAL CYSTECTOMY. <i>Journal of Urology</i> , 2009 , 181, 635	2.5	14
62	Impact of lymph node dissection on cancer specific survival in patients with upper tract urothelial carcinoma treated with radical nephroureterectomy. <i>Journal of Urology</i> , 2009 , 181, 2482-9	2.5	154
61	VALIDATION OF THE CRITERIA SUGGESTED BY CURRENT GUIDELINES TO INDICATE THE NEED FOR BASELINE STAGING BONE SCAN IN PATIENTS WITH NEWLY DIAGNOSED PROSTATE CANCER. <i>Journal of Urology</i> , 2009 , 181, 782-782	2.5	3
60	Conditional survival predictions after nephrectomy for renal cell carcinoma. <i>Journal of Urology</i> , 2009 , 182, 2607-12	2.5	45
59	Holmium laser enucleation of the prostate and holmium laser ablation of the prostate: indications and outcome. <i>Current Opinion in Urology</i> , 2009 , 19, 38-43	2.8	37
58	Prophylaxis of erectile function after radical prostatectomy with phosphodiesterase type 5 inhibitors. <i>Current Pharmaceutical Design</i> , 2009 , 15, 3496-501	3.3	4
57	Survival after radical prostatectomy and radiotherapy for prostate cancer: a population-based study. <i>Canadian Urological Association Journal</i> , 2009 , 3, 13-21	1.2	14
56	Age at diagnosis is a determinant factor of renal cell carcinoma-specific survival in patients treated with nephrectomy. <i>Canadian Urological Association Journal</i> , 2008 , 2, 610-7	1.2	29
55	A nomogram predicting metastatic progression after radical prostatectomy. <i>International Journal of Urology</i> , 2008 , 15, 889-94	2.3	10
54	A nomogram is more accurate than a regression tree in predicting lymph node invasion in prostate cancer. <i>BJU International</i> , 2008 , 101, 556-60	5.6	21
53	The effect of surgical volume, age and comorbidities on 30-day mortality after radical prostatectomy: a population-based analysis of 9208 consecutive cases. <i>BJU International</i> , 2008 , 101, 826-32	5.6	24

52	Comparison of stage migration patterns between Europe and the USA: an analysis of 11 350 men treated with radical prostatectomy for prostate cancer. <i>BJU International</i> , 2008 , 101, 1513-8	5.6	122
51	Adjuvant chemotherapy for bladder cancer does not alter cancer-specific survival after cystectomy in a matched case-control study. <i>BJU International</i> , 2008 , 101, 1356-61	5.6	14
50	Accuracy of life tables in predicting overall survival in patients after radical prostatectomy. <i>BJU International</i> , 2008 , 102, 33-8	5.6	24
49	High surgical volume is associated with a lower rate of secondary therapy after radical prostatectomy for localized prostate cancer. <i>BJU International</i> , 2008 , 102, 463-7	5.6	17
48	Nerve-sparing radical retropubic prostatectomy in patients previously submitted to holmium laser enucleation of the prostate for bladder outlet obstruction due to benign prostatic enlargement. <i>European Urology</i> , 2008 , 53, 1180-5	10.2	30
47	Acceptance of and discontinuation rate from erectile dysfunction oral treatment in patients following bilateral nerve-sparing radical prostatectomy. <i>European Urology</i> , 2008 , 53, 564-70	10.2	70
46	Poor overall survival in septa- and octogenarian patients after radical prostatectomy and radiotherapy for prostate cancer: a population-based study of 6183 men. <i>European Urology</i> , 2008 , 54, 107-16	10.2	26
45	High provider volume is associated with lower rate of secondary therapies after definitive radiotherapy for localized prostate cancer. <i>European Urology</i> , 2008 , 54, 97-105	10.2	14
44	Neoadjuvant sunitinib induction therapy may effectively down-stage renal cell carcinoma atrial thrombi. <i>European Urology</i> , 2008 , 53, 845-8	10.2	93
43	Toxicities associated with the administration of sorafenib, sunitinib, and temsirolimus and their management in patients with metastatic renal cell carcinoma. <i>European Urology</i> , 2008 , 53, 917-30	10.2	175
42	Validation of the contemporary epstein criteria for insignificant prostate cancer in European men. <i>European Urology</i> , 2008 , 54, 1306-13	10.2	95
41	Reply to Michael Staehler@ Letter to the Editor re: Pierre I Karakiewicz, Nazareno Suardi, Claudio Jeldres, et al. Neoadjuvant Sunitinib Induction Therapy May Effectively Down-Stage Renal Cell Carcinoma Atrial Thrombi. <i>Eur Urol</i> 2008;53:845-848. <i>European Urology</i> , 2008 , 54, 951-953	10.2	
40	Impact of surgical volume on the rate of lymph node metastases in patients undergoing radical prostatectomy and extended pelvic lymph node dissection for clinically localized prostate cancer. <i>European Urology</i> , 2008 , 54, 794-802	10.2	52
39	Pregnant woman presenting with a gross retroperitoneal mass: surgical treatment with caval replacement. <i>European Urology</i> , 2008 , 54, 677-80	10.2	2
38	Baseline prevalence of erectile dysfunction in a prostate cancer screening population. <i>Journal of Sexual Medicine</i> , 2008 , 5, 428-35	1.1	20
37	The effect of comorbidities and socioeconomic status on sexual and urinary function in men undergoing prostate cancer screening. <i>Journal of Sexual Medicine</i> , 2008 , 5, 668-76	1.1	12
36	Body mass index and its association with genitourinary disorders in men undergoing prostate cancer screening. <i>Journal of Sexual Medicine</i> , 2008 , 5, 2141-51	1.1	9
35	COMBINATION OF CELL CYCLE REGULATING BIO-MARKERS IMPROVES PROGNOSIS IN PATIENTS WITH ORGAN CONFINED UROTHELIAL CANCER AT RADICAL CYSTECTOMY. <i>Journal of Urology</i> , 2008 , 179, 578-578	2.5	7

34	External validation of the updated Partin tables in a cohort of North American men. <i>Journal of Urology</i> , 2008 , 180, 898-902; discussion 902-3	2.5	34
33	Holmium laser enucleation versus transurethral resection of the prostate: results from a 2-center prospective randomized trial in patients with obstructive benign prostatic hyperplasia. <i>Journal of Urology</i> , 2008 , 179, S87-90	2.5	79
32	EXTERNAL-BEAM RADIATION THERAPY INCREASES THE RATE OF SECONDARY MALIGNANCIES RELATIVE TO RADICAL PROSTATECTOMY IN MEN WITH PROSTATE CANCER. <i>Journal of Urology</i> , 2008 , 179, 113-113	2.5	2
31	NEPHRON-SPARING SURGERY VS. RADICAL NEPHRECTOMY IN PATIENTS WITH RENAL CELL CARCINOMA >7 CM. WITH NO EVIDENCE OF NODAL OR DISTANT METASTASIS. <i>Journal of Urology</i> , 2008 , 179, 417-418	2.5	2
30	ADJUVANT CHEMOTHERAPY FOR UPPER TRACT TRANSITIONAL CELL CARCINOMA: RESULTS FROM THE UPPER TRACT UCC CONSORTIUM. <i>Journal of Urology</i> , 2008 , 179, 119-119	2.5	2
29	CURRENTLY USED CRITERIA FOR ACTIVE SURVEILLANCE IN MEN WITH LOW RISK PROSTATE CANCER. AN ANALYSIS OF PATHOLOGICAL FEATURES. <i>Journal of Urology</i> , 2008 , 179, 152-152	2.5	71
28	THE EFFECT OF ANDROGEN DEPRIVATION THERAPY ON THE RATE OF SUBSEQUENT NON-CANCER MORBIDITIES. <i>Journal of Urology</i> , 2008 , 179, 186-186	2.5	35
27	Comparison of nomograms with other methods for predicting outcomes in prostate cancer: a critical analysis of the literature. <i>Clinical Cancer Research</i> , 2008 , 14, 4400-7	12.9	197
26	External validation of a biomarker-based preoperative nomogram predicts biochemical recurrence after radical prostatectomy. <i>Journal of Clinical Oncology</i> , 2008 , 26, 1526-31	2.2	50
25	Lower urinary tract symptoms affect one-third of men in a prostate cancer screening population. <i>Journal of Endourology</i> , 2008 , 22, 369-76	2.7	7
24	Biopsies performed at tertiary care centers are superior to referral biopsies in predicting pathologic Gleason sum. <i>Journal of Endourology</i> , 2008 , 22, 533-8	2.7	6
23	A nomogram predicting long-term biochemical recurrence after radical prostatectomy. <i>Cancer</i> , 2008 , 112, 1254-63	6.4	95
22	Critical assessment of tools to predict clinically insignificant prostate cancer at radical prostatectomy in contemporary men. <i>Cancer</i> , 2008 , 113, 701-9	6.4	74
21	Currently used criteria for active surveillance in men with low-risk prostate cancer: an analysis of pathologic features. <i>Cancer</i> , 2008 , 113, 2068-72	6.4	88
20	Mortality at 120 days after prostatic biopsy: a population-based study of 22,175 men. <i>International Journal of Cancer</i> , 2008 , 123, 647-52	7.5	32
19	Obesity does not predispose to more aggressive prostate cancer either at biopsy or radical prostatectomy in European men. <i>International Journal of Cancer</i> , 2007 , 121, 791-5	7.5	42
18	Impact of chronic prostatitis-like symptoms on the quality of life in a large group of men. <i>BJU International</i> , 2007 , 100, 1307-11	5.6	57
17	A nomogram for staging of exclusive nonobturator lymph node metastases in men with localized prostate cancer. <i>European Urology</i> , 2007 , 51, 112-9; discussion 119-20	10.2	54

16	Detection of lymph-node metastases with integrated [11C]choline PET/CT in patients with PSA failure after radical retropubic prostatectomy: results confirmed by open pelvic-retroperitoneal lymphadenectomy. <i>European Urology</i> , 2007 , 52, 423-9	10.2	199
15	Preserved postoperative penile size correlates well with maintained erectile function after bilateral nerve-sparing radical retropubic prostatectomy. <i>European Urology</i> , 2007 , 52, 702-7	10.2	56
14	Body mass index does not predict prostate-specific antigen or percent free prostate-specific antigen in men undergoing prostate cancer screening. <i>European Journal of Cancer</i> , 2007 , 43, 1180-7	7.5	24
13	Prostate volume and adverse prostate cancer features: fact not artifact. <i>European Journal of Cancer</i> , 2007 , 43, 2669-77	7.5	66
12	Holmium laser enucleation of the prostate versus open prostatectomy for prostates >70 g: 24-month follow-up. <i>European Urology</i> , 2006 , 50, 563-8	10.2	264
11	Complications and other surgical outcomes associated with extended pelvic lymphadenectomy in men with localized prostate cancer. <i>European Urology</i> , 2006 , 50, 1006-13	10.2	266
10	Diagnosis and Treatment of the Circumcaval Ureter. <i>European Urology Supplements</i> , 2006 , 5, 449-462	0.9	20
9	Radical prostatectomy after previous prostate surgery: clinical and functional outcomes. <i>Journal of Urology</i> , 2006 , 176, 2459-63; discussion 2463	2.5	86
8	Holmium laser enucleation versus open prostatectomy for benign prostatic hyperplasia: an inpatient cost analysis. <i>Urology</i> , 2006 , 68, 302-6	1.6	85
7	General versus spinal anesthesia with different forms of sedation in patients undergoing radical retropubic prostatectomy: results of a prospective, randomized study. <i>International Journal of Urology</i> , 2006 , 13, 1185-90	2.3	16
6	Safety and tolerability of oral erectile dysfunction treatments in the elderly. <i>Drugs and Aging</i> , 2005 , 22, 323-38	4.7	5
5	Pfannenstiel versus vertical laparotomy in patients undergoing radical retropubic prostatectomy with spinal anesthesia: results of a prospective, randomized trial. <i>European Urology</i> , 2005 , 47, 202-8	10.2	13
4	Vardenafil for the treatment of erectile dysfunction: a critical review of the literature based on personal clinical experience. <i>European Urology</i> , 2005 , 47, 612-21	10.2	31
3	Improving the preservation of the urethral sphincter and neurovascular bundles during open radical retropubic prostatectomy. <i>European Urology</i> , 2005 , 48, 938-45	10.2	118
2	Holmium laser enucleation versus transurethral resection of the prostate: results from a 2-center, prospective, randomized trial in patients with obstructive benign prostatic hyperplasia. <i>Journal of Urology</i> , 2004 , 172, 1926-9	2.5	239
1	General versus spinal anesthesia in patients undergoing radical retropubic prostatectomy: results of a prospective, randomized study. <i>Urology</i> , 2004 , 64, 95-100	1.6	40