Grant N Wheeler

List of Publications by Year
in descending order

60 papers	2,791 citations	257101 24 h-index	182168 51 g-index
68 all docs	68 docs citations	68 times ranked	4294 citing authors

#	Article	IF	Citations
1	An efficient miRNA knockout approach using CRISPR-Cas9 in Xenopus. Developmental Biology, 2022, 483, 66-75.	0.9	8
2	MicroRNAs in neural crest development and neurocristopathies. Biochemical Society Transactions, 2022, 50, 965-974.	1.6	5
3	Characterising open chromatin in chick embryos identifies cis-regulatory elements important for paraxial mesoderm formation and axis extension. Nature Communications, 2021, 12, 1157.	5.8	8
4	Expression analysis of chick Frizzled receptors during spinal cord development. Gene Expression Patterns, 2021, 39, 119167.	0.3	1
5	In Vivo Assessment of Drug-Induced Hepatotoxicity Using <i>Xenopus</i> Embryos. Cold Spring Harbor Protocols, 2020, 2020, pdb.prot106096.	0.2	4
6	FZD10 regulates cell proliferation and mediates Wnt1 induced neurogenesis in the developing spinal cord. PLoS ONE, 2020, 15, e0219721.	1.1	11
7	Toxicity and biodegradation of zinc ferrite nanoparticles in Xenopus laevis. Journal of Nanoparticle Research, 2019, 21, 1.	0.8	6
8	Microsyntenic Clusters Reveal Conservation of lncRNAs in Chordates Despite Absence of Sequence Conservation. Biology, 2019, 8, 61.	1.3	19
9	Solid lipid nanoparticles for the delivery of anti-microbial oligonucleotides. European Journal of Pharmaceutics and Biopharmaceutics, 2019, 134, 166-177.	2.0	42
10	Paracetamol-induced liver injury modelled in Xenopus laevis embryos. Toxicology Letters, 2019, 302, 83-91.	0.4	13
11	Unravelling the mechanisms that determine the uptake and metabolism of magnetic single and multicore nanoparticles in a<i>Xenopus laevis</i>model. Nanoscale, 2018, 10, 690-704.	2.8	21
12	microRNAs associated with early neural crest development in Xenopus laevis. BMC Genomics, 2018, 19, 59.	1.2	22
13	miR-133 mediated regulation of the hedgehog pathway orchestrates embryo myogenesis. Development (Cambridge), 2018, 145, .	1.2	28
14	ADAMTS9, a member of the ADAMTS family, in Xenopus development. Gene Expression Patterns, 2018, 29, 72-81.	0.3	12
15	The anti-rheumatic drug, leflunomide, synergizes with MEK inhibition to suppress melanoma growth. Oncotarget, 2018, 9, 3815-3829.	0.8	17
16	Cationic liposomal vectors incorporating a bolaamphiphile for oligonucleotide antimicrobials. Biochimica Et Biophysica Acta - Biomembranes, 2017, 1859, 1767-1777.	1.4	22
17	Combining Cytotoxicity Assessment and Xenopus laevis Phenotypic Abnormality Assay as a Predictor of Nanomaterial Safety. Current Protocols in Toxicology / Editorial Board, Mahin D Maines (editor-in-chief) [et Al], 2017, 73, 20.13.1-20.13.33.	1.1	3
18	Frizzled-7 is required for Xenopus heart development. Biology Open, 2017, 6, 1861-1868.	0.6	8

#	Article	IF	Citations
19	A functional approach to understanding the role of NCKX5 in Xenopus pigmentation. PLoS ONE, 2017, 12, e0180465.	1.1	11
20	The positive transcriptional elongation factor (P-TEFb) is required for neural crest specification. Developmental Biology, 2016, 416, 361-372.	0.9	12
21	An early developmental vertebrate model for nanomaterial safety: bridging cell-based and mammalian toxicity assessment. Nanomedicine, 2016, 11, 643-656.	1.7	21
22	Sprouty2 mediated tuning of signalling is essential for somite myogenesis. BMC Medical Genomics, 2015, 8, S8.	0.7	2
23	A Database of microRNA Expression Patterns in Xenopus laevis. PLoS ONE, 2015, 10, e0138313.	1.1	21
24	The ADAMTS (A Disintegrin and Metalloproteinase with Thrombospondin motifs) family. Genome Biology, 2015, 16, 113.	3.8	471
25	It's about time for neural crest. Science, 2015, 348, 1316-1317.	6.0	7
26	Klhl31 attenuates Î²-catenin dependent Wnt signaling and regulates embryo myogenesis. Developmental Biology, 2015, 402, 61-71.	0.9	17
27	myomiR-dependent switching of BAF60 variant incorporation into Brg1 chromatin remodeling complexes during embryo myogenesis. Development (Cambridge), 2014, 141, 3378-3387.	1.2	58
28	Comparative mode-of-action analysis following manual and automated phenotype detection in<i>Xenopus laevis</i>. MedChemComm, 2014, 5, 386-396.	3.5	2
29	Smad1 transcription factor integrates BMP2 and Wnt3a signals in migrating cardiac progenitor cells. Proceedings of the National Academy of Sciences of the United States of America, 2014, 111, 7337-7342.	3.3	40
30	Frizzled-7 expression during early cardiogenesis of Xenopus laevisembryo. BMC Genomics, 2014, 15, .	1.2	1
31	Extensions to In Silico Bioactivity Predictions Using Pathway Annotations and Differential Pharmacology Analysis: Application to <i>Xenopus laevis</i> Phenotypic Readouts. Molecular Informatics, 2013, 32, 1009-1024.	1.4	13
32	Phenotypic Screens with Model Organisms. , 2012, , 121-136.		2
33	Chemical Genetics and Drug Discovery in Xenopus. Methods in Molecular Biology, 2012, 917, 155-166.	0.4	24
34	<i>Xenopus</i>: An ideal system for chemical genetics. Genesis, 2012, 50, 207-218.	0.8	27
35	DHODH modulates transcriptional elongation in the neural crest and melanoma. Nature, 2011, 471, 518-522.	13.7	411
36	Simple vertebrate models for chemical genetics and drug discovery screens: Lessons from zebrafish and <i>Xenopus</i>. Developmental Dynamics, 2009, 238, 1287-1308.	0.8	156

#	Article	IF	Citations
37	A Chemical Genomic Approach Identifies Matrix Metalloproteinases as Playing an Essential and Specific Role in Xenopus Melanophore Migration. Chemistry and Biology, 2009, 16, 93-104.	6.2	44
38	Frizzled-10 promotes sensory neuron development in Xenopus embryos. Developmental Biology, 2009, 335, 143-155.	0.9	27
39	13-P092 Klhl31 is regulated by myogenic signals in developing somites and modulates Wnt signaling in vitro and in vivo. Mechanisms of Development, 2009, 126, S222.	1.7	0
40	Chemical genomics identifies compounds affecting Xenopus laevis pigment cell development. Molecular BioSystems, 2009, 5, 376.	2.9	35
41	Wnt6 expression in epidermis and epithelial tissues during <i>Xenopus</i> organogenesis. Developmental Dynamics, 2008, 237, 768-779.	0.8	24
42	Three matrix metalloproteinases are required in vivo for macrophage migration during embryonic development. Mechanisms of Development, 2008, 125, 1059-1070.	1.7	48
43	Inducible Gene Expression in Transient Transgenic Xenopus Embryos. Methods in Molecular Biology, 2008, 469, 431-449.	0.4	3
44	The MH1 domain of Smad3 interacts with Pax6 and represses autoregulation of the Pax6 P1 promoter. Nucleic Acids Research, 2007, 35, 890-901.	6.5	44
45	Frizzled7 mediates canonical Wnt signaling in neural crest induction. Developmental Biology, 2006, 298, 285-298.	0.9	66
46	FGF-4 signaling is involved in mir-206 expression in developing somites of chicken embryos. Developmental Dynamics, 2006, 235, 2185-2191.	0.8	82
47	Xenopus as a model organism in developmental chemical genetic screens. Molecular BioSystems, 2005, 1, 223.	2.9	46
48	Matrix metalloproteinase genes inXenopus development. Developmental Dynamics, 2004, 231, 214-220.	0.8	58
49	The cloning, genomic organization and expression of the focal contact protein paxillin in Drosophila. Gene, 2001, 262, 291-299.	1.0	16
50	Difference in XTcf-3 dependency accounts for change in response to Î²-catenin-mediated Wnt signalling in Xenopus blastula. Development (Cambridge), 2001, 128, 2063-2073.	1.2	50
51	Inducible gene expression in transgenic Xenopus embryos. Current Biology, 2000, 10, 849-852.	1.8	66
52	Binding of Integrin Î±6Î²4 to Plectin Prevents Plectin Association with F-Actin but Does Not Interfere with Intermediate Filament Binding. Journal of Cell Biology, 1999, 147, 417-434.	2.3	171
53	Two novel Xenopus frizzled genes expressed in developing heart and brain. Mechanisms of Development, 1999, 86, 203-207.	1.7	51
54	Mouse Desmocollin (Dsc3) and Desmoglein (Dsg1) Genes Are Closely Linked in the Proximal Region of Chromosome 18. Genomics, 1994, 21, 510-516.	1.3	41

#	Article	IF	Citations
55	Desmosomal glycoproteins I, II and III: novel members of the cadherin superfamily. Biochemical Society Transactions, 1991, 19, 1060-1064.	1.6	23
56	Chromosomal assignment of the human genes coding for the major proteins of the desmosome junction, desmoglein DGI (DSG), desmocollins DGIIIII (DSC), desmoplakins DPIII (DSP), and plakoglobin DPIII (JUP). Genomics, 1991, 10, 640-645.	1.3	69
57	Desmosomal glycoprotein DGI, a component of intercellular desmosome junctions, is related to the cadherin family of cell adhesion molecules.. Proceedings of the National Academy of Sciences of the United States of America, 1991, 88, 4796-4800.	3.3	168
58	Desmosomal glycoproteins II and III. Cadherin-like junctional molecules generated by alternative splicing. Journal of Biological Chemistry, 1991, 266, 10438-45.	1.6	90
59	Identification of the B1 and B2 subunits of human placental laminin and rat parietal-yolk-sac laminin using antisera specific for murine laminin-<i>Î²</i>-galactosidase fusion proteins. Biochemical Journal, 1990, 270, 463-468.	1.7	17
60	The use of bacterial fusion proteins in the production of anti-laminin antibodies. Biochemical Society Transactions, 1989, 17, 185-186.	1.6	2