

Jim Holland

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4474684/jim-holland-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

133
papers

12,084
citations

42
h-index

109
g-index

143
ext. papers

14,904
ext. citations

6.8
avg. IF

6.1
L-index

#	Paper	IF	Citations
133	Outlook for Implementation of Genomics-Based Selection in Public Cotton Breeding Programs. <i>Plants</i> , 2022 , 11, 1446	4.5	1
132	Eleven biosynthetic genes explain the majority of natural variation in carotenoid levels in maize grain. <i>Plant Cell</i> , 2021 , 33, 882-900	11.6	3
131	Characterizing the oligogenic architecture of plant growth phenotypes informs genomic selection approaches in a common wheat population. <i>BMC Genomics</i> , 2021 , 22, 402	4.5	3
130	Effects of artificial inoculation on trait correlations with resistance to Fusarium ear rot and fumonisin contamination in maize. <i>Crop Science</i> , 2021 , 61, 2522	2.4	0
129	The importance of dominance and genotype-by-environment interactions on grain yield variation in a large-scale public cooperative maize experiment. <i>G3: Genes, Genomes, Genetics</i> , 2021 , 11,	3.2	12
128	Domestication reshaped the genetic basis of inbreeding depression in a maize landrace compared to its wild relative, teosinte.. <i>PLoS Genetics</i> , 2021 , 17, e1009797	6	1
127	Environment-specific genomic prediction ability in maize using environmental covariates depends on environmental similarity to training data.. <i>G3: Genes, Genomes, Genetics</i> , 2021 ,	3.2	2
126	The genetic architecture of the maize progenitor, teosinte, and how it was altered during maize domestication. <i>PLoS Genetics</i> , 2020 , 16, e1008791	6	17
125	Heterosis of leaf and rhizosphere microbiomes in field-grown maize. <i>New Phytologist</i> , 2020 , 228, 1055-1069	10.69	25
124	Genetic variation for response to mixed triazole and strobilurin application in diverse maize 2020 , 3, e20054		
123	Maize genomes to fields (G2F): 2014-2017 field seasons: genotype, phenotype, climatic, soil, and inbred ear image datasets. <i>BMC Research Notes</i> , 2020 , 13, 71	2.3	16
122	Genomic prediction for resistance to Fusarium ear rot and fumonisin contamination in maize. <i>Crop Science</i> , 2020 , 60, 1863-1875	2.4	10
121	Dominance Effects and Functional Enrichments Improve Prediction of Agronomic Traits in Hybrid Maize. <i>Genetics</i> , 2020 , 215, 215-230	4	12
120	Enhancing Crop Breeding Using Population Genomics Approaches. <i>Population Genomics</i> , 2020 , 1	1.4	
119	Relative utility of agronomic, phenological, and morphological traits for assessing genotype-by-environment interaction in maize inbreds. <i>Crop Science</i> , 2020 , 60, 62-81	2.4	7
118	Association mapping and genomic prediction for ear rot disease caused by <i>Fusarium verticillioides</i> in a tropical maize germplasm. <i>Crop Science</i> , 2020 , 60, 2867-2881	2.4	3
117	Genome-wide association analysis of the strength of the MAMP-elicited defense response and resistance to target leaf spot in sorghum. <i>Scientific Reports</i> , 2020 , 10, 20817	4.9	2

116	Utility of Climatic Information via Combining Ability Models to Improve Genomic Prediction for Yield Within the Genomes to Fields Maize Project. <i>Frontiers in Genetics</i> , 2020 , 11, 592769	4.5	11
115	A Genome-Wide Association Study To Understand the Effect of Infection on Seedlings of a Maize Diversity Panel. <i>G3: Genes, Genomes, Genetics</i> , 2020 , 10, 1685-1696	3.2	2
114	Training population selection and use of fixed effects to optimize genomic predictions in a historical USA winter wheat panel. <i>Theoretical and Applied Genetics</i> , 2019 , 132, 1247-1261	6	41
113	Optimal Designs for Genomic Selection in Hybrid Crops. <i>Molecular Plant</i> , 2019 , 12, 390-401	14.4	41
112	The genetic architecture of teosinte catalyzed and constrained maize domestication. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019 , 116, 5643-5652	11.5	34
111	Diverse Components of Resistance to Infection and Fumonisin Contamination in Four Maize Recombinant Inbred Families. <i>Toxins</i> , 2019 , 11,	4.9	11
110	Genomic-Wide Association Study of Popping Expansion in Tropical Popcorn and Field Corn Germplasm. <i>Crop Science</i> , 2019 , 59, 2007-2019	2.4	14
109	The Genomic Basis for Short-Term Evolution of Environmental Adaptation in Maize. <i>Genetics</i> , 2019 , 213, 1479-1494	4	11
108	Validation and Characterization of Maize Multiple Disease Resistance QTL. <i>G3: Genes, Genomes, Genetics</i> , 2019 , 9, 2905-2912	3.2	7
107	A Genome Wide Association Study Reveals Markers and Genes Associated with Resistance to Infection of Seedlings in a Maize Diversity Panel. <i>G3: Genes, Genomes, Genetics</i> , 2019 , 9, 571-579	3.2	16
106	Accounting for Genotype-by-Environment Interactions and Residual Genetic Variation in Genomic Selection for Water-Soluble Carbohydrate Concentration in Wheat. <i>G3: Genes, Genomes, Genetics</i> , 2018 , 8, 1909-1919	3.2	7
105	Defining the Role of the MADS-Box Gene, <i>Zea Agamous-like1</i> , a Target of Selection During Maize Domestication. <i>Journal of Heredity</i> , 2018 , 109, 333-338	2.4	15
104	Dissecting Symptomatology and Fumonisin Contamination Produced by <i>Fusarium verticillioides</i> in Maize Ears. <i>Phytopathology</i> , 2018 , 108, 1475-1485	3.8	9
103	Maize Genomes to Fields: 2014 and 2015 field season genotype, phenotype, environment, and inbred ear image datasets. <i>BMC Research Notes</i> , 2018 , 11, 452	2.3	16
102	Ecogeography of teosinte. <i>PLoS ONE</i> , 2018 , 13, e0192676	3.7	36
101	Harnessing Maize Biodiversity. <i>Compendium of Plant Genomes</i> , 2018 , 335-366	0.8	5
100	Plant Genetics: Two Steps on the Path to Maize Adaptation. <i>Current Biology</i> , 2018 , 28, R1098-R1101	6.3	1
99	Enhancing genomic prediction with genome-wide association studies in multiparental maize populations. <i>Heredity</i> , 2017 , 118, 585-593	3.6	43

98	Modifications to a LATE MERISTEM IDENTITY1 gene are responsible for the major leaf shapes of Upland cotton (<i>Gossypium hirsutum</i> L.). <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017 , 114, E57-E66	11.5	48
97	Selection for water-soluble carbohydrate accumulation and investigation of genetic environment interactions in an elite wheat breeding population. <i>Theoretical and Applied Genetics</i> , 2017 , 130, 2445-2461	6	26
96	High-Throughput Resequencing of Maize Landraces at Genomic Regions Associated with Flowering Time. <i>PLoS ONE</i> , 2017 , 12, e0168910	3.7	8
95	Genomic Relationships and GBLUP 2017 , 311-354		2
94	Genomic Selection 2017 , 355-384		
93	Imputing Missing Genotypes 2017 , 287-309		
92	Multivariate Models 2017 , 165-201		
91	Genomic estimation of complex traits reveals ancient maize adaptation to temperate North America. <i>Science</i> , 2017 , 357, 512-515	33.3	110
90	The effect of artificial selection on phenotypic plasticity in maize. <i>Nature Communications</i> , 2017 , 8, 13481	17.4	58
89	Genome-Wide Associations for Water-Soluble Carbohydrate Concentration and Relative Maturity in Wheat Using SNP and DArT Marker Arrays. <i>G3: Genes, Genomes, Genetics</i> , 2017 , 7, 2821-2830	3.2	13
88	Genetic Data Analysis for Plant and Animal Breeding 2017 ,		85
87	Genetic Architecture of Domestication-Related Traits in Maize. <i>Genetics</i> , 2016 , 204, 99-113	4	31
86	Responses to Recurrent Index Selection for Reduced Fusarium Ear Rot and Lodging and for Increased Yield in Maize. <i>Crop Science</i> , 2016 , 56, 85-94	2.4	11
85	The Genetics of Leaf Flecking in Maize and Its Relationship to Plant Defense and Disease Resistance. <i>Plant Physiology</i> , 2016 , 172, 1787-1803	6.6	14
84	Genome-wide association study reveals a set of genes associated with resistance to the Mediterranean corn borer (<i>Sesamia nonagrioides</i> L.) in a maize diversity panel. <i>BMC Plant Biology</i> , 2015 , 15, 35	5.3	55
83	Ensemble Learning of QTL Models Improves Prediction of Complex Traits. <i>G3: Genes, Genomes, Genetics</i> , 2015 , 5, 2073-84	3.2	6
82	Hallauer's Tussock: a decade of selection for tropical-to-temperate phenological adaptation in maize. <i>Heredity</i> , 2015 , 114, 229-40	3.6	19
81	Charles W. Stuber: Maize Geneticist and Pioneer of Marker-Assisted Selection 2015 , 1-22		

80	MAGIC maize: a new resource for plant genetics. <i>Genome Biology</i> , 2015 , 16, 163	18.3	24
79	Charles W. Stuber: Maize Geneticist and Pioneer of Marker-Assisted Selection 2015 , 1-22		
78	Joint-multiple family linkage analysis predicts within-family variation better than single-family analysis of the maize nested association mapping population. <i>Heredity</i> , 2015 , 114, 552-63	3.6	55
77	New insight into a complex plant-fungal pathogen interaction. <i>Nature Genetics</i> , 2015 , 47, 101-3	36.3	6
76	QTL mapping using high-throughput sequencing. <i>Methods in Molecular Biology</i> , 2015 , 1284, 257-85	1.4	18
75	Insights into the effects of long-term artificial selection on seed size in maize. <i>Genetics</i> , 2014 , 198, 409-21		25
74	Yield Effects of Two Southern Leaf Blight Resistance Loci in Maize Hybrids. <i>Crop Science</i> , 2014 , 54, 882-894		3
73	Protein Profiling Reveals Novel Proteins in Pollen and Pistil of W22 (ga1; Ga1) in Maize. <i>Proteomes</i> , 2014 , 2, 258-271	4.6	3
72	A genome-wide association study of the maize hypersensitive defense response identifies genes that cluster in related pathways. <i>PLoS Genetics</i> , 2014 , 10, e1004562	6	43
71	Mining Natural Variation for Maize Improvement: Selection on Phenotypes and Genes 2014 , 615-649		17
70	Limits on the reproducibility of marker associations with southern leaf blight resistance in the maize nested association mapping population. <i>BMC Genomics</i> , 2014 , 15, 1068	4.5	25
69	Genome-wide association study of Fusarium ear rot disease in the U.S.A. maize inbred line collection. <i>BMC Plant Biology</i> , 2014 , 14, 372	5.3	62
68	The genetic architecture of maize height. <i>Genetics</i> , 2014 , 196, 1337-56	4	211
67	Comprehensive genotyping of the USA national maize inbred seed bank. <i>Genome Biology</i> , 2013 , 14, R55	18.3	337
66	Confirming quantitative trait loci for aflatoxin resistance from Mp313E in different genetic backgrounds. <i>Molecular Breeding</i> , 2013 , 32, 15-26	3.4	38
65	A genome-wide association study reveals genes associated with fusarium ear rot resistance in a maize core diversity panel. <i>G3: Genes, Genomes, Genetics</i> , 2013 , 3, 2095-104	3.2	64
64	A connected set of genes associated with programmed cell death implicated in controlling the hypersensitive response in maize. <i>Genetics</i> , 2013 , 193, 609-20	4	40
63	Genetic architecture of maize kernel composition in the nested association mapping and inbred association panels. <i>Plant Physiology</i> , 2012 , 158, 824-34	6.6	242

62	Multivariate mixed linear model analysis of longitudinal data: an information-rich statistical technique for analyzing plant disease resistance. <i>Phytopathology</i> , 2012 , 102, 1016-25	3.8	4
61	Maize HapMap2 identifies extant variation from a genome in flux. <i>Nature Genetics</i> , 2012 , 44, 803-7	36.3	470
60	Diallel Analysis of Resistance to Fusarium Ear Rot and Fumonisin Contamination in Maize. <i>Crop Science</i> , 2012 , 52, 2173-2181	2.4	46
59	ZmCCT and the genetic basis of day-length adaptation underlying the postdomestication spread of maize. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012 , 109, E1913-215	11.5	226
58	The relationship between parental genetic or phenotypic divergence and progeny variation in the maize nested association mapping population. <i>Heredity</i> , 2012 , 108, 490-9	3.6	96
57	Genome-wide association study of leaf architecture in the maize nested association mapping population. <i>Nature Genetics</i> , 2011 , 43, 159-62	36.3	746
56	Genome-wide association study of quantitative resistance to southern leaf blight in the maize nested association mapping population. <i>Nature Genetics</i> , 2011 , 43, 163-8	36.3	444
55	A novel genetic framework for studying response to artificial selection. <i>Plant Genetic Resources: Characterisation and Utilisation</i> , 2011 , 9, 281-283	1	8
54	Three new teosintes (<i>Zea</i> spp., Poaceae) from Mexico. <i>American Journal of Botany</i> , 2011 , 98, 1537-48	2.7	22
53	Multivariate analysis of maize disease resistances suggests a pleiotropic genetic basis and implicates a GST gene. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 7339-44	11.5	115
52	Distinct genetic architectures for male and female inflorescence traits of maize. <i>PLoS Genetics</i> , 2011 , 7, e1002383	6	175
51	Allelic Effect Variation at Key Photoperiod Response Quantitative Trait Loci in Maize. <i>Crop Science</i> , 2011 , 51, 1036-1049	2.4	16
50	Seed Dormancy in Mexican Teosinte. <i>Crop Science</i> , 2011 , 51, 2056-2066	2.4	15
49	Mapping QTL Controlling Southern Leaf Blight Resistance by Joint Analysis of Three Related Recombinant Inbred Line Populations. <i>Crop Science</i> , 2011 , 51, 1571-1579	2.4	25
48	QTL Controlling Masculinization of Ear Tips in a Maize (<i>Zea mays</i> L.) Intraspecific Cross. <i>G3: Genes, Genomes, Genetics</i> , 2011 , 1, 337-41	3.2	5
47	Comparison of Conventional, Modified Single Seed Descent, and Doubled Haploid Breeding Methods for Maize Inbred Line Development Using Germplasm Enhancement of Maize Breeding Crosses. <i>Crop Science</i> , 2011 , 51, 1534-1543	2.4	10
46	Direct mapping of density response in a population of B73 x Mo17 recombinant inbred lines of maize (<i>Zea Mays</i> L.). <i>Heredity</i> , 2010 , 104, 583-99	3.6	25
45	Joint Analysis of Near-Isogenic and Recombinant Inbred Line Populations Yields Precise Positional Estimates for Quantitative Trait Loci. <i>Plant Genome</i> , 2010 , 3,	4.4	12

44	Agriculture. Increased food and ecosystem security via perennial grains. <i>Science</i> , 2010 , 328, 1638-9	33.3	303
43	Genetic control of photoperiod sensitivity in maize revealed by joint multiple population analysis. <i>Genetics</i> , 2010 , 184, 799-812	4	88
42	Estimating and Interpreting Heritability for Plant Breeding: An Update 2010 , 9-112		140
41	Selection for Reduced Fusarium Ear Rot and Fumonisin Content in Advanced Backcross Maize Lines and Their Topcross Hybrids. <i>Crop Science</i> , 2010 , 50, 2249-2260	2.4	25
40	Mapping resistance quantitative trait Loci for three foliar diseases in a maize recombinant inbred line population-evidence for multiple disease resistance?. <i>Phytopathology</i> , 2010 , 100, 72-9	3.8	73
39	A heritability-adjusted GGE biplot for test environment evaluation. <i>Euphytica</i> , 2010 , 171, 355-369	2.1	97
38	Genetic properties of the maize nested association mapping population. <i>Science</i> , 2009 , 325, 737-40	33.3	775
37	The genetic architecture of maize flowering time. <i>Science</i> , 2009 , 325, 714-8	33.3	1043
36	Harnessing quantitative genetics and genomics for understanding and improving complex traits in crops 2009 , 123-136		
35	Increasing Yield 2009 , 469-482		1
34	BREEDING FOR IMPROVED RESISTANCE TO FUMONISIN CONTAMINATION IN MAIZE. <i>Toxin Reviews</i> , 2008 , 27, 371-389	2.3	41
33	Genetic design and statistical power of nested association mapping in maize. <i>Genetics</i> , 2008 , 178, 539-514		753
32	Molecular Characterization of Maize Inbreds with Expired U.S. Plant Variety Protection. <i>Crop Science</i> , 2008 , 48, 1673-1685	2.4	57
31	Climatic Adaptation and Ecological Descriptors of 42 Mexican Maize Races. <i>Crop Science</i> , 2008 , 48, 1502-1512	2.4	64
30	Quantitative Trait Loci and Epistasis for Crown Freezing Tolerance in the Kanota Ogle Hexaploid Oat Mapping Population. <i>Crop Science</i> , 2008 , 48, 149-157	2.4	13
29	Johnsongrass (<i>Sorghum halepense</i>) Pollen Expresses Accase Target-site Resistance. <i>Weed Technology</i> , 2007 , 21, 384-388	1.4	6
28	Genotypic Correlation and Multivariate QTL Analyses for Cell Wall Components and Resistance to Stalk Tunneling by the European Corn Borer in Maize. <i>Crop Science</i> , 2007 , 47, 485-488	2.4	18
27	Relationships of Resistance to Fusarium Ear Rot and Fumonisin Contamination with Agronomic Performance of Maize. <i>Crop Science</i> , 2007 , 47, 1770-1778	2.4	19

26	Mapping reciprocal effects and interactions with plant density stress in <i>Zea mays</i> L. <i>Heredity</i> , 2007 , 99, 14-30	3.6	14
25	Genetic architecture of complex traits in plants. <i>Current Opinion in Plant Biology</i> , 2007 , 10, 156-61	9.9	310
24	Mapping resistance to Southern rust in a tropical by temperate maize recombinant inbred topcross population. <i>Theoretical and Applied Genetics</i> , 2007 , 114, 659-67	6	31
23	QTL mapping with near-isogenic lines in maize. <i>Theoretical and Applied Genetics</i> , 2007 , 114, 1211-28	6	93
22	Precise mapping of quantitative trait loci for resistance to southern leaf blight, caused by <i>Cochliobolus heterostrophus</i> race O, and flowering time using advanced intercross maize lines. <i>Genetics</i> , 2007 , 176, 645-57	4	88
21	Relationships among resistances to Fusarium and Aspergillus ear rots and contamination by fumonisin and aflatoxin in maize. <i>Phytopathology</i> , 2007 , 97, 311-7	3.8	44
20	Panzea: a database and resource for molecular and functional diversity in the maize genome. <i>Nucleic Acids Research</i> , 2006 , 34, D752-7	20.1	74
19	Mapping density response in maize: a direct approach for testing genotype and treatment interactions. <i>Genetics</i> , 2006 , 173, 331-48	4	18
18	Heritabilities and Correlations of Fusarium Ear Rot Resistance and Fumonisin Contamination Resistance in Two Maize Populations. <i>Crop Science</i> , 2006 , 46, 353-361	2.4	88
17	QTL Mapping for Fusarium Ear Rot and Fumonisin Contamination Resistance in Two Maize Populations. <i>Crop Science</i> , 2006 , 46, 1734-1743	2.4	101
16	Responses to Selection for Partial Resistance to Crown Rust in Oat. <i>Crop Science</i> , 2006 , 46, 1260-1265	2.4	12
15	Estimating Genotypic Correlations and Their Standard Errors Using Multivariate Restricted Maximum Likelihood Estimation with SAS Proc MIXED. <i>Crop Science</i> , 2006 , 46, 642-654	2.4	230
14	Identification of quantitative trait Loci for resistance to southern leaf blight and days to anthesis in a maize recombinant inbred line population. <i>Phytopathology</i> , 2006 , 96, 1067-71	3.8	33
13	A unified mixed-model method for association mapping that accounts for multiple levels of relatedness. <i>Nature Genetics</i> , 2006 , 38, 203-8	36.3	2595
12	Marker-Assisted Breeding for Host Resistance to Mycotoxin Contamination. <i>Food Additives</i> , 2005 , 423-436		9
11	Genomic regions controlling vernalization and photoperiod responses in oat. <i>Theoretical and Applied Genetics</i> , 2002 , 105, 113-126	6	54
10	Correlated Responses to Selection for Greater β -Glucan Content in Two Oat Populations. <i>Crop Science</i> , 2002 , 42, 730	2.4	10
9	Correlated responses of fatty acid composition, grain quality, and agronomic traits to nine cycles of recurrent selection for increased oil content in oat. <i>Euphytica</i> , 2001 , 122, 69-79	2.1	36

8	Genetic Relationships of Crown Rust Resistance, Grain Yield, Test Weight, and Seed Weight in Oat. <i>Crop Science</i> , 2001 , 41, 1041-1050	2.4	28
7	Inheritance of resistance to southern corn rust in tropical-by-corn-belt maize populations. <i>Theoretical and Applied Genetics</i> , 1998 , 96, 232-241	6	36
6	Genetic Improvement for Yield and Fertility of Alfalfa Cultivars Representing Different Eras of Breeding. <i>Crop Science</i> , 1994 , 34, 953-957	2.4	47
5	Genetic Analysis of Lodging in Diverse Maize Hybrids		4
4	Heterosis of leaf and rhizosphere microbiomes in field-grown maize		2
3	Eleven biosynthetic genes explain the majority of natural variation for carotenoid levels in maize grain		2
2	Dedication: Major M. Goodman: Maize Geneticist and Breeder1-29		
1	Epistasis and Plant Breeding27-92		14