

Andrew Tom Bates

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4476761/publications.pdf>

Version: 2024-02-01

9
papers

476
citations

1162889
8
h-index

1474057
9
g-index

9
all docs

9
docs citations

9
times ranked

959
citing authors

#	ARTICLE	IF	CITATIONS
1	Breast Cancer Risk After Supradiaphragmatic Radiotherapy for Hodgkin's Lymphoma in England and Wales: A National Cohort Study. <i>Journal of Clinical Oncology</i> , 2012, 30, 2745-2752.	0.8	142
2	Randomized Trial of Erlotinib Plus Whole-Brain Radiotherapy for NSCLC Patients With Multiple Brain Metastases. <i>Journal of the National Cancer Institute</i> , 2014, 106, .	3.0	105
3	Effect of Single-Fraction vs Multifraction Radiotherapy on Ambulatory Status Among Patients With Spinal Canal Compression From Metastatic Cancer. <i>JAMA - Journal of the American Medical Association</i> , 2019, 322, 2084.	3.8	71
4	Study protocol for the SARON trial: a multicentre, randomised controlled phase III trial comparing the addition of stereotactic ablative radiotherapy and radical radiotherapy with standard chemotherapy alone for oligometastatic non-small cell lung cancer. <i>BMJ Open</i> , 2018, 8, e020690.	0.8	56
5	Risk of Premature Menopause After Treatment for Hodgkin's Lymphoma. <i>Journal of the National Cancer Institute</i> , 2014, 106, .	3.0	48
6	PSA doubling time kinetics during prostate cancer biochemical relapse after external beam radiation therapy. <i>International Journal of Radiation Oncology Biology Physics</i> , 2005, 62, 148-153.	0.4	19
7	Variation in geographical treatment intensity affects survival of non-small cell lung cancer patients in England. <i>Cancer Epidemiology</i> , 2018, 57, 13-23.	0.8	18
8	Short duration immunochemotherapy followed by radioimmunotherapy consolidation is effective and well tolerated in relapsed follicular lymphoma: 5-year results from a UK National Cancer Research Institute Lymphoma Group study. <i>British Journal of Haematology</i> , 2016, 173, 274-282.	1.2	12
9	Fractionated 90y Ibritumomab Tiuxetan (Zevalin, ^{90}Y) Radioimmunotherapy As An Initial Therapy of Follicular Lymphoma - First Results From a Phase II Study in Patients Requiring Treatment According to GELF/BNLI Criteria. <i>Blood</i> , 2011, 118, 102-102.	0.6	5