

Julia Hankel

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4477253/publications.pdf>

Version: 2024-02-01

16
papers

150
citations

1307594

7
h-index

1199594

12
g-index

16
all docs

16
docs citations

16
times ranked

235
citing authors

#	ARTICLE	IF	CITATIONS
1	Lauric acid as feed additive – An approach to reducing <i>Campylobacter</i> spp. in broiler meat. PLoS ONE, 2017, 12, e0175693.	2.5	34
2	Caecal Microbiota of Experimentally <i>Campylobacter jejuni</i> -Infected Chickens at Different Ages. Frontiers in Microbiology, 2019, 10, 2303.	3.5	19
3	Influence of lauric acid on the susceptibility of chickens to an experimental <i>Campylobacter jejuni</i> colonisation. PLoS ONE, 2018, 13, e0204483.	2.5	16
4	Influence of a specific amino acid pattern in the diet on the course of an experimental <i>Campylobacter jejuni</i> infection in broilers. Poultry Science, 2018, 97, 4020-4030.	3.4	15
5	Intestinal Microbiota of Fattening Pigs Offered Non-Fermented and Fermented Liquid Feed with and without the Supplementation of Non-Fermented Coarse Cereals. Microorganisms, 2020, 8, 638.	3.6	15
6	Performance, Fermentation Characteristics and Composition of the Microbiome in the Digest of Piglets Kept on a Feed With Humic Acid-Rich Peat. Frontiers in Veterinary Science, 2019, 6, 29.	2.2	12
7	The Effects of Feed Particle Size and Floor Type on the Growth Performance, GIT Development, and Pododermatitis in Broiler Chickens. Animals, 2020, 10, 1256.	2.3	10
8	Faecal Microbiota of Dogs Offered a Vegetarian Diet with or without the Supplementation of Feather Meal and either Cornmeal, Rye or Fermented Rye: A Preliminary Study. Microorganisms, 2020, 8, 1363.	3.6	6
9	Influence of different protein sources in the broiler diet on the presence of <i>Campylobacter</i> spp. in excreta and caecal content. Journal of Animal Physiology and Animal Nutrition, 2017, 101, 95-104.	2.2	5
10	Comparison of Chicken Cecal Microbiota after Metaphylactic Treatment or Following Administration of Feed Additives in a Broiler Farm with Enterococcal Spondylitis History. Pathogens, 2021, 10, 1068.	2.8	5
11	Feed Choice Led to Higher Protein Intake in Broiler Chickens Experimentally Infected With <i>Campylobacter jejuni</i> . Frontiers in Nutrition, 2018, 5, 79.	3.7	4
12	Feeding a <i>Saccharomyces cerevisiae</i> Fermentation Product (Olimond BB) Does Not Alter the Fecal Microbiota of Thoroughbred Racehorses. Animals, 2022, 12, 1496.	2.3	4
13	Energy Consumption of Young Military Working Dogs in Pre-Training in Germany. Animals, 2020, 10, 1753.	2.3	3
14	Effects of a carbohydrase complex added in different inclusion rates in feeds for broilers on growth performance, digesta viscosity and foot pad health. Journal of Animal Physiology and Animal Nutrition, 2017, 101, 105-109.	2.2	1
15	Fermentation Characteristics of Rye and Sorghum Depending on Water:Feed Ratio. Fermentation, 2022, 8, 155.	3.0	1
16	Mitigating the Spread and Translocation of <i>Salmonella</i> Enteritidis in Experimentally Infected Broilers under the Influence of Different Flooring Housing Systems and Feed Particle Sizes. Microorganisms, 2021, 9, 874.	3.6	0