

# Cheol-Keun Park

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/448489/publications.pdf>

Version: 2024-02-01

29  
papers

530  
citations

623734

14  
h-index

677142

22  
g-index

29  
all docs

29  
docs citations

29  
times ranked

1261  
citing authors

#	ARTICLE	IF	CITATIONS
1	Down-regulation of TRPV6 Is Associated With Adverse Prognosis in Hepatocellular Carcinoma Treated With Curative Resection. <i>Cancer Genomics and Proteomics</i> , 2022, 19, 259-269.	2.0	1
2	Validation of ZMYND8 as a new treatment target in hepatocellular carcinoma. <i>Journal of Cancer Research and Clinical Oncology</i> , 2021, 147, 3517-3534.	2.5	5
3	Expression of Pregnancy Up-regulated Non-ubiquitous Calmodulin Kinase (PNCK) in Hepatocellular Carcinoma. <i>Cancer Genomics and Proteomics</i> , 2020, 17, 747-755.	2.0	15
4	Hepatic Angiosarcoma: Clinicopathologic Study With an Investigation of ROS1 Gene Rearrangements. <i>In Vivo</i> , 2020, 34, 1463-1467.	1.3	2
5	Prognostic effect of preoperative neutrophil-lymphocyte ratio is related with tumor necrosis and tumor-infiltrating lymphocytes in hepatocellular carcinoma. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2020, 477, 807-816.	2.8	18
6	Clinical Significance of Trk Receptor Expression as a New Therapeutic Target in Hepatocellular Carcinoma. <i>Pathology and Oncology Research</i> , 2020, 26, 2587-2595.	1.9	4
7	Evaluation of the American Joint Committee on Cancer (AJCC) 8th Edition Staging System for Hepatocellular Carcinoma in 1,008 Patients with Curative Resection. <i>Cancer Research and Treatment</i> , 2020, 52, 1145-1152.	3.0	24
8	Prognostic significance of miR-122 expression after curative resection in patients with hepatocellular carcinoma. <i>Scientific Reports</i> , 2019, 9, 14738.	3.3	18
9	Prognostic Role of Apelin Receptor Expression in Hepatocellular Carcinoma Treated With Curative Surgical Resection. <i>Anticancer Research</i> , 2019, 39, 3025-3031.	1.1	14
10	CAMTA-1 Expression in 24 Cases of Hepatic Epithelioid Hemangioendothelioma in a Single Institute: Diagnostic Utility for Differential Diagnosis from Hepatic Angiosarcoma. <i>In Vivo</i> , 2019, 33, 2293-2297.	1.3	10
11	Clinical importance of TERT overexpression in hepatocellular carcinoma treated with curative surgical resection in HBV endemic area. <i>Scientific Reports</i> , 2017, 7, 12258.	3.3	11
12	Nonlinear tumor evolution from dysplastic nodules to hepatocellular carcinoma. <i>Oncotarget</i> , 2017, 8, 2076-2082.	1.8	8
13	Genes co-amplified with <i>ERBB2</i> or <i>MET</i> as novel potential cancer-promoting genes in gastric cancer. <i>Oncotarget</i> , 2017, 8, 92209-92226.	1.8	26
14	Hepatocyte homeostasis for chromosome ploidy and liver function is regulated by Ssu72 protein phosphatase. <i>Hepatology</i> , 2016, 63, 247-259.	7.3	23
15	Expression of DBC1 is associated with poor prognosis in hepatitis virus-related hepatocellular carcinoma. <i>Pathology Research and Practice</i> , 2016, 212, 616-621.	2.3	13
16	Loss of Tuberous Sclerosis Complex 2 (TSC2) as a Predictive Biomarker of Response to mTOR Inhibitor Treatment in Patients with Hepatocellular Carcinoma. <i>Translational Oncology</i> , 2016, 9, 466-471.	3.7	13
17	The Prognostic Role of Mitotic Index in Hepatocellular Carcinoma Patients after Curative Hepatectomy. <i>Cancer Research and Treatment</i> , 2016, 48, 180-189.	3.0	19
18	The Overexpression of CCAR1 in Hepatocellular Carcinoma Associates with Poor Prognosis. <i>Cancer Research and Treatment</i> , 2016, 48, 1065-1073.	3.0	15

#	ARTICLE	IF	CITATIONS
19	NADPH Oxidase 1 and NADPH Oxidase 4 Have Opposite Prognostic Effects for Patients with Hepatocellular Carcinoma after Hepatectomy. <i>Gut and Liver</i> , 2016, 10, 826-835.	2.9	26
20	HULC and H19 Played Different Roles in Overall and Disease-Free Survival from Hepatocellular Carcinoma after Curative Hepatectomy: A Preliminary Analysis from Gene Expression Omnibus. <i>Disease Markers</i> , 2015, 2015, 1-9.	1.3	66
21	Two sample test for high-dimensional partially paired data. <i>Journal of Applied Statistics</i> , 2015, 42, 1946-1961.	1.3	3
22	Microsteatosis may not interact with macrosteatosis in living donor liver transplantation. <i>Journal of Hepatology</i> , 2015, 62, 556-562.	3.7	28
23	The prognosis of hepatocellular carcinoma after curative hepatectomy in young patients. <i>Oncotarget</i> , 2015, 6, 18664-18673.	1.8	18
24	Expression of PEG10 Is Associated with Poor Survival and Tumor Recurrence in Hepatocellular Carcinoma. <i>Cancer Research and Treatment</i> , 2015, 47, 844-852.	3.0	39
25	ATAD2 as a Poor Prognostic Marker for Hepatocellular Carcinoma after Curative Resection. <i>Cancer Research and Treatment</i> , 2015, 47, 853-861.	3.0	35
26	Gastroenteropancreatic Neuroendocrine Tumors with Liver Metastases in Korea: A Clinicopathological Analysis of 72 Cases in a Single Institute. <i>Cancer Research and Treatment</i> , 2015, 47, 738-746.	3.0	10
27	High Expression of Ribonucleotide Reductase Subunit M2 Correlates with Poor Prognosis of Hepatocellular Carcinoma. <i>Gut and Liver</i> , 2014, 8, 662-668.	2.9	36
28	High Expression of Aldo-Keto Reductase 1B10 Is an Independent Predictor of Favorable Prognosis in Patients with Hepatocellular Carcinoma. <i>Gut and Liver</i> , 2014, 8, 648-654.	2.9	24
29	Primary hepatic neuroendocrine carcinoma. <i>Korean Journal of Hepato-biliary-pancreatic Surgery</i> , 2013, 17, 34.	1.0	6