

Peter Reeh

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4486581/publications.pdf>

Version: 2024-02-01

212
papers

15,231
citations

13068

68
h-index

20900

115
g-index

217
all docs

217
docs citations

217
times ranked

11062
citing authors

#	ARTICLE	IF	CITATIONS
1	Cannabinoids mediate analgesia largely via peripheral type 1 cannabinoid receptors in nociceptors. <i>Nature Neuroscience</i> , 2007, 10, 870-879.	7.1	504
2	Protons selectively induce lasting excitation and sensitization to mechanical stimulation of nociceptors in rat skin, in vitro. <i>Journal of Neuroscience</i> , 1992, 12, 86-95.	1.7	477
3	Methylglyoxal modification of Nav1.8 facilitates nociceptive neuron firing and causes hyperalgesia in diabetic neuropathy. <i>Nature Medicine</i> , 2012, 18, 926-933.	15.2	414
4	TREK-1, a K ⁺ channel involved in polymodal pain perception. <i>EMBO Journal</i> , 2006, 25, 2368-2376.	3.5	363
5	TRPA1 channels mediate acute neurogenic inflammation and pain produced by bacterial endotoxins. <i>Nature Communications</i> , 2014, 5, 3125.	5.8	361
6	Sensory neuron sodium channel Nav1.8 is essential for pain at low temperatures. <i>Nature</i> , 2007, 447, 856-859.	13.7	355
7	H ₂ S and NO cooperatively regulate vascular tone by activating a neuroendocrine HNO ⁺ “TRPA1”CGRP signalling pathway. <i>Nature Communications</i> , 2014, 5, 4381.	5.8	324
8	The mechano-activated K ⁺ channels TRAAK and TREK-1 control both warm and cold perception. <i>EMBO Journal</i> , 2009, 28, 1308-1318.	3.5	309
9	Chemosensitivity of fine afferents from rat skin in vitro. <i>Journal of Neurophysiology</i> , 1990, 63, 887-901.	0.9	298
10	Responsiveness and functional attributes of electrically localized terminals of cutaneous C-fibers in vivo and in vitro. <i>Journal of Neurophysiology</i> , 1992, 68, 581-595.	0.9	289
11	Sensory receptors in mammalian skin in an in vitro preparation. <i>Neuroscience Letters</i> , 1986, 66, 141-146.	1.0	272
12	A dominant role of acid pH in inflammatory excitation and sensitization of nociceptors in rat skin, in vitro. <i>Journal of Neuroscience</i> , 1995, 15, 3982-3989.	1.7	268
13	Selective excitation by capsaicin of mechano-heat sensitive nociceptors in rat skin. <i>Brain Research</i> , 1988, 446, 262-268.	1.1	242
14	Sensory and Signaling Mechanisms of Bradykinin, Eicosanoids, Platelet-Activating Factor, and Nitric Oxide in Peripheral Nociceptors. <i>Physiological Reviews</i> , 2012, 92, 1699-1775.	13.1	239
15	Chapter 8. Tissue acidosis in nociception and pain. <i>Progress in Brain Research</i> , 1996, 113, 143-151.	0.9	236
16	Pain due to tissue acidosis: a mechanism for inflammatory and ischemic myalgia?. <i>Neuroscience Letters</i> , 1996, 208, 191-194.	1.0	215
17	Does neurogenic inflammation alter the sensitivity of unmyelinated nociceptors in the rat?. <i>Brain Research</i> , 1986, 384, 42-50.	1.1	204
18	<i>Sox10</i> is required for Schwann cell identity and progression beyond the immature Schwann cell stage. <i>Journal of Cell Biology</i> , 2010, 189, 701-712.	2.3	198

#	ARTICLE	IF	CITATIONS
19	TRPA1 and Substance P Mediate Colitis in Mice. <i>Gastroenterology</i> , 2011, 141, 1346-1358.	0.6	197
20	Sustained graded pain and hyperalgesia from harmless experimental tissue acidosis in human skin. <i>Neuroscience Letters</i> , 1993, 154, 113-116.	1.0	181
21	Release of substance P, calcitonin gene-related peptide and prostaglandin E2 from rat dura mater encephali following electrical and chemical stimulation in vitro. <i>Neuroscience</i> , 1999, 89, 901-907.	1.1	178
22	The nociceptor sensitization by bradykinin does not depend on sympathetic neurons. <i>Neuroscience</i> , 1992, 46, 465-473.	1.1	174
23	Methylglyoxal Activates Nociceptors through Transient Receptor Potential Channel A1 (TRPA1). <i>Journal of Biological Chemistry</i> , 2012, 287, 28291-28306.	1.6	166
24	Pattern of monosynaptic Ia connections in the cat forelimb.. <i>Journal of Physiology</i> , 1989, 419, 321-351.	1.3	155
25	Phenotyping sensory nerve endings in vitro in the mouse. <i>Nature Protocols</i> , 2009, 4, 174-196.	5.5	152
26	Unresponsive afferent nerve fibres in the sural nerve of the rat.. <i>Journal of Physiology</i> , 1991, 435, 229-242.	1.3	151
27	Molecular physiology of proton transduction in nociceptors. <i>Current Opinion in Pharmacology</i> , 2001, 1, 45-51.	1.7	151
28	Variable sensitivity to noxious heat is mediated by differential expression of the CGRP gene. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 12938-12943.	3.3	151
29	Excitation of cutaneous afferent nerve endings in vitro by a combination of inflammatory mediators and conditioning effect of substance P. <i>Experimental Brain Research</i> , 1992, 91, 467-76.	0.7	144
30	The vanilloid receptor TRPV1 is activated and sensitized by local anesthetics in rodent sensory neurons. <i>Journal of Clinical Investigation</i> , 2008, 118, 763-76.	3.9	134
31	New mechanism underlying IL-31-induced atopic dermatitis. <i>Journal of Allergy and Clinical Immunology</i> , 2018, 141, 1677-1689.e8.	1.5	131
32	The effect of carrageenan-induced inflammation on the sensitivity of unmyelinated skin nociceptors in the rat. <i>Pain</i> , 1987, 29, 363-373.	2.0	130
33	Beyond H ₂ S and NO Interplay: Hydrogen Sulfide and Nitroprusside React Directly to Give Nitroxyl (HNO). A New Pharmacological Source of HNO. <i>Journal of Medicinal Chemistry</i> , 2013, 56, 1499-1508.	2.9	126
34	Actions of cholinergic agonists and antagonists on sensory nerve endings in rat skin, in vitro. <i>Journal of Neurophysiology</i> , 1993, 70, 397-405.	0.9	122
35	Sustained sensitization and recruitment of rat cutaneous nociceptors by bradykinin and a novel theory of its excitatory action. <i>Journal of Physiology</i> , 2001, 532, 229-239.	1.3	119
36	Rat peripheral nerve components release calcitonin gene-related peptide and prostaglandin E2 in response to noxious stimuli: evidence that nervi nervorum are nociceptors. <i>Neuroscience</i> , 1999, 92, 319-325.	1.1	117

#	ARTICLE	IF	CITATIONS
37	<i>Sox10</i> is required for Schwann cell homeostasis and myelin maintenance in the adult peripheral nerve. <i>Glia</i> , 2011, 59, 1022-1032.	2.5	113
38	Interactions of inflammatory mediators stimulating release of calcitonin gene-related peptide, substance P and prostaglandin E2 from isolated rat skin. <i>Neuropharmacology</i> , 2001, 40, 416-423.	2.0	111
39	Sodium channelopathies and pain. <i>Pflugers Archiv European Journal of Physiology</i> , 2010, 460, 249-263.	1.3	110
40	Excitatory Nicotinic and Desensitizing Muscarinic (M2) Effects on C-Nociceptors in Isolated Rat Skin. <i>Journal of Neuroscience</i> , 2001, 21, 3295-3302.	1.7	108
41	Discharge patterns of afferent cutaneous nerve fibers from the rat's tail during prolonged noxious mechanical stimulation. <i>Experimental Brain Research</i> , 1987, 65, 493-504.	0.7	106
42	Pain due to experimental acidosis in human skin: evidence for non-adapting nociceptor excitation. <i>Neuroscience Letters</i> , 1995, 199, 29-32.	1.0	106
43	Inflammatory mediators potentiate pain induced by experimental tissue acidosis. <i>Pain</i> , 1996, 66, 163-170.	2.0	105
44	Activation of TRPM3 by a potent synthetic ligand reveals a role in peptide release. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015, 112, E1363-72.	3.3	105
45	Inflammatory Mediators at Acidic pH Activate Capsaicin Receptors in Cultured Sensory Neurons From Newborn Rats. <i>Journal of Neurophysiology</i> , 1998, 79, 670-676.	0.9	103
46	Ciguatoxins activate specific cold pain pathways to elicit burning pain from cooling. <i>EMBO Journal</i> , 2012, 31, 3795-3808.	3.5	103
47	5,6-EET Is Released upon Neuronal Activity and Induces Mechanical Pain Hypersensitivity via TRPA1 on Central Afferent Terminals. <i>Journal of Neuroscience</i> , 2012, 32, 6364-6372.	1.7	103
48	Human TRPA1 is a heat sensor displaying intrinsic U-shaped thermosensitivity. <i>Scientific Reports</i> , 2016, 6, 28763.	1.6	103
49	Direct evidence for functional TRPV1/TRPA1 heteromers. <i>Pflugers Archiv European Journal of Physiology</i> , 2014, 466, 2229-2241.	1.3	98
50	Sensitization of nociceptive cutaneous nerve fibers from the rat's tail by noxious mechanical stimulation. <i>Experimental Brain Research</i> , 1987, 65, 505-12.	0.7	97
51	Calcitonin gene-related peptide and prostaglandin E2 but not substance P release induced by antidromic nerve stimulation from rat skin in vitro. <i>Neuroscience</i> , 1999, 89, 303-310.	1.1	95
52	Muscarinic receptor subtypes mediating central and peripheral antinociception studied with muscarinic receptor knockout mice. <i>Life Sciences</i> , 2003, 72, 2047-2054.	2.0	93
53	Local Anesthetic-like Inhibition of Voltage-gated Na ⁺ Channels by the Partial μ -opioid Receptor Agonist Buprenorphine. <i>Anesthesiology</i> , 2012, 116, 1335-1346.	1.3	88
54	Transient opening of the perineurial barrier for analgesic drug delivery. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012, 109, E2018-27.	3.3	87

#	ARTICLE	IF	CITATIONS
55	Phenotyping the Function of TRPV1-Expressing Sensory Neurons by Targeted Axonal Silencing. <i>Journal of Neuroscience</i> , 2013, 33, 315-326.	1.7	86
56	Morphological evidence for functional capsaicin receptor expression and calcitonin gene-related peptide exocytosis in isolated peripheral nerve axons of the mouse. <i>Neuroscience</i> , 2004, 126, 585-590.	1.1	85
57	Role of sensory neurons in colitis: increasing evidence for a neuroimmune link in the gut. <i>Inflammatory Bowel Diseases</i> , 2011, 17, 1030-1033.	0.9	82
58	The Molecular Basis for Species-specific Activation of Human TRPA1 Protein by Protons Involves Poorly Conserved Residues within Transmembrane Domains 5 and 6. <i>Journal of Biological Chemistry</i> , 2013, 288, 20280-20292.	1.6	82
59	Receptors, cells and circuits involved in pruritus of systemic disorders. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2014, 1842, 869-892.	1.8	82
60	Improved superfusion technique for rapid cooling or heating of cultured cells under patch-clamp conditions. <i>Journal of Neuroscience Methods</i> , 2006, 151, 178-185.	1.3	79
61	The General Anesthetic Propofol Excites Nociceptors by Activating TRPV1 and TRPA1 Rather than GABAA Receptors. <i>Journal of Biological Chemistry</i> , 2010, 285, 34781-34792.	1.6	79
62	Sensitized peripheral nociception in experimental diabetes of the rat. <i>Pain</i> , 2010, 151, 496-505.	2.0	78
63	Systemic desensitization through TRPA1 channels by capsazepine and mustard oil - a novel strategy against inflammation and pain. <i>Scientific Reports</i> , 2016, 6, 28621.	1.6	78
64	Rises in [Ca ²⁺] _i mediate capsaicin- and proton-induced heat sensitization of rat primary nociceptive neurons. <i>European Journal of Neuroscience</i> , 1999, 11, 3143-3150.	1.2	77
65	The proximodistal aggravation of colitis depends on substance P released from TRPV1-expressing sensory neurons. <i>Journal of Gastroenterology</i> , 2012, 47, 256-265.	2.3	75
66	A technique for fast application of heated solutions of different composition to cultured neurones. <i>Journal of Neuroscience Methods</i> , 1998, 82, 195-201.	1.3	72
67	Heat-induced release of CGRP from isolated rat skin and effects of bradykinin and the protein kinase C activator PMA. <i>Pain</i> , 1999, 83, 289-295.	2.0	72
68	The interphase of the formalin test. <i>Pain</i> , 2014, 155, 511-521.	2.0	71
69	Protease Activated Receptors 1 and 4 Sensitize TRPV1 in Nociceptive Neurones. <i>Molecular Pain</i> , 2010, 6, 1744-8069-6-61.	1.0	69
70	Injection pain of rocuronium and vecuronium is evoked by direct activation of nociceptive nerve endings. <i>European Journal of Anaesthesiology</i> , 2003, 20, 245-253.	0.7	67
71	Muscarinic M2 Receptors on Peripheral Nerve Endings: A Molecular Target of Antinociception. <i>Journal of Neuroscience</i> , 2002, 22, RC229-RC229.	1.7	66
72	Photosensitization in Porphyrins and Photodynamic Therapy Involves TRPA1 and TRPV1. <i>Journal of Neuroscience</i> , 2016, 36, 5264-5278.	1.7	66

#	ARTICLE	IF	CITATIONS
73	Transient receptor potential melastatin 8 ion channel in macrophages modulates colitis through a balance-shift in TNF-alpha and interleukin-10 production. <i>Mucosal Immunology</i> , 2016, 9, 1500-1513.	2.7	65
74	Stable analogues of cyclic AMP but not cyclic GMP sensitize unmyelinated primary afferents in rat skin to heat stimulation but not to inflammatory mediators, in vitro. <i>Neuroscience</i> , 1996, 74, 609-617.	1.1	64
75	The TRPV1/2/3 activator 2-aminoethoxydiphenyl borate sensitizes native nociceptive neurons to heat in wildtype but not TRPV1 deficient mice. <i>Neuroscience</i> , 2005, 135, 1277-1284.	1.1	64
76	ATP can enhance the proton-induced CGRP release through P2Y receptors and secondary PGE2 release in isolated rat dura mater. <i>Pain</i> , 2002, 97, 259-265.	2.0	60
77	Proton-induced calcitonin gene-related peptide release from rat sciatic nerve axons, in vitro, involving TRPV1. <i>European Journal of Neuroscience</i> , 2003, 18, 803-810.	1.2	60
78	A high-threshold heat-activated channel in cultured rat dorsal root ganglion neurons resembles TRPV2 and is blocked by gadolinium. <i>European Journal of Neuroscience</i> , 2007, 26, 12-22.	1.2	60
79	Injection pain of rocuronium and vecuronium is evoked by direct activation of nociceptive nerve endings. <i>European Journal of Anaesthesiology</i> , 2003, 20, 245-253.	0.7	59
80	Streptozotocin Stimulates the Ion Channel TRPA1 Directly. <i>Journal of Biological Chemistry</i> , 2015, 290, 15185-15196.	1.6	59
81	HCN2 channels account for mechanical (but not heat) hyperalgesia during long-standing inflammation. <i>Pain</i> , 2014, 155, 1079-1090.	2.0	58
82	Nociceptor excitation by thermal sensitization â€” A hypothesis. <i>Progress in Brain Research</i> , 2000, 129, 39-50.	0.9	57
83	Substance P, calcitonin gene related peptide and PGE2 co-released from the mouse colon: a new model to study nociceptive and inflammatory responses in viscera, in vitro. <i>Pain</i> , 2001, 93, 213-219.	2.0	56
84	Recordings From Brain Stem Neurons Responding to Chemical Stimulation of the Subarachnoid Space. <i>Journal of Neurophysiology</i> , 1997, 77, 3122-3133.	0.9	55
85	Bupivacaineâ€”induced cellular entry of Ca^{2+} and its contribution to differential nerve block. <i>British Journal of Pharmacology</i> , 2014, 171, 438-451.	2.7	55
86	Inflammatory pain control by blocking oxidized phospholipid-mediated TRP channel activation. <i>Scientific Reports</i> , 2017, 7, 5447.	1.6	53
87	Bradykinin-induced nociceptor sensitization to heat is mediated by cyclooxygenase products in isolated rat skin. <i>European Journal of Neuroscience</i> , 2001, 14, 210-218.	1.2	52
88	TRPV1, TRPA1, and CB1 in the isolated vagus nerve â€” Axonal chemosensitivity and control of neuropeptide release. <i>Neuropeptides</i> , 2011, 45, 391-400.	0.9	52
89	Differential Contribution of TRPA1, TRPV4 and TRPM8 to Colonic Nociception in Mice. <i>PLoS ONE</i> , 2015, 10, e0128242.	1.1	52
90	An interaction of inflammatory mediators and protons in small diameter dorsal root ganglion neurons of the rat. <i>Neuroscience Letters</i> , 1997, 224, 37-40.	1.0	51

#	ARTICLE	IF	CITATIONS
91	Pro- and anti-inflammatory actions of ricinoleic acid: similarities and differences with capsaicin. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2001, 364, 87-95.	1.4	51
92	Analgesic treatment of ciguatoxin-induced cold allodynia. <i>Pain</i> , 2013, 154, 1999-2006.	2.0	51
93	Low pH facilitates capsaicin responses in isolated sensory neurons of the rat. <i>Neuroscience Letters</i> , 1996, 211, 5-8.	1.0	49
94	Topical acetylsalicylic, salicylic acid and indomethacin suppress pain from experimental tissue acidosis in human skin. <i>Pain</i> , 1995, 62, 339-347.	2.0	48
95	Noxious heat-induced CGRP release from rat sciatic nerve axons in vitro. <i>European Journal of Neuroscience</i> , 2001, 14, 1203-1208.	1.2	47
96	Calcitonin gene-related peptide release from intact isolated dorsal root and trigeminal ganglia. <i>Neuropeptides</i> , 2008, 42, 311-317.	0.9	47
97	Carrageenan inflammation increases bradykinin sensitivity of rat cutaneous nociceptors. <i>Neuroscience Letters</i> , 1990, 111, 206-210.	1.0	46
98	Role of nitric oxide in zymosan induced paw inflammation and thermal hyperalgesia. <i>Inflammation Research</i> , 2001, 50, 83-88.	1.6	46
99	Angiotensin II facilitates sympathetic transmission in rat hind limb circulation.. <i>Hypertension</i> , 1993, 21, 322-328.	1.3	45
100	Muscarinic M2 receptors inhibit heat-induced CGRP release from isolated rat skin. <i>NeuroReport</i> , 2001, 12, 2457-2460.	0.6	45
101	Differential effects of TRPV channel block on polymodal activation of rat cutaneous nociceptors in vitro. <i>Experimental Brain Research</i> , 2009, 196, 31-44.	0.7	45
102	Opposite effects of substance P and calcitonin gene-related peptide in oxazolone colitis. <i>Digestive and Liver Disease</i> , 2012, 44, 24-29.	0.4	45
103	Measurement of the analgesic effects of aspirin with a new experimental algometric procedure. <i>Pain</i> , 1988, 32, 215-222.	2.0	44
104	Scratching an itch. <i>Nature Neuroscience</i> , 2013, 16, 117-118.	7.1	44
105	Soluble Epoxide Hydrolase Limits Mechanical Hyperalgesia during Inflammation. <i>Molecular Pain</i> , 2011, 7, 1744-8069-7-78.	1.0	43
106	<sc>TRPA1</sc> and <sc>TRPV1</sc> are differentially involved in heat nociception of mice. <i>European Journal of Pain</i> , 2013, 17, 1472-1482.	1.4	43
107	Sensory Transduction in Peripheral Nerve Axons Elicits Ectopic Action Potentials. <i>Journal of Neuroscience</i> , 2008, 28, 6281-6284.	1.7	41
108	The anti-diabetic drug glibenclamide is an agonist of the transient receptor potential Ankyrin 1 (TRPA1) ion channel. <i>European Journal of Pharmacology</i> , 2013, 704, 15-22.	1.7	41

#	ARTICLE	IF	CITATIONS
109	Activated human platelets in plasma excite nociceptors in rat skin, in vitro. <i>Neuroscience Letters</i> , 1994, 170, 103-106.	1.0	38
110	The pH response of rat cutaneous nociceptors correlates with extracellular [Na ⁺] and is increased under amiloride. <i>European Journal of Neuroscience</i> , 1999, 11, 2783-2792.	1.2	38
111	TRPV1 controls acid- and heat-induced calcitonin gene-related peptide release and sensitization by bradykinin in the isolated mouse trachea. <i>European Journal of Neuroscience</i> , 2009, 29, 1896-1904.	1.2	38
112	Tonic Postganglionic Sympathetic Inhibition Induced by Afferent Renal Nerves?. <i>Hypertension</i> , 2012, 59, 467-476.	1.3	38
113	Bimodal Concentration-Response of Nicotine Involves the Nicotinic Acetylcholine Receptor, Transient Receptor Potential Vanilloid Type 1, and Transient Receptor Potential Ankyrin 1 Channels in Mouse Trachea and Sensory Neurons. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2013, 347, 529-539.	1.3	38
114	Conditioning of histamine by bradykinin alters responses of rat nociceptor and human itch sensation. <i>Neuroscience Letters</i> , 1993, 152, 117-120.	1.0	37
115	Sensitization to heat through G-protein-coupled receptor pathways in the isolated sciatic mouse nerve. <i>European Journal of Neuroscience</i> , 2007, 25, 3570-3575.	1.2	37
116	Inhibitory CB1 and activating/desensitizing TRPV1-mediated cannabinoid actions on CGRP release in rodent skin. <i>Neuropeptides</i> , 2011, 45, 229-237.	0.9	37
117	Amplified Cold Transduction in Native Nociceptors by M-Channel Inhibition. <i>Journal of Neuroscience</i> , 2013, 33, 16627-16641.	1.7	37
118	TRPA1 and TRPV1 Antagonists Do Not Inhibit Human Acidosis-Induced Pain. <i>Journal of Pain</i> , 2017, 18, 526-534.	0.7	37
119	Chapter 31 Sensory receptors in a mammalian skin "nerve in vitro" preparation. <i>Progress in Brain Research</i> , 1988, 74, 271-276.	0.9	36
120	Effects of TRPV1 receptor antagonists on stimulated iCGRP release from isolated skin of rats and TRPV1 mutant mice. <i>Pain</i> , 2004, 109, 284-290.	2.0	36
121	Role of different proton-sensitive channels in releasing calcitonin gene-related peptide from isolated hearts of mutant mice. <i>Cardiovascular Research</i> , 2005, 65, 405-410.	1.8	36
122	Establishment of myelinating schwann cells and barrier integrity between central and peripheral nervous systems depend on Sox10. <i>Glia</i> , 2012, 60, 806-819.	2.5	36
123	Cigarette smoke has sensory effects through nicotinic and TRPA1 but not TRPV1 receptors on the isolated mouse trachea and larynx. <i>American Journal of Physiology - Lung Cellular and Molecular Physiology</i> , 2015, 309, L812-L820.	1.3	36
124	Quaternary Lidocaine Derivative QX-314 Activates and Permeates Human TRPV1 and TRPA1 to Produce Inhibition of Sodium Channels and Cytotoxicity. <i>Anesthesiology</i> , 2016, 124, 1153-1165.	1.3	35
125	Stimulated prostaglandin E2 release from rat skin, in vitro. <i>Life Sciences</i> , 1998, 62, 2045-2055.	2.0	33
126	Sodium Channel Na _v 1.8 Underlies TTX-Resistant Axonal Action Potential Conduction in Somatosensory C-Fibers of Distal Cutaneous Nerves. <i>Journal of Neuroscience</i> , 2017, 37, 5204-5214.	1.7	33

#	ARTICLE	IF	CITATIONS
127	Mechanisms of potassium- and capsaicin-induced axonal calcitonin gene-related peptide release: Involvement of L- and T-type calcium channels and TRPV1 but not sodium channels. <i>Neuroscience</i> , 2008, 151, 836-842.	1.1	32
128	High Concentrations of Morphine Sensitize and Activate Mouse Dorsal Root Ganglia via TRPV1 and TRPA1 Receptors. <i>Molecular Pain</i> , 2009, 5, 1744-8069-5-17.	1.0	32
129	Location of motoneurons projecting to the cat distal forelimb. II. Median and ulnar motornuclei. <i>Journal of Comparative Neurology</i> , 1986, 244, 302-312.	0.9	31
130	Modulation of CGRP and PGE2 release from isolated rat skin by $\hat{1}\pm$ -adrenoceptors and $\hat{1}\pm$ -opioid-receptors. <i>NeuroReport</i> , 2001, 12, 2097-2100.	0.6	31
131	Do distinct populations of dorsal root ganglion neurons account for the sensory peptidergic innervation of the kidney?. <i>American Journal of Physiology - Renal Physiology</i> , 2009, 297, F1427-F1434.	1.3	31
132	A New Paradigm to Understand and Treat Diabetic Neuropathy. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2014, 122, 201-207.	0.6	31
133	Crotalphine desensitizes TRPA1 ion channels to alleviate inflammatory hyperalgesia. <i>Pain</i> , 2016, 157, 2504-2516.	2.0	31
134	Effects of classical algogens. <i>Seminars in Neuroscience</i> , 1995, 7, 221-226.	2.3	30
135	Dose-dependent competitive block by topical acetylsalicylic and salicylic acid of low pH-induced cutaneous pain. <i>Pain</i> , 1996, 64, 71-82.	2.0	29
136	Plasma levels after peroral and topical ibuprofen and effects upon low pH-induced cutaneous and muscle pain. <i>European Journal of Pain</i> , 2000, 4, 195-209.	1.4	28
137	Morphological characterization of rat Mas-related G-protein-coupled receptor C and functional analysis of agonists. <i>Neuroscience</i> , 2008, 151, 242-254.	1.1	27
138	Irritant Volatile Anesthetics Induce Neurogenic Inflammation Through TRPA1 and TRPV1 Channels in the Isolated Mouse Trachea. <i>Anesthesia and Analgesia</i> , 2015, 120, 467-471.	1.1	27
139	The roles of TRPV1, TRPA1 and TRPM8 channels in chemical and thermal sensitivity of the mouse oral mucosa. <i>European Journal of Neuroscience</i> , 2018, 47, 201-210.	1.2	27
140	Diltiazem blocks the PH-induced excitation of rat nociceptors together with their mechanical and electrical excitability in vitro. <i>Journal of Neurophysiology</i> , 1996, 75, 1-10.	0.9	26
141	More sensory competence for nociceptive neurons in culture. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996, 93, 14995-14997.	3.3	26
142	Intracutaneous injections of platelets cause acute pain and protracted hyperalgesia. <i>Neuroscience Letters</i> , 1997, 226, 171-174.	1.0	26
143	NaV1.7 and pain: contribution of peripheral nerves. <i>Pain</i> , 2018, 159, 496-506.	2.0	26
144	Chemical Excitation and Sensitization of Nociceptors. , 1994, , 119-131.		26

#	ARTICLE	IF	CITATIONS
145	Electrophysiological characterization of vagal afferents relevant to mucosal nociception in the rat upper oesophagus. <i>Journal of Physiology</i> , 2007, 582, 229-242.	1.3	25
146	Interactions of inflammatory mediators and low pH not influenced by capsazepine in rat cutaneous nociceptors. <i>NeuroReport</i> , 2000, 11, 973-976.	0.6	24
147	Location of median and ulnar motornuclei in the cat. <i>Neuroscience Letters</i> , 1982, 30, 103-108.	1.0	23
148	An oral TRPV1 antagonist attenuates laser radiant heat-evoked potentials and pain ratings from UVB-inflamed and normal skin. <i>British Journal of Clinical Pharmacology</i> , 2013, 75, 404-414.	1.1	23
149	Functional and structural characterization of axonal opioid receptors as targets for analgesia. <i>Molecular Pain</i> , 2016, 12, 174480691662873.	1.0	22
150	Lactate is a potent inhibitor of the capsaicin receptor TRPV1. <i>Scientific Reports</i> , 2016, 6, 36740.	1.6	22
151	Bradykinin-induced nociceptor sensitisation to heat depends on cox-1 and cox-2 in isolated rat skin. <i>Pain</i> , 2007, 130, 14-24.	2.0	20
152	Methylene blue induces ongoing activity in rat cutaneous primary afferents and depolarization of DRG neurons via a photosensitive mechanism. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 1997, 356, 619-625.	1.4	19
153	Interactions of histamine and bradykinin on polymodal C-fibres in isolated rat skin. <i>European Journal of Pain</i> , 2001, 5, 97-106.	1.4	19
154	TRPA1 and TRPV1 are required for lidocaine-evoked calcium influx and neuropeptide release but not cytotoxicity in mouse sensory neurons. <i>PLoS ONE</i> , 2017, 12, e0188008.	1.1	19
155	Bitter taste signaling in tracheal epithelial brush cells elicits innate immune responses to bacterial infection. <i>Journal of Clinical Investigation</i> , 2022, 132, .	3.9	19
156	Effects of oxygen radicals on nociceptive afferents in the rat skin in vitro. <i>Pain</i> , 1995, 62, 87-94.	2.0	18
157	Responsiveness of C-fiber nociceptors to punctate force-controlled stimuli in isolated rat skin: lack of modulation by inflammatory mediators and flurbiprofen. <i>Neuroscience Letters</i> , 2004, 361, 163-167.	1.0	18
158	S(+)-flurbiprofen but not 5-HT1 agonists suppress basal and stimulated CGRP and PGE2 release from isolated rat dura mater. <i>Pain</i> , 2003, 103, 313-320.	2.0	17
159	Why cooling is beneficial: non-linear temperature-dependency of stimulated iCGRP release from isolated rat skin. <i>Pain</i> , 2004, 110, 215-219.	2.0	17
160	The tetrodotoxin-resistant Na ⁺ channel Nav1.8 reduces the potency of local anesthetics in blocking C-fiber nociceptors. <i>Pflügers Archiv European Journal of Physiology</i> , 2010, 459, 751-763.	1.3	17
161	Heat-resistant action potentials require TTX-resistant sodium channels Nav1.8 and Nav1.9. <i>Journal of General Physiology</i> , 2018, 150, 1125-1144.	0.9	17
162	Reduced excitability and impaired nociception in peripheral unmyelinated fibers from Nav1.9-null mice. <i>Pain</i> , 2017, 158, 58-67.	2.0	16

#	ARTICLE	IF	CITATIONS
163	Ciguatoxins Evoke Potent CGRP Release by Activation of Voltage-Gated Sodium Channel Subtypes NaV1.9, NaV1.7 and NaV1.1. <i>Marine Drugs</i> , 2017, 15, 269.	2.2	16
164	Complementary roles of murine NaV1.7, NaV1.8 and NaV1.9 in acute itch signalling. <i>Scientific Reports</i> , 2020, 10, 2326.	1.6	16
165	Formalin Evokes Calcium Transients from the Endoplasmatic Reticulum. <i>PLoS ONE</i> , 2015, 10, e0123762.	1.1	16
166	Electrophysiological and Neurochemical Techniques to Investigate Sensory Neurons in Analgesia Research. <i>Methods in Molecular Biology</i> , 2010, 617, 237-259.	0.4	15
167	Analgesic Effect of Acetylsalicylic Acid (ASA) Versus a Lithium-ASA combination: An Evoked Potential Study Employing Radiant Heat Stimulation with a CO ₂ Laser. <i>Pharmacopsychiatry</i> , 1987, 20, 217-221.	1.7	14
168	Pulmonary serotonin 5-HT ₃ -sensitive afferent fibers modulate renal sympathetic nerve activity in rats. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 1997, 272, H979-H986.	1.5	13
169	Denervation and NK1 receptor block modulate stimulated CGRP and PGE2 release from rat skin. <i>NeuroReport</i> , 2000, 11, 283-286.	0.6	13
170	The effects of excessive heat on heat-activated membrane currents in cultured dorsal root ganglia neurons from neonatal rat. <i>Pain</i> , 2002, 95, 207-214.	2.0	13
171	Inflammatory mediators do not stimulate CGRP release if prostaglandin synthesis is blocked by S(+)-flurbiprofen in isolated rat skin. <i>Inflammation Research</i> , 2003, 52, 519-523.	1.6	13
172	Acid-induced CGRP release from the stomach does not depend on TRPV1 or ASIC3. <i>Neurogastroenterology and Motility</i> , 2010, 22, 680-687.	1.6	13
173	TRPA1-dependent calcium transients and CGRP release in DRG neurons require extracellular calcium. <i>Journal of Cell Biology</i> , 2020, 219, .	2.3	13
174	Nobel somatosensations and pain. <i>Pflügers Archiv European Journal of Physiology</i> , 2022, 474, 405-420.	1.3	13
175	Release of calcitonin gene-related peptide from the isolated mouse heart: Methodological validation of a new model. <i>Neuropeptides</i> , 2006, 40, 107-113.	0.9	12
176	Vitamin B complex attenuated heat hyperalgesia following infraorbital nerve constriction in rats and reduced capsaicin in vivo and in vitro effects. <i>European Journal of Pharmacology</i> , 2015, 762, 326-332.	1.7	12
177	A Randomized, Double-Blind, Placebo- and Active Comparator-Controlled Phase I Study of Analgesic/Antihyperalgesic Properties of ASP8477, a Fatty Acid Amide Hydrolase Inhibitor, in Healthy Female Subjects. <i>Pain Medicine</i> , 2018, 19, 1206-1218.	0.9	12
178	Local NGF and GDNF levels modulate morphology and function of porcine DRG neurites, In Vitro. <i>PLoS ONE</i> , 2018, 13, e0203215.	1.1	12
179	Comparison of Two Different Preparations of Ibuprofen with Regard to the Time Course of Their Analgesic Effect. <i>Arzneimittelforschung</i> , 2004, 54, 444-451.	0.5	10
180	Prostaglandin E2 and I2 facilitate noxious heat-induced spike discharge but not iCGRP release from rat cutaneous nociceptors. <i>Life Sciences</i> , 2007, 81, 1685-1693.	2.0	10

#	ARTICLE	IF	CITATIONS
181	Projected pain from noxious heat stimulation of an exposed peripheral nerve – A case report. <i>European Journal of Pain</i> , 2009, 13, 35-37.	1.4	10
182	Mice and rats differ with respect to activity-dependent slowing of conduction velocity in the saphenous peripheral nerve. <i>Neuroscience Letters</i> , 2015, 592, 12-16.	1.0	10
183	The phospholipase C inhibitor U73122 is a potent agonist of the polymodal transient receptor potential ankyrin type 1 (TRPA1) receptor channel. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2020, 393, 177-189.	1.4	10
184	Can Receptor Potentials Be Detected With Threshold Tracking in Rat Cutaneous Nociceptive Terminals?. <i>Journal of Neurophysiology</i> , 2005, 94, 219-225.	0.9	9
185	Photosensitization of TRPA1 and TRPV1 by 7-dehydrocholesterol: implications for the Smith-Lemli-Opitz syndrome. <i>Pain</i> , 2017, 158, 2475-2486.	2.0	9
186	Painful diabetic neuropathy leads to functional CaV3.2 expression and spontaneous activity in skin nociceptors of mice. <i>Experimental Neurology</i> , 2021, 346, 113838.	2.0	9
187	The formalin test does not probe inflammatory pain but excitotoxicity in rodent skin. <i>Physiological Reports</i> , 2022, 10, e15194.	0.7	9
188	Inflammation and Hypersensitivity in the Context of the Sensory Functions of Axonal Membranes: What Are the Molecular Mechanisms?. <i>Digestive Diseases</i> , 2009, 27, 11-15.	0.8	8
189	Etomidate and propylene glycol activate nociceptive TRP ion channels. <i>Molecular Pain</i> , 2018, 14, 174480691881169.	1.0	8
190	Psoralens activate and photosensitize Transient Receptor Potential channels Ankyrin type 1 (TRPA1) and Vanilloid type 1 (TRPV1). <i>European Journal of Pain</i> , 2021, 25, 122-135.	1.4	8
191	Lysophosphatidic acid activates nociceptors and causes pain or itch depending on the application mode in human skin. <i>Pain</i> , 2022, 163, 445-460.	2.0	8
192	Nociceptive neurons in the rat caudal trigeminal nucleus respond to blood plasma perfusion of the subarachnoid space: the involvement of complement. <i>Pain</i> , 1999, 81, 283-288.	2.0	7
193	Afferent renal innervation in anti-Thy1.1 nephritis in rats. <i>American Journal of Physiology - Renal Physiology</i> , 2020, 319, F822-F832.	1.3	7
194	Effects of Bradykinin on Nociceptors. <i>NeuroImmune Biology</i> , 2009, 8, 135-168.	0.2	6
195	Taurolidine and congeners activate hTRPA1 but not hTRPV1 channels and stimulate CGRP release from mouse tracheal sensory nerves. <i>Pharmacology Research and Perspectives</i> , 2016, 4, e00204.	1.1	6
196	Norepinephrine reduces $\% \text{Ca}^{2+}$ -conotoxin-sensitive Ca^{2+} currents in renal afferent neurons in rats. <i>American Journal of Physiology - Renal Physiology</i> , 2012, 302, F350-F357.	1.3	5
197	The prokineticin Bv8 sensitizes cutaneous terminals of female mice to heat. <i>European Journal of Pain</i> , 2016, 20, 1326-1334.	1.4	5
198	Use dependence of peripheral nociceptive conduction in the absence of tetrodotoxin-resistant sodium channel subtypes. <i>Journal of Physiology</i> , 2016, 594, 5529-5541.	1.3	5

#	ARTICLE	IF	CITATIONS
199	Boole's algebra and the Gordian knot in peripheral nociception. Pain Forum, 1998, 7, 84-86.	1.1	4
200	Stimulated release of calcitonin gene-related peptide from the human right atrium in patients with and without diabetes mellitus. Peptides, 2006, 27, 3255-3260.	1.2	4
201	Reactive dicarbonyl compounds cause Calcitonin Gene-Related Peptide release and synergize with inflammatory conditions in mouse skin and peritoneum. Journal of Biological Chemistry, 2020, 295, 6330-6343.	1.6	4
202	TRPA1-mediated nociception: Response to letter by Fischer et al. Neuroscience, 2008, 155, 339.	1.1	3
203	Nociceptor Excitation by Sensitization: A Novel Hypothesis, Its Cellular and Molecular Background. , 2000, , 9-19.		3
204	Acetylsalicylic acid reduces pH-induced excitation of rat cutaneous nociceptors in vitro. Primary Sensory Neuron: the International Interdisciplinary Journal Reporting Basic and Clinical Research on Sensory Receptors and Primary Afferent Neurons, 1997, 2, 77-94.	0.3	2
205	Intradermally applied LPA but not bile salts induce itch and pain in humans depending on the mode of application. Journal of Hepatology, 2017, 66, S355.	1.8	2
206	Activity and connectivity changes of central projection areas revealed by functional magnetic resonance imaging in Nav1.8-deficient mice upon cold signaling. Scientific Reports, 2017, 7, 543.	1.6	1
207	Imaging the influence of peripheral TRPV1-signaling on cerebral nociceptive processing applying fMRI-based graph theory in a resiniferatoxin rat model. PLoS ONE, 2022, 17, e0266669.	1.1	1
208	Effects of ReN1869, a CNS-available antihistamine, on capsaicin- and histamine-induced neurogenic inflammation in healthy subjects. Drug Development Research, 2002, 57, 193-199.	1.4	0
209	MUDr. Ladislav Vyklický, DrSc. (1925-2008). European Journal of Pain, 2009, 13, 327-328.	1.4	0
210	T114 NAV1.9 IS A TRANSFORMATION AMPLIFIER REDUCING POLYMODAL RECEPTIVE THRESHOLDS. European Journal of Pain Supplements, 2011, 5, 19.	0.0	0
211	(368) Sodium channel Nav1.8 mediates axonal TTX-resistant conduction in peripheral C-fibres in mice. Journal of Pain, 2016, 17, S67.	0.7	0
212	The Role of TRP Channels in Nicotinic Provoked Pain and Irritation from the Oral Cavity and Throat: Translating Animal Data to Humans. Nicotine and Tobacco Research, 2022, , .	1.4	0