Gerardo Gamba

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/4492225/gerardo-gamba-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

98 papers

9,678 citations

46 h-index

98 g-index

102 ext. papers

10,618 ext. citations

7.9 avg, IF

5.99 L-index

#	Paper	IF	Citations
98	SIRT7 modulates the stability and activity of the renal K-Cl cotransporter KCC4 through deacetylation. <i>EMBO Reports</i> , 2021 , 22, e50766	6.5	3
97	Role of KLHL3 and dietary K in regulating KS-WNK1 expression. <i>American Journal of Physiology - Renal Physiology</i> , 2021 , 320, F734-F747	4.3	2
96	Molecular mechanisms for the modulation of blood pressure and potassium homeostasis by the distal convoluted tubule <i>EMBO Molecular Medicine</i> , 2021 , e14273	12	2
95	WNK3 and WNK4 exhibit opposite sensitivity with respect to cell volume and intracellular chloride concentration. <i>American Journal of Physiology - Cell Physiology</i> , 2020 , 319, C371-C380	5.4	7
94	Mutation affecting the conserved acidic WNK1 motif causes inherited hyperkalemic hyperchloremic acidosis. <i>Journal of Clinical Investigation</i> , 2020 , 130, 6379-6394	15.9	11
93	Familial Hyperkalemic Hypertension Genotype With a Negative Phenotype: A CUL3 Mosaicism. <i>American Journal of Hypertension</i> , 2020 , 33, 278-281	2.3	1
92	Structure-function relationships in the renal NaCl cotransporter (NCC). <i>Current Topics in Membranes</i> , 2019 , 83, 177-204	2.2	3
91	Regulation of the renal NaCl cotransporter by the WNK/SPAK pathway: lessons learned from genetically altered animals. <i>American Journal of Physiology - Renal Physiology</i> , 2019 , 316, F146-F158	4.3	9
90	Resilience to acute kidney injury in offspring of maternal protein restriction. <i>American Journal of Physiology - Renal Physiology</i> , 2019 , 317, F1637-F1648	4.3	3
89	Geraniin is a diuretic by inhibiting the Na-K-2Cl cotransporter NKCC2. <i>American Journal of Physiology - Renal Physiology</i> , 2018 , 314, F240-F250	4.3	5
88	Inactivation of SPAK kinase reduces body weight gain in mice fed a high-fat diet by improving energy expenditure and insulin sensitivity. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2018 , 314, E53-E65	6	9
87	Disruption of the with no lysine kinase-STE20-proline alanine-rich kinase pathway reduces the hypertension induced by angiotensin II. <i>Journal of Hypertension</i> , 2018 , 36, 361-367	1.9	1
86	Kidney-specific WNK1 isoform (KS-WNK1) is a potent activator of WNK4 and NCC. <i>American Journal of Physiology - Renal Physiology</i> , 2018 , 315, F734-F745	4.3	23
85	Letter to the editor: Remembering Steve Hebert (1946-2008). <i>American Journal of Physiology - Cell Physiology</i> , 2018 , 315, C122-C123	5.4	
84	The Calcium-Sensing Receptor Increases Activity of the Renal NCC through the WNK4-SPAK Pathway. <i>Journal of the American Society of Nephrology: JASN</i> , 2018 , 29, 1838-1848	12.7	15
83	Phosphorylation by PKC and PKA regulate the kinase activity and downstream signaling of WNK4. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017 , 114, E879-E886	11.5	28
82	In Reply to lassessing the Effect of Spironolactone on Acute Kidney Injury After Cardiac Surgery U American Journal of Kidney Diseases, 2017 , 70, 152-153	7.4	

(2015-2017)

81	Gender Differences in the Acute Kidney Injury to Chronic Kidney Disease Transition. <i>Scientific Reports</i> , 2017 , 7, 12270	4.9	28
8o	The thiazide sensitive sodium chloride co-transporter NCC is modulated by site-specific ubiquitylation. <i>Scientific Reports</i> , 2017 , 7, 12981	4.9	10
79	Mexneurin is a novel precursor of peptides in the central nervous system of rodents. <i>FEBS Letters</i> , 2017 , 591, 1627-1636	3.8	5
78	The Effect of Spironolactone on Acute Kidney Injury After Cardiac Surgery: A Randomized, Placebo-Controlled Trial. <i>American Journal of Kidney Diseases</i> , 2017 , 69, 192-199	7.4	18
77	With no lysine L-WNK1 isoforms are negative regulators of the K+-Cl- cotransporters. <i>American Journal of Physiology - Cell Physiology</i> , 2016 , 311, C54-66	5.4	9
76	The European Eel NCCIGene Encodes a Thiazide-resistant Na-Cl Cotransporter. <i>Journal of Biological Chemistry</i> , 2016 , 291, 22472-22481	5.4	4
75	On the molecular mechanism of renal salt excretion modulation by extracellular potassium. <i>Journal of Physiology</i> , 2016 , 594, 6071-6072	3.9	2
74	The regulation of Na+Cl- cotransporter by with-no-lysine kinase 4. <i>Current Opinion in Nephrology and Hypertension</i> , 2016 , 25, 417-23	3.5	7
73	SPAK and OSR1 play essential roles in potassium homeostasis through actions on the distal convoluted tubule. <i>Journal of Physiology</i> , 2016 , 594, 4945-66	3.9	36
7²	Mini-review: regulation of the renal NaCl cotransporter by hormones. <i>American Journal of Physiology - Renal Physiology</i> , 2016 , 310, F10-4	4.3	33
71	Regulation of Renal Electrolyte Transport by WNK and SPAK-OSR1 Kinases. <i>Annual Review of Physiology</i> , 2016 , 78, 367-89	23.1	123
7°	Gain-of-function missense variant in SLC12A2, encoding the bumetanide-sensitive NKCC1 cotransporter, identified in human schizophrenia. <i>Journal of Psychiatric Research</i> , 2016 , 77, 22-6	5.2	31
69	Unique chloride-sensing properties of WNK4 permit the distal nephron to modulate potassium homeostasis. <i>Kidney International</i> , 2016 , 89, 127-34	9.9	148
68	AT1 receptor antagonism before ischemia prevents the transition of acute kidney injury to chronic kidney disease. <i>Kidney International</i> , 2016 , 89, 363-73	9.9	62
67	Potassium and Its Discontents: New Insight, New Treatments. <i>Journal of the American Society of Nephrology: JASN</i> , 2016 , 27, 981-9	12.7	51
66	Physiological role of SLC12 family members in the kidney. <i>American Journal of Physiology - Renal Physiology</i> , 2016 , 311, F131-44	4.3	24
65	Insulin and SGK1 reduce the function of Na+/monocarboxylate transporter 1 (SMCT1/SLC5A8). <i>American Journal of Physiology - Cell Physiology</i> , 2016 , 311, C720-C734	5.4	6
64	Ovarian hormones and prolactin increase renal NaCl cotransporter phosphorylation. <i>American Journal of Physiology - Renal Physiology</i> , 2015 , 308, F799-808	4.3	32

63	The Effect of WNK4 on the Na+-Cl- Cotransporter Is Modulated by Intracellular Chloride. <i>Journal of the American Society of Nephrology: JASN</i> , 2015 , 26, 1781-6	12.7	102
62	Heat shock protein 72 (Hsp72) specific induction and temporal stability in urine samples as a reliable biomarker of acute kidney injury (AKI). <i>Biomarkers</i> , 2015 , 20, 453-9	2.6	15
61	Increased phosphorylation of the renal Na+-Cl- cotransporter in male kidney transplant recipient patients with hypertension: a prospective cohort. <i>American Journal of Physiology - Renal Physiology</i> , 2015 , 309, F836-42	4.3	18
60	Exonic mutations in the SLC12A3 gene cause exon skipping and premature termination in Gitelman syndrome. <i>Journal of the American Society of Nephrology: JASN</i> , 2015 , 26, 271-9	12.7	26
59	Revisiting the NaCl cotransporter regulation by with-no-lysine kinases. <i>American Journal of Physiology - Cell Physiology</i> , 2015 , 308, C779-91	5.4	38
58	Modulation of NCC activity by low and high K(+) intake: insights into the signaling pathways involved. <i>American Journal of Physiology - Renal Physiology</i> , 2014 , 306, F1507-19	4.3	73
57	WNK-SPAK-NCC cascade revisited: WNK1 stimulates the activity of the Na-Cl cotransporter via SPAK, an effect antagonized by WNK4. <i>Hypertension</i> , 2014 , 64, 1047-53	8.5	60
56	Intra-renal transfection of heat shock protein 90 alpha or beta (Hsp90lbr Hsp90llprotects against ischemia/reperfusion injury. <i>Nephrology Dialysis Transplantation</i> , 2014 , 29, 301-12	4.3	13
55	WNK3 abrogates the NEDD4-2-mediated inhibition of the renal Na+-Cl- cotransporter. <i>American Journal of Physiology - Renal Physiology</i> , 2014 , 307, F275-86	4.3	18
54	Hyperkalemic hypertension-associated cullin 3 promotes WNK signaling by degrading KLHL3. <i>Journal of Clinical Investigation</i> , 2014 , 124, 4723-36	15.9	87
53	Mineralocorticoid receptor phosphorylation regulates ligand binding and renal response to volume depletion and hyperkalemia. <i>Cell Metabolism</i> , 2013 , 18, 660-71	24.6	122
52	The SLC12 family of electroneutral cation-coupled chloride cotransporters. <i>Molecular Aspects of Medicine</i> , 2013 , 34, 288-98	16.7	93
51	Molecular evidence for a role for K(+)-Cl(-) cotransporters in the kidney. <i>American Journal of Physiology - Renal Physiology</i> , 2013 , 305, F1402-11	4.3	27
50	N-terminal serine dephosphorylation is required for KCC3 cotransporter full activation by cell swelling. <i>Journal of Biological Chemistry</i> , 2013 , 288, 31468-76	5.4	35
49	Spironolactone prevents chronic kidney disease caused by ischemic acute kidney injury. <i>Kidney International</i> , 2013 , 83, 93-103	9.9	81
48	Insulin increases the functional activity of the renal NaCl cotransporter. <i>Journal of Hypertension</i> , 2013 , 31, 303-11	1.9	54
47	Regulation of the renal Na+-Cl- cotransporter by phosphorylation and ubiquitylation. <i>American Journal of Physiology - Renal Physiology</i> , 2012 , 303, F1573-83	4.3	55
46	Recovery from ischemic acute kidney injury by spironolactone administration. <i>Nephrology Dialysis Transplantation</i> , 2012 , 27, 3160-9	4.3	46

(2007-2012)

45	Mechanisms of sodium-chloride cotransporter modulation by angiotensin II. <i>Current Opinion in Nephrology and Hypertension</i> , 2012 , 21, 516-22	3.5	29
44	Activation of the renal Na+:Cl- cotransporter by angiotensin II is a WNK4-dependent process. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012 , 109, 7929-34	11.5	191
43	WNK3-SPAK interaction is required for the modulation of NCC and other members of the SLC12 family. <i>Cellular Physiology and Biochemistry</i> , 2012 , 29, 291-302	3.9	29
42	WNK3 is a putative chloride-sensing kinase. <i>Cellular Physiology and Biochemistry</i> , 2011 , 28, 1123-34	3.9	32
41	Aldosterone paradox: differential regulation of ion transport in distal nephron. <i>Physiology</i> , 2011 , 26, 115-23	9.8	88
40	Hsp72 is an early and sensitive biomarker to detect acute kidney injury. <i>EMBO Molecular Medicine</i> , 2011 , 3, 5-20	12	49
39	Nedd4-2 modulates renal Na+-Cl- cotransporter via the aldosterone-SGK1-Nedd4-2 pathway. Journal of the American Society of Nephrology: JASN, 2011 , 22, 1707-19	12.7	127
38	Activation of the bumetanide-sensitive Na+,K+,2Cl- cotransporter (NKCC2) is facilitated by Tamm-Horsfall protein in a chloride-sensitive manner. <i>Journal of Biological Chemistry</i> , 2011 , 286, 30200-	·1 ⁵ 0 ⁴	119
37	WNK2 kinase is a novel regulator of essential neuronal cation-chloride cotransporters. <i>Journal of Biological Chemistry</i> , 2011 , 286, 30171-80	5.4	67
36	The nanopeptide hormone vasopressin is a new player in the modulation of renal Na(+)-Cl(-) cotransporter activity. <i>Kidney International</i> , 2010 , 78, 127-9	9.9	5
35	The thiazide-sensitive Na+-Cl- cotransporter: molecular biology, functional properties, and regulation by WNKs. <i>American Journal of Physiology - Renal Physiology</i> , 2009 , 297, F838-48	4.3	68
34	Angiotensin II signaling increases activity of the renal Na-Cl cotransporter through a WNK4-SPAK-dependent pathway. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009 , 106, 4384-9	11.5	200
33	Thick ascending limb: the Na(+):K (+):2Cl (-) co-transporter, NKCC2, and the calcium-sensing receptor, CaSR. <i>Pflugers Archiv European Journal of Physiology</i> , 2009 , 458, 61-76	4.6	101
32	A novel protein kinase signaling pathway essential for blood pressure regulation in humans. <i>Trends in Endocrinology and Metabolism</i> , 2008 , 19, 91-5	8.8	33
31	Regulation of NKCC2 by a chloride-sensing mechanism involving the WNK3 and SPAK kinases. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008 , 105, 8458-63	11.5	174
30	WNK kinases, renal ion transport and hypertension. <i>American Journal of Nephrology</i> , 2008 , 28, 860-70	4.6	40
29	Roles of the cation-chloride cotransporters in neurological disease. <i>Nature Clinical Practice Neurology</i> , 2008 , 4, 490-503		294
28	Identification and functional characterization of cation-chloride cotransporters in plants. <i>Plant Journal</i> , 2007 , 50, 278-92	6.9	164

27	NKCC2 surface expression in mammalian cells: down-regulation by novel interaction with aldolase B. <i>Journal of Biological Chemistry</i> , 2007 , 282, 33817-33830	5.4	28
26	Thiazide diuretics directly induce osteoblast differentiation and mineralized nodule formation by interacting with a sodium chloride co-transporter in bone. <i>Journal of the American Society of Nephrology: JASN</i> , 2007 , 18, 2509-16	12.7	82
25	Phorbol ester stimulation of RasGRP1 regulates the sodium-chloride cotransporter by a PKC-independent pathway. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007 , 104, 20120-5	11.5	34
24	Affinity-defining domains in the Na-Cl cotransporter: a different location for Cl- and thiazide binding. <i>Journal of Biological Chemistry</i> , 2006 , 281, 17266-17275	5.4	42
23	The Na+:Cl- cotransporter is activated and phosphorylated at the amino-terminal domain upon intracellular chloride depletion. <i>Journal of Biological Chemistry</i> , 2006 , 281, 28755-63	5.4	182
22	WNK3 bypasses the tonicity requirement for K-Cl cotransporter activation via a phosphatase-dependent pathway. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006 , 103, 1976-81	11.5	100
21	WNK protein kinases modulate cellular Cl- flux by altering the phosphorylation state of the Na-K-Cl and K-Cl cotransporters. <i>Physiology</i> , 2006 , 21, 326-35	9.8	98
20	Role of WNK kinases in regulating tubular salt and potassium transport and in the development of hypertension. <i>American Journal of Physiology - Renal Physiology</i> , 2005 , 288, F245-52	4.3	85
19	WNK3 kinase is a positive regulator of NKCC2 and NCC, renal cation-Cl- cotransporters required for normal blood pressure homeostasis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005 , 102, 16777-82	11.5	156
18	Molecular physiology and pathophysiology of electroneutral cation-chloride cotransporters. <i>Physiological Reviews</i> , 2005 , 85, 423-93	47.9	613
17	WNK3 modulates transport of Cl- in and out of cells: implications for control of cell volume and neuronal excitability. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005 , 102, 16783-8	11.5	178
16	A single nucleotide polymorphism alters the activity of the renal Na+:Cl- cotransporter and reveals a role for transmembrane segment 4 in chloride and thiazide affinity. <i>Journal of Biological Chemistry</i> , 2004 , 279, 16553-60	5.4	31
15	Electroneutral cation-chloride cotransporters in the central nervous system. <i>Neurochemical Research</i> , 2004 , 29, 17-25	4.6	61
14	Molecular physiology of cation-coupled Cl- cotransport: the SLC12 family. <i>Pflugers Archiv European Journal of Physiology</i> , 2004 , 447, 580-93	4.6	205
13	N-Glycosylation at two sites critically alters thiazide binding and activity of the rat thiazide-sensitive Na(+):Cl(-) cotransporter. <i>Journal of the American Society of Nephrology: JASN</i> , 2003 , 14, 271-82	12.7	107
12	Molecular pathogenesis of inherited hypertension with hyperkalemia: the Na-Cl cotransporter is inhibited by wild-type but not mutant WNK4. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003 , 100, 680-4	11.5	342
11	Functional properties of the apical Na+-K+-2Cl- cotransporter isoforms. <i>Journal of Biological Chemistry</i> , 2002 , 277, 11004-12	5.4	96
10	Molecular, functional, and genomic characterization of human KCC2, the neuronal K-Cl cotransporter. <i>Molecular Brain Research</i> , 2002 , 103, 91-105		98

LIST OF PUBLICATIONS

9	Renal potassium-chloride cotransporters. Current Opinion in Nephrology and Hypertension, 2001, 10, 685	-9. \$	34
8	Deranged transcriptional regulation of cell-volume-sensitive kinase hSGK in diabetic nephropathy. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000 , 97, 8157-62	11.5	194
7	Functional comparison of the K+-Cl- cotransporters KCC1 and KCC4. <i>Journal of Biological Chemistry</i> , 2000 , 275, 30326-34	5.4	138
6	Cloning and characterization of KCC3 and KCC4, new members of the cation-chloride cotransporter gene family. <i>Journal of Biological Chemistry</i> , 1999 , 274, 16355-62	5.4	229
5	Molecular biology of distal nephron sodium transport mechanisms. <i>Kidney International</i> , 1999 , 56, 1606	- 2 ,2 ₀	30
4	The many roles of the calcium-sensing receptor in health and disease. <i>Archives of Medical Research</i> , 1999 , 30, 436-48	6.6	28
3	Cloning and functional expression of a rat kidney extracellular calcium/polyvalent cation-sensing receptor. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995 , 92, 131-	. 5 11.5	421
2	Primary structure and functional expression of a cDNA encoding the thiazide-sensitive, electroneutral sodium-chloride cotransporter. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993 , 90, 2749-53	11.5	316
1	Cloning and characterization of an extracellular Ca(2+)-sensing receptor from bovine parathyroid. Nature 1993, 366, 575-80	50.4	2224