## Courtney D Sandau

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4496198/publications.pdf

Version: 2024-02-01

33 papers

1,945 citations

331538 21 h-index 33 g-index

35 all docs 35 does citations

35 times ranked 2244 citing authors

#	Article	IF	CITATIONS
1	Polychlorinated biphenyl (PCB) concentrations and profiles in marine mammals from the North Atlantic Ocean. Chemosphere, 2022, 288, 132639.	4.2	13
2	Development of retention time indices for comprehensive multidimensional gas chromatography and application to ignitable liquid residue mapping in wildfire investigations. Journal of Chromatography A, 2021, 1635, 461717.	1.8	18
3	The application of comprehensive two-dimensional gas chromatography to the analysis of wildfire debris for ignitable liquid residue. Forensic Science International, 2020, 310, 110256.	1.3	10
4	Evaluation of the effectiveness of different indicator PCBs to estimating total PCB concentrations in environmental investigations. Chemosphere, 2019, 237, 124429.	4.2	38
5	Background levels of dioxin-like polychlorinated biphenyls (dlPCBs), polychlorinated, polybrominated and mixed halogenated dibenzo-p-dioxins and dibenzofurans (PCDD/Fs, PBDD/Fs & PXDD/Fs) in sera of pregnant women in Accra, Ghana. Science of the Total Environment, 2019, 673, 631-642.	3.9	23
6	Soil gas investigation of an alleged gas migration issue on a residential farm located above the Weyburn-Midale CO2 enhanced oil recovery project. International Journal of Greenhouse Gas Control, 2019, 81, 11-20.	2.3	8
7	Forensic source attribution for toluene in environmental samples. Environmental Toxicology and Chemistry, 2018, 37, 729-737.	2.2	2
8	Comprehensive two-dimensional gas chromatography high-resolution mass spectrometry for the analysis of substituted and unsubstituted polycyclic aromatic compounds in environmental samples. Journal of Chromatography A, 2018, 1579, 106-114.	1.8	9
9	Developing deep high-resolution concentration and 13C isotope profiles for methane, ethane, and propane. Journal of Petroleum Science and Engineering, 2018, 170, 280-290.	2.1	7
10	Snapping Turtles (Chelydra serpentina) from Canadian Areas of Concern across the southern Laurentian Great Lakes: Chlorinated and brominated hydrocarbon contaminants and metabolites in relation to circulating concentrations of thyroxine and vitamin A. Environmental Research, 2015, 143, 266-278.	3.7	14
11	Elucidating the structural properties that influence the persistence of PCBs in humans using the National Health and Nutrition Examination Survey (NHANES) dataset. Science of the Total Environment, 2013, 461-462, 99-107.	3.9	35
12	Paraben Levels in an Urban Community of Western Canada. ISRN Toxicology, 2013, 2013, 1-8.	2.7	13
13	Data Comparability Between Biomonitoring Studies for PCDD/Fsâ€"Issues for the Use of the National Health and Nutrition Examination Survey (NHANES) Data. Epidemiology, 2011, 22, S33-S34.	1.2	O
14	Polyhalogenated aromatic hydrocarbons and metabolites: Relation to circulating thyroid hormone and retinol in nestling bald eagles ( <i>Haliaeetus leucocephalus</i> ). Environmental Toxicology and Chemistry, 2010, 29, 1301-1310.	2,2	22
15	Forensic Geo-Gas Investigation of Methane: Characterization of Sources within an Urban Setting. Environmental Forensics, 2010, 11, 108-116.	1.3	5
16	Thyroid Hormone Levels of Pregnant Inuit Women and Their Infants Exposed to Environmental Contaminants. Environmental Health Perspectives, 2009, 117, 1014-1020.	2.8	77
17	Concentration and profile of 22 urinary polycyclic aromatic hydrocarbon metabolites in the US population. Environmental Research, 2008, 107, 320-331.	3.7	266
18	Plasma Concentrations of Selected Organobromine Compounds and Polychlorinated Biphenyls in Postmenopausal Women of Québec, Canada. Environmental Health Perspectives, 2007, 115, 1429-1434.	2.8	71

#	Article	IF	CITATIONS
19	Levels of persistent organic pollutants (POPs) in a coastal northern Norwegian population with high fish-liver intake. Journal of Environmental Monitoring, 2006, 8, 552.	2.1	25
20	Automated solid-phase extraction method for measuring urinary polycyclic aromatic hydrocarbon metabolites in human biomonitoring using isotope-dilution gas chromatography high-resolution mass spectrometry. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2006, 835, 47-54.	1.2	96
21	Exposure to Polycyclic Aromatic Hydrocarbons in Children in the United States. Epidemiology, 2006, 17, S34.	1.2	0
22	An assessment of the toxicological significance of anthropogenic contaminants in Canadian arctic wildlife. Science of the Total Environment, 2005, 351-352, 57-93.	3.9	160
23	A reassessment of the nomenclature of polychlorinated biphenyl (PCB) metabolites Environmental Health Perspectives, 2004, 112, 291-294.	2.8	60
24	Development of a accelerated solvent extraction and gel permeation chromatography analytical method for measuring persistent organohalogen compounds in adipose and organ tissue analysis. Chemosphere, 2004, 57, 373-381.	4.2	104
25	New high-resolution mass spectrometric approach for the measurement of polychlorinated biphenyls and organochlorine pesticides in human serum. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2003, 794, 137-148.	1.2	104
26	Comprehensive Solid-Phase Extraction Method for Persistent Organic Pollutants. Validation and Application to the Analysis of Persistent Chlorinated Pesticides. Analytical Chemistry, 2003, 75, 71-77.	3.2	100
27	Pentachlorophenol and hydroxylated polychlorinated biphenyl metabolites in umbilical cord plasma of neonates from coastal populations in Québec Environmental Health Perspectives, 2002, 110, 411-417.	2.8	140
28	Methylsulfone polycglorinated biphenyl and 2,2â€bis(chlorophenyl)â€1,1â€dichloroethylene metabolites in beluga whale ( <i>Delphinapterus leucas</i> ) from the St. Lawrence river estuary and western Hudson bay, Canada. Environmental Toxicology and Chemistry, 2000, 19, 1378-1388.	2.2	12
29	Analysis of hydroxylated metabolites of PCBs (OH-PCBs) and other chlorinated phenolic compounds in whole blood from Canadian inuit Environmental Health Perspectives, 2000, 108, 611-616.	2.8	182
30	The Enantioselective Bioaccumulation of Chiral Chlordane and $\hat{l}_{\pm}$ -HCH Contaminants in the Polar Bear Food Chain. Environmental Science & Eamp; Technology, 2000, 34, 2668-2674.	4.6	130
31	Identification of 4-Hydroxyheptachlorostyrene in Polar Bear Plasma and Its Binding Affinity to Transthyretin:  A Metabolite of Octachlorostyrene?. Environmental Science &	4.6	74
32	METHYLSULFONE POLYCHLORINATED BIPHENYL AND 2,2-BIS(CHLOROPHENYL)-1,1-DICHLOROETHYLENE METABOLITES IN BELUGA WHALE (DELPHINAPTERUS LEUCAS) FROM THE ST. LAWRENCE RIVER ESTUARY AND WESTERN HUDSON BAY, CANADA. Environmental Toxicology and Chemistry, 2000, 19, 1378.	2.2	25
33	Enantioselective Gas Chromatography/Mass Spectrometry of Methylsulfonyl PCBs with Application to Arctic Marine Mammals. Analytical Chemistry, 1998, 70, 3845-3852.	3.2	53