Guido Lattuada

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4497437/publications.pdf

Version: 2024-02-01

45 2,599 24 45 papers citations h-index g-index

46 46 46 3993
all docs docs citations times ranked citing authors

#	Article	IF	CITATIONS
1	Prevalence, Metabolic Features, and Prognosis of Metabolically Healthy Obese Italian Individuals. Diabetes Care, 2011, 34, 210-215.	4.3	335
2	Habitual Physical Activity Is Associated With Intrahepatic Fat Content in Humans. Diabetes Care, 2007, 30, 683-688.	4.3	273
3	Effects of metabolic modulation by trimetazidine on left ventricular function and phosphocreatine/adenosine triphosphate ratio in patients with heart failure. European Heart Journal, 2006, 27, 942-948.	1.0	210
4	Fatty liver index and mortality: The cremona study in the 15th year of follow-up. Hepatology, 2011, 54, 145-152.	3 . 6	208
5	Increased mediastinal fat and impaired left ventricular energy metabolism in young men with newly found fatty liver. Hepatology, 2008, 47, 51-58.	3.6	182
6	Insulin resistance, intramyocellular lipid content, and plasma adiponectin in patients with type 1 diabetes. American Journal of Physiology - Endocrinology and Metabolism, 2003, 285, E1174-E1181.	1.8	150
7	Fasting Plasma Leptin, Tumor Necrosis Factor-Â Receptor 2, and Monocyte Chemoattracting Protein 1 Concentration in a Population of Glucose-Tolerant and Glucose-Intolerant Women: Impact on cardiovascular mortality. Diabetes Care, 2003, 26, 2883-2889.	4.3	117
8	Insulin resistance and whole body energy homeostasis in obese adolescents with fatty liver disease. American Journal of Physiology - Endocrinology and Metabolism, 2006, 291, E697-E703.	1.8	105
9	Insulin resistance/hyperinsulinemia and cancer mortality: the Cremona study at the 15th year of follow-up. Acta Diabetologica, 2012, 49, 421-428.	1.2	89
10	Association Between Plasma Monocyte Chemoattractant Protein-1 Concentration and Cardiovascular Disease Mortality in Middle-Aged Diabetic and Nondiabetic Individuals. Diabetes Care, 2009, 32, 2105-2110.	4.3	80
11	Screening for non-alcoholic fatty liver disease in type 2 diabetes using non-invasive scores and association with diabetic complications. BMJ Open Diabetes Research and Care, 2020, 8, e000904.	1.2	71
12	Reduced intrahepatic fat content is associated with increased whole-body lipid oxidation in patients with type 1 diabetes. Diabetologia, 2005, 48, 2615-2621.	2.9	65
13	Effect of partial inhibition of fatty acid oxidation by trimetazidine on whole body energy metabolism in patients with chronic heart failure. Heart, 2011, 97, 1495-1500.	1.2	60
14	Abnormal Left Ventricular Energy Metabolism in Obese Men With Preserved Systolic and Diastolic Functions Is Associated With Insulin Resistance. Diabetes Care, 2007, 30, 1520-1526.	4.3	59
15	Effect of the sporting discipline on the right and left ventricular morphology and function of elite male track runners: A magnetic resonance imaging and phosphorus 31 spectroscopy study. American Heart Journal, 2007, 154, 937-942.	1.2	56
16	NAFLD/NASH in patients with type 2 diabetes and related treatment options. Journal of Endocrinological Investigation, 2018, 41, 509-521.	1.8	50
17	Serum Retinol-Binding Protein-4, Leptin, and Adiponectin Concentrations Are Related to Ectopic Fat Accumulation. Journal of Clinical Endocrinology and Metabolism, 2007, 92, 4883-4888.	1.8	49
18	Serum Resistin and Hepatic Fat Content in Nondiabetic Individuals. Journal of Clinical Endocrinology and Metabolism, 2006, 91, 5122-5125.	1.8	43

#	Article	IF	CITATIONS
19	Impact of diabetes on COVID-19-related in-hospital mortality: a retrospective study from Northern Italy. Journal of Endocrinological Investigation, 2021, 44, 843-850.	1.8	41
20	Reduced whole-body lipid oxidation is associated with insulin resistance, but not with intramyocellular lipid content in offspring of type 2 diabetic patients. Diabetologia, 2005, 48, 741-747.	2.9	37
21	Increased serum resistin in elite endurance athletes with high insulin sensitivity. Diabetologia, 2006, 49, 1893-1900.	2.9	34
22	Postabsorptive and insulin-stimulated energy and protein metabolism in patients with myotonic dystrophy type 1. American Journal of Clinical Nutrition, 2004, 80, 357-364.	2.2	30
23	Beneficial effects of betaâ€blockers on left ventricular function and cellular energy reserve in patients with heart failure. Fundamental and Clinical Pharmacology, 2013, 27, 455-464.	1.0	27
24	Why Does NAFLD Predict Type 2 Diabetes?. Current Diabetes Reports, 2011, 11, 167-172.	1.7	26
25	Left ventricular function and energy metabolism in middle-aged men undergoing long-lasting sustained aerobic oxidative training. Heart, 2008, 95, 630-635.	1.2	19
26	Postabsorptive and Insulin-Stimulated Energy Homeostasis and Leucine Turnover in Offspring of Type 2 Diabetic Patients. Diabetes Care, 2004, 27, 2716-2722.	4.3	18
27	Lack of association of apoE Îμ4 allele with insulin resistance. Acta Diabetologica, 2012, 49, 25-32.	1.2	18
28	Assessment of insulin sensitivity based on a fasting blood sample in men with liver cirrhosis before and after liver transplantation 1. Transplantation, 2003, 76, 697-702.	0.5	15
29	Free leptin index and thyroid function in male highly trained athletes. European Journal of Endocrinology, 2009, 161, 871-876.	1.9	15
30	Lipid accumulation in overweight type 2 diabetic subjects: relationships with insulin sensitivity and adipokines. Acta Diabetologica, 2013, 50, 301-307.	1.2	15
31	Effects of short-term manipulation of serum FFA concentrations on left ventricular energy metabolism and function in patients with heart failure: no association with circulating bio-markers of inflammation. Acta Diabetologica, 2015, 52, 753-761.	1.2	14
32	Impact of using different biomarkers of liver fibrosis on hepatologic referral of individuals with severe obesity and NAFLD. Journal of Endocrinological Investigation, 2020, 43, 1019-1026.	1.8	13
33	Resting cardiac energy metabolism is inversely associated with heart rate in healthy young adult men. American Heart Journal, 2011, 162, 136-141.	1.2	12
34	Resting Energy Expenditure in Obese Women with Primary Hypothyroidism and Appropriate Levothyroxine Replacement Therapy. Journal of Clinical Endocrinology and Metabolism, 2020, 105, e1741-e1748.	1.8	12
35	The anti-ischemic effect of trimetazidine in patients with postprandial myocardial ischemia is unrelated to meal composition. American Heart Journal, 2006, 151, 1238.e1-1238.e8.	1.2	8
36	Elevated fasting plasma Câ€peptide occurs in nonâ€diabetic individuals with fatty liver, irrespective of insulin resistance. Diabetic Medicine, 2009, 26, 847-854.	1.2	7

#	Article	IF	CITATIONS
37	Baseline TSH levels and short-term weight loss after different procedures of bariatric surgery. International Journal of Obesity, 2021, 45, 326-330.	1.6	7
38	Increased low-grade inflammation is associated with lack of functional response to carvedilol in patients with systolic heart failure. Journal of Cardiovascular Medicine, 2013, 14, 49-56.	0.6	6
39	Hypercortisolism and altered glucose homeostasis in obese patients in the preâ€bariatric surgery assessment. Diabetes/Metabolism Research and Reviews, 2021, 37, e3389.	1.7	5
40	A high carbohydrate meal yields a lower ischemic threshold than a high fat meal in patients with stable coronary disease. International Journal of Cardiology, 2011, 147, 209-213.	0.8	3
41	Left ventricular function and energy homeostasis in patients with type 1 diabetes with and without microvascular complications. International Journal of Cardiology, 2012, 154, 111-115.	0.8	3
42	Fasting Whole-Body Energy Homeostasis and Hepatic Energy Metabolism in Nondiabetic Humans with Fatty Liver. Oxidative Medicine and Cellular Longevity, 2019, 2019, 1-7.	1.9	3
43	Resting Whole Body Energy Metabolism in Class 3 Obesity; from Preserved Insulin Sensitivity to Overt Type 2 Diabetes. Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy, 2020, Volume 13, 489-497.	1.1	3
44	Effects of endurance exercise training on metabolic and inflammatory parameters in HIV-1-infected patients with secondary lipodystrophy and diabetes. Sport Sciences for Health, 2010, 6, 23-25.	0.4	1
45	Metabolic and Psychological Features are Associated with Weight Loss 12 Months After Sleeve Gastrectomy. Journal of Clinical Endocrinology and Metabolism, 2021, 106, e3087-e3097.	1.8	1