

S Bruce Archibald

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4500921/publications.pdf>

Version: 2024-02-01

36
papers

908
citations

623734

14
h-index

454955

30
g-index

36
all docs

36
docs citations

36
times ranked

794
citing authors

#	ARTICLE	IF	CITATIONS
1	The Palaeorehniidae (Orthoptera, Ensifera, <i>Zeuneropterinae</i>), and new taxa from the early Eocene Okanagan Highlands, western North America. <i>Zootaxa</i> , 2022, 5100, 559-572.	0.5	1
2	<i>Kishenehna prima</i> , a new genus and species of darner dragonfly (Odonata, Aeshnidae). <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 707 Td (Go</i> 5099, 496-500.	0.5	0
3	<i>Danowhetaksa</i> gen. nov. with two species from the early Eocene <i>Å</i> lst Formation from Denmark, the first Palearctic <i>Whetwhetaksidae</i> (Odonata: Cephalozygoptera). <i>Zootaxa</i> , 2022, 5099, 586-592.	0.5	3
4	A revision of Chrysopidae (Neuroptera) from the late Eocene Florissant Formation, Colorado, with description of new species. <i>Zootaxa</i> , 2022, 5133, 301-345.	0.5	2
5	The first Odonata from the early Eocene Allenby Formation of the Okanagan Highlands, British Columbia, Canada (Anisoptera, Aeshnidae and <i>cf</i>). <i>Cephalozygoptera, Dysagrionidae</i> . <i>Canadian Entomologist</i> , 2022, 154, .	0.8	3
6	The Cephalozygoptera, a new, extinct suborder of Odonata with new taxa from the early Eocene Okanagan Highlands, western North America. <i>Zootaxa</i> , 2021, 4934, <i>zootaxa.4934.1.1</i> .	0.5	11
7	<i>New Protosmylinae</i> (Neuroptera: Osmylidae) from the early Eocene of western North America, with taxonomic remarks. <i>Zootaxa</i> , 2021, 4980, 142-156.	0.5	0
8	The second North American fossil horntail wood-wasp (Hymenoptera: Siricidae), from the early Eocene Green River Formation. <i>Zootaxa</i> , 2021, 4999, 325-334.	0.5	2
9	The head of Cephalozygoptera (Odonata). <i>Zootaxa</i> , 2021, 5047, 97-100.	0.5	4
10	A new genus and species of split-footed lacewings (Neuroptera) from the early Eocene of western Canada and revision of the subfamily affinities of Mesozoic Nymphidae. <i>Canadian Entomologist</i> , 2020, 152, 269-287.	0.8	4
11	Fossil dragonflies (Odonata: Anisoptera) from the early Eocene Okanagan Highlands, western North America. <i>Canadian Entomologist</i> , 2019, 151, 783-816.	0.8	8
12	The oldest Inocelliidae (Raphidioptera) from the Eocene of western North America. <i>Canadian Entomologist</i> , 2019, 151, 521-530.	0.8	5
13	Early Eocene (Ypresian) birds from the Okanagan Highlands, British Columbia (Canada) and Washington State (USA). <i>Canadian Journal of Earth Sciences</i> , 2019, 56, 803-813.	1.3	6
14	Two new species of fossil <i>Eomerope</i> (Mecoptera: Eomeropidae) from the Ypresian Okanagan Highlands, far-western North America, and Eocene Holarctic dispersal of the genus. <i>Canadian Entomologist</i> , 2018, 150, 393-403.	0.8	10
15	New early Eocene Siricomorpha (Hymenoptera: Symphyta: Pamphiliidae, Siricidae, Cephidae) from the Okanagan Highlands, western North America. <i>Canadian Entomologist</i> , 2016, 148, 209-228.	0.8	13
16	A new species of <i>Archaeochrysa</i> Adams (Neuroptera: Chrysopidae) from the early Eocene of Driftwood Canyon, British Columbia, Canada. <i>Canadian Entomologist</i> , 2015, 147, 359-369.	0.8	12
17	Stem-group stick insects (Phasmatodea) in the early Eocene at McAbee, British Columbia, Canada, and Republic, Washington, United States of America. <i>Canadian Entomologist</i> , 2015, 147, 744-753.	0.8	21
18	The second genus and species of the extinct neuropteroid family <i>Corydasialidae</i> , from early Eocene McAbee, British Columbia, Canada. <i>Zootaxa</i> , 2015, 4040, 569.	0.5	5

#	ARTICLE	IF	CITATIONS
19	A revision of the late Eocene snakeflies (Raphidioptera) of the Florissant Formation, Colorado, with special reference to the wing venation of the Raphidiomorpha. <i>Zootaxa</i> , 2014, 3784, 401-44.	0.5	13
20	Early Eocene big headed flies (Diptera: Pipunculidae) from the Okanagan Highlands, western North America. <i>Canadian Entomologist</i> , 2014, 146, 429-443.	0.8	4
21	Fossil palm beetles refine upland winter temperatures in the Early Eocene Climatic Optimum. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 8095-8100.	7.1	48
22	The Eocene apex of panorpoid scorpionfly family diversity. <i>Journal of Paleontology</i> , 2013, 87, 677-695.	0.8	26
23	A diverse new assemblage of green lacewings (Insecta, Neuroptera, Chrysopidae) from the early Eocene Okanagan Highlands, western North America. <i>Journal of Paleontology</i> , 2013, 87, 123-146.	0.8	26
24	New species of Nymphites Haase (Neuroptera: Nymphidae) from the Middle Jurassic of China, with a redescription of the type species of the genus. <i>Zootaxa</i> , 2013, 3700, 393.	0.5	9
25	Intercontinental dispersal of giant thermophilic ants across the Arctic during early Eocene hyperthermals. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2011, 278, 3679-3686.	2.6	63
26	Seasonality, the latitudinal gradient of diversity, and Eocene insects. <i>Paleobiology</i> , 2010, 36, 374-398.	2.0	127
27	New Cimbrophlebiidae (Insecta: Mecoptera) from the Early Eocene at McAbee, British Columbia, Canada and Republic, Washington, USA. <i>Zootaxa</i> , 2009, 2249, 51-62.	0.5	24
28	New fossil species of Nymphidae (Neuroptera) from the Eocene of North America and Europe. <i>Zootaxa</i> , 2009, 2157, 59-68.	0.5	13
29	Bulldog Ants of the Eocene Okanagan Highlands and History of the Subfamily (Hymenoptera:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 10 1	2.5	57
30	Ecology and Distribution of Cenozoic Eomeropidae (Mecoptera), and a New Species of <I>Eomerope</I> Cockerell from the Early Eocene McAbee Locality, British Columbia, Canada. <i>Annals of the Entomological Society of America</i> , 2005, 98, 503-514.	2.5	28
31	Fossil biotas from the Okanagan Highlands, southern British Columbia and northeastern Washington State: climates and ecosystems across an Eocene landscape. <i>Canadian Journal of Earth Sciences</i> , 2005, 42, 167-185.	1.3	179
32	Regional and local vegetation community dynamics of the Eocene Okanagan Highlands (British) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 2	1.3	79
33	New genus of minute Berothidae (Neuroptera) from Early Eocene amber of British Columbia. <i>Canadian Entomologist</i> , 2004, 136, 61-76.	0.8	21
34	New Early Eocene brown lacewings (Neuroptera: Hemerobiidae) from western North America. <i>Canadian Entomologist</i> , 2003, 135, 637-653.	0.8	19
35	An Early Eocene bee (Hymenoptera: Halictidae) from Quilchena, British Columbia. <i>Canadian Entomologist</i> , 2003, 135, 63-69.	0.8	14
36	Early Eocene insects from Quilchena, British Columbia, and their paleoclimatic implications. <i>Canadian Journal of Zoology</i> , 2000, 78, 1441-1462.	1.0	48