Yoshio Saito

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4502286/publications.pdf Version: 2024-02-01



Υσεμίο δλιτο

#	Article	IF	CITATIONS
1	FRET-based dual-labeled oligonucleotide probes detect target DNA by probing a minor groove environment. Organic and Biomolecular Chemistry, 2022, 20, 1699-1706.	2.8	2
2	2-Ethynylnaphthalene-modified 8-aza-3,7-dideaza-2′-deoxyadenosine derivative discriminates thymine in target DNA via changes in the fluorescence wavelength. Tetrahedron Letters, 2020, 61, 151841.	1.4	2
3	Synthesis of ethynylpyrene-modified 3-deaza-2′-deoxyguanosines as environmentally sensitive fluorescent nucleosides: Target DNA-sequence detection via changes in the fluorescence wavelength. Tetrahedron Letters, 2019, 60, 825-830.	1.4	3
4	A fluorescent 3,7-bis-(naphthalen-1-ylethynylated)-2′-deoxyadenosine analogue reports thymidine in complementary DNA by a large emission Stokes shift. Organic and Biomolecular Chemistry, 2018, 16, 1496-1507.	2.8	11
5	Base-modified fluorescent purine nucleosides and nucleotides for use in oligonucleotide probes. Journal of Photochemistry and Photobiology C: Photochemistry Reviews, 2018, 36, 48-73.	11.6	42
6	The fluorescently responsive 3-(naphthalen-1-ylethynyl)-3-deaza-2′-deoxyguanosine discriminates cytidine via the DNA minor groove. Organic and Biomolecular Chemistry, 2017, 15, 7853-7859.	2.8	11
7	C2-substituted 8-aza-7-deaza-2′-deoxyadenosines as environmentally sensitive fluorescent nucleosides for discriminating apurinic/apyrimidinic sites in DNA duplex. Tetrahedron Letters, 2017, 58, 117-120.	1.4	11
8	Fluorescent Purine Nucleosides and Their Applications. Nucleic Acids and Molecular Biology, 2016, , 27-61.	0.2	0
9	Design and synthesis of a novel fluorescent benzo[g]imidazo[4,5-c]quinoline nucleoside for monitoring base-pair-induced protonation with cytosine: distinguishing cytosine via changes in the intensity and wavelength of fluorescence. Organic and Biomolecular Chemistry, 2016, 14, 3934-3942.	2.8	13
10	Design and synthesis of environmentally sensitive fluorescent 2-naphthylethynylated 2′-deoxyadenosines: Detection of target DNA via changes in fluorescence intensity and wavelength. Bioorganic and Medicinal Chemistry Letters, 2016, 26, 684-689.	2.2	7
11	Design and synthesis of 7-naphthyl-8-aza-7-deaza-2′-deoxyadenosines as environmentally sensitive fluorescent nucleosides. Tetrahedron Letters, 2015, 56, 3034-3038.	1.4	16
12	Synthesis of 8-aza-3,7-dideaza-2′-deoxyadenosines possessing a new adenosine skeleton as an environmentally sensitive fluorescent nucleoside for monitoring the DNA minor groove. Organic and Biomolecular Chemistry, 2015, 13, 7459-7468.	2.8	21
13	Synthesis and photophysical properties of pyrene-labeled 3-deaza-2′-deoxyadenosines comprising a non-ï€-conjugated linker: fluorescence quenching-based oligodeoxynucleotide probes for thymine identification. Organic and Biomolecular Chemistry, 2015, 13, 10540-10547.	2.8	9
14	Molecular Design of an Environmentally Sensitive Fluorescent Nucleoside, 3â€Deazaâ€2â€2â€Deoxyadenosine Derivative: Distinguishing Thymine by Probing the DNA Minor Groove. ChemBioChem, 2014, 15, 1638-1644.	2.6	24
15	Design of an environmentally sensitive fluorescent 8-aza-7-deaza-2′-deoxyadenosine derivative with dual fluorescence for the specific detection of thymine. Organic and Biomolecular Chemistry, 2014, 12, 660-666.	2.8	40
16	Naphthalene-based environmentally sensitive fluorescent 8-substituted 2′-deoxyadenosines: Application to DNA detection. Bioorganic and Medicinal Chemistry Letters, 2013, 23, 886-892.	2.2	27
17	Synthesis of solvatofluorochromic 7-arylethynylated 7-deaza-2′-deoxyadenosine derivatives: application to the design of environmentally sensitive fluorescent probes forming stable DNA duplexes. Tetrahedron Letters, 2013, 54, 2348-2352.	1.4	17
18	An environmentally sensitive fluorescent purine nucleoside that changes emission wavelength upon hybridization. Chemical Communications, 2013, 49, 5684.	4.1	38

Υοςηίο Saito

#	Article	IF	CITATIONS
19	Fluorescent nucleosides with â€~on–off' switching function, pH-responsive fluorescent uridine derivatives. Bioorganic and Medicinal Chemistry Letters, 2012, 22, 2753-2756.	2.2	10
20	Fluorometric detection of adenine in target DNA by exciplex formation with fluorescent 8-arylethynylated deoxyguanosine. Bioorganic and Medicinal Chemistry Letters, 2012, 22, 3723-3726.	2.2	17
21	Synthesis and photophysical properties of novel push–pull-type solvatochromic 7-deaza-2′-deoxypurine nucleosides. Tetrahedron Letters, 2011, 52, 4726-4729.	1.4	22
22	Stabilization of DNA duplex by 2-substituted adenine as a minor groove modifier. Bioorganic and Medicinal Chemistry Letters, 2011, 21, 7021-7024.	2.2	7
23	Design and synthesis of highly solvatochromic fluorescent 2′-deoxyguanosine and 2′-deoxyadenosine analogs. Bioorganic and Medicinal Chemistry Letters, 2011, 21, 1275-1278.	2.2	38
24	Synthesis and photophysical properties of 8-arylbutadienyl 2′-deoxyguanosines. Tetrahedron Letters, 2011, 52, 491-494.	1.4	16
25	Synthesis of environmentally sensitive 2′-deoxyguanosine containing solvatochromic pyrene fluorophore. Tetrahedron Letters, 2011, 52, 2359-2361.	1.4	21
26	Synthesis of novel push–pull-type solvatochromic 2′-deoxyguanosine derivatives with longer wavelength emission. Tetrahedron Letters, 2010, 51, 2606-2609.	1.4	34
27	Design of environmentally sensitive fluorescent 2′-deoxyguanosine containing arylethynyl moieties: Distinction of thymine base by base-discriminating fluorescent (BDF) probe. Bioorganic and Medicinal Chemistry Letters, 2010, 20, 2817-2820.	2.2	31
28	Singly and doubly labeled base-discriminating fluorescent oligonucleotide probes containing oxo-pyrene chromophore. Bioorganic and Medicinal Chemistry Letters, 2010, 20, 3227-3230.	2.2	30
29	Photo-switching of vinylpyrene-substituted 2'-deoxyguanosine and its application. Nucleic Acids Symposium Series, 2009, 53, 193-194.	0.3	0
30	Fluorescence switching of photochromic vinylpyrene-substituted 2′-deoxyguanosine. Tetrahedron Letters, 2009, 50, 1403-1406.	1.4	18
31	Ends free and self-quenched molecular beacon with pyrene labeled pyrrolocytidine in the middle of the stem. Tetrahedron, 2009, 65, 934-939.	1.9	33
32	Pyrene-labeled deoxyguanosine as a fluorescence sensor to discriminate single and double stranded DNA structures: Design of ends free molecular beacons. Bioorganic and Medicinal Chemistry Letters, 2009, 19, 6392-6395.	2.2	40
33	Anthracene based base-discriminating fluorescent oligonucleotide probes for SNPs typing: Synthesis and photophysical properties. Bioorganic and Medicinal Chemistry, 2008, 16, 107-113.	3.0	38
34	C8-alkynyl- and alkylamino substituted 2′-deoxyguanosines: a universal linker for nucleic acids modification. Tetrahedron, 2008, 64, 3578-3588.	1.9	31
35	Fluorometric sensing of conformational switching of DNA; The use of fluorescence labeled C8-alkylamino substituted 2'-deoxyguanosine. Nucleic Acids Symposium Series, 2008, 52, 357-358.	0.3	1
36	Design of an ultimate quencher free molecular beacon containing pyrrolocytidine-guanine base pair. Nucleic Acids Symposium Series, 2008, 52, 361-362.	0.3	3

Υοςηίο Saito

#	Article	IF	CITATIONS
37	Design of an efficient self-quenched molecular beacon for SNPs genotyping. Nucleic Acids Symposium Series, 2008, 52, 359-360.	0.3	1
38	Design of dual-labeled oligonucleotide probes for SNPs genotyping. Nucleic Acids Symposium Series, 2007, 51, 23-24.	0.3	0
39	Synthesis of C8-alkylamino substituted 2'-deoxyguanosine. Nucleic Acids Symposium Series, 2007, 51, 147-148.	0.3	0
40	Design of a novel G-quenched molecular beacon: A simple and efficient strategy for DNA sequence analysis. Chemical Communications, 2007, , 4492.	4.1	26
41	Dual-labeled oligonucleotide probe for sensing adenosine via FRET: A novel alternative to SNPs genotyping. Chemical Communications, 2007, , 2133.	4.1	18
42	Acridone-labeled Base-discriminating Fluorescence (BDF) Nucleoside: Synthesis and Their Photophysical Properties. Chemistry Letters, 2006, 35, 1182-1183.	1.3	14
43	Highly selective fluorescent nucleobases for designing base-discriminating fluorescent probes. Pure and Applied Chemistry, 2006, 78, 2305-2312.	1.9	9
44	Synthesis and properties of acridone-labeled base-discriminating fluorescent (BDF) nucleosides. Nucleic Acids Symposium Series, 2006, 50, 181-182.	0.3	1
45	Design of base-discriminating fluorescent nucleosides. Journal of Photochemistry and Photobiology C: Photochemistry Reviews, 2005, 6, 108-122.	11.6	181
46	Synthesis and properties of purine-type base-discriminating fluorescent (BDF) nucleosides: distinction of thymine by fluorescence-labeled deoxyadenosine derivatives. Tetrahedron Letters, 2005, 46, 7605-7608.	1.4	30
47	Synthesis and properties of novel base-discriminating fluorescent (BDF) nucleosides. Nucleic Acids Symposium Series, 2005, 49, 153-154.	0.3	3
48	Design of base-discriminating fluorescent (BDF) nucleobase for SNP typing. Nucleic Acids Symposium Series, 2004, 48, 243-244.	0.3	2
49	Synthesis and properties of novel base-discriminating fluorescent (BDF) nucleosides: a highly polarity-sensitive fluorophore for SNP typing. Tetrahedron Letters, 2004, 45, 7827-7831.	1.4	32
50	Base-discriminating fluorescent (BDF) nucleoside: distinction of thymine by fluorescence quenchingElectronic supplementary information (ESI) available: Experimental procedures for new compounds, UV and excitation spectra. See http://www.rsc.org/suppdata/cc/b4/b405832a/. Chemical Communications, 2004, , 1704.	4.1	59