

# Bingbing Wang

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4502955/publications.pdf>

Version: 2024-02-01

17  
papers

626  
citations

687363

13  
h-index

888059

17  
g-index

17  
all docs

17  
docs citations

17  
times ranked

820  
citing authors

#	ARTICLE	IF	CITATIONS
1	Recapitulation of Short RNA-Directed Translational Gene Silencing In Vitro. <i>Molecular Cell</i> , 2006, 22, 553-560.	9.7	166
2	Distinct passenger strand and mRNA cleavage activities of human Argonaute proteins. <i>Nature Structural and Molecular Biology</i> , 2009, 16, 1259-1266.	8.2	121
3	MicroRNA-repressed mRNAs contain 40S but not 60S components. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 5343-5348.	7.1	94
4	RelB/p52-mediated NF- $\kappa$ B signaling alters histone acetylation to increase the abundance of corticotropin-releasing hormone in human placenta. <i>Science Signaling</i> , 2015, 8, ra85.	3.6	36
5	RelB/NF- $\kappa$ B2 Regulates Corticotropin-Releasing Hormone in the Human Placenta. <i>Molecular Endocrinology</i> , 2012, 26, 1356-1369.	3.7	34
6	Glucocorticoid Receptor Signaling Contributes to Constitutive Activation of the Noncanonical NF- $\kappa$ B Pathway in Term Human Placenta. <i>Molecular Endocrinology</i> , 2013, 27, 203-211.	3.7	22
7	Integrated RNA-seq and ChIP-seq analysis reveals a feed-forward loop regulating H3K9ac and key labor drivers in human placenta. <i>Placenta</i> , 2019, 76, 40-50.	1.5	21
8	Mono-(2-Ethylhexyl) Phthalate Promotes Pro-Labor Gene Expression in the Human Placenta. <i>PLoS ONE</i> , 2016, 11, e0147013.	2.5	21
9	Analysis of microRNA effector functions in vitro. <i>Methods</i> , 2007, 43, 91-104.	3.8	20
10	Anandamide down-regulates placental transporter expression through CB2 receptor-mediated inhibition of cAMP synthesis. <i>Pharmacological Research</i> , 2019, 141, 331-342.	7.1	17
11	miR-15b-5p promotes expression of proinflammatory cytokines in human placenta by inhibiting Apelin signaling pathway. <i>Placenta</i> , 2021, 104, 8-15.	1.5	17
12	Negative Effects of Progesterone Receptor Isoform-A on Human Placental Activity of the Noncanonical NF- $\kappa$ B Signaling. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014, 99, E320-E328.	3.6	16
13	Vitamin D stimulates multiple microRNAs to inhibit CRH and other pro-labor genes in human placenta. <i>Endocrine Connections</i> , 2018, 7, 1380-1388.	1.9	16
14	Fetal lung C4BPA induces p100 processing in human placenta. <i>Scientific Reports</i> , 2019, 9, 5519.	3.3	12
15	Screening a small molecule library to identify inhibitors of NF- $\kappa$ B inducing kinase and pro-labor genes in human placenta. <i>Scientific Reports</i> , 2018, 8, 1657.	3.3	6
16	Fetal Exosomal Platelet-activating Factor Triggers Functional Progesterone Withdrawal in Human Placenta. <i>Reproductive Sciences</i> , 2021, 28, 252-262.	2.5	5
17	Comparative Immunohistochemistry of Placental Corticotropin-Releasing Hormone and the Transcription Factor RelB-NF- $\kappa$ B2 Between Humans and Nonhuman Primates. <i>Comparative Medicine</i> , 2015, 65, 140-3.	1.0	2