

Shou-Wei Li

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4506867/publications.pdf>

Version: 2024-02-01

15
papers

691
citations

1305906

8
h-index

1255698

13
g-index

15
all docs

15
docs citations

15
times ranked

1298
citing authors

#	ARTICLE	IF	CITATIONS
1	Development and Validation of a Nomogram Model Based on Hematological Indicators for Predicting the Prognosis of Diffused Gliomas. <i>Frontiers in Surgery</i> , 2022, 9, 803237.	0.6	2
2	Alterations in the RTK/Ras/PI3K/AKT pathway serve as potential biomarkers for immunotherapy outcome of diffuse gliomas. <i>Aging</i> , 2021, 13, 15444-15458.	1.4	10
3	MET overexpression contributes to STAT4-PD-L1 signaling activation associated with tumor-associated, macrophages-mediated immunosuppression in primary glioblastomas. , 2021, 9, e002451.		13
4	Neutrophil depletion enhances the therapeutic effect of PD-1 antibody on glioma. <i>Aging</i> , 2020, 12, 15290-15301.	1.4	11
5	Intratumoral Susceptibility Signals Reflect Biomarker Status in Gliomas. <i>Scientific Reports</i> , 2019, 9, 17080.	1.6	15
6	Sanbo Scoring System, Based on Age and Pre-treatment Hematological Markers, is a Non-invasive and Independent Prognostic Predictor for Patients with Primary Glioblastomas: A Retrospective Multicenter Study. <i>Journal of Cancer</i> , 2019, 10, 5654-5660.	1.2	4
7	Phosphorylated Hsp27 is mutually exclusive with ATRX loss and the IDH1^{R132H} mutation and may predict better prognosis among glioblastomas without the IDH1 mutation and ATRX loss. <i>Journal of Clinical Pathology</i> , 2018, 71, 702-707.	1.0	16
8	Prognostic role of pretreatment circulating MDSCs in patients with solid malignancies: A meta-analysis of 40 studies. <i>Oncolimmunology</i> , 2018, 7, e1494113.	2.1	83
9	Higher Plasma Fibrinogen Levels are Associated with Malignant Phenotype and Worse Survival in Patients with Glioblastomas. <i>Journal of Cancer</i> , 2018, 9, 2024-2029.	1.2	6
10	Molecular and clinical characterization of PTPN2 expression from RNA-seq data of 996 brain gliomas. <i>Journal of Neuroinflammation</i> , 2018, 15, 145.	3.1	15
11	Preoperative Changes in Hematological Markers and Predictors of Glioma Grade and Survival. <i>Frontiers in Pharmacology</i> , 2018, 9, 886.	1.6	52
12	Allergy reduces the risk of meningioma: a meta-analysis. <i>Scientific Reports</i> , 2017, 7, 40333.	1.6	6
13	Immune-Related Adverse Events Associated with Anti-PD-1/PD-L1 Treatment for Malignancies: A Meta-Analysis. <i>Frontiers in Pharmacology</i> , 2017, 8, 730.	1.6	350
14	Preoperative inflammation markers and IDH mutation status predict glioblastoma patient survival. <i>Oncotarget</i> , 2017, 8, 50117-50123.	0.8	80
15	IDH-1R132H mutation status in diffuse glioma patients: implications for classification. <i>Oncotarget</i> , 2016, 7, 31393-31400.	0.8	28