

Naoko P Kato

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4519840/naoko-p-kato-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

53
papers

1,312
citations

19
h-index

34
g-index

66
ext. papers

1,554
ext. citations

2.5
avg, IF

3.81
L-index

#	Paper	IF	Citations
53	Methodological quality of studies assessing validity and reliability of the European Heart Failure Self-care Behaviour Scale: a systematic review using the COSMIN methodology. <i>European Journal of Cardiovascular Nursing</i> , 2021 , 20, 501-512	3.3	1
52	The consequences of the COVID-19 pandemic for self-care in patients supported with a left ventricular assist device. <i>European Journal of Heart Failure</i> , 2020 , 22, 933-936	12.3	10
51	Trajectory of self-care behaviour in patients with heart failure: the impact on clinical outcomes and influencing factors. <i>European Journal of Cardiovascular Nursing</i> , 2020 , 19, 421-432	3.3	7
50	Psychometric Testing of the Hebrew Version of the European Heart Failure Self-Care Behaviour Scale. <i>Heart Lung and Circulation</i> , 2020 , 29, e121-e130	1.8	1
49	Development of an Instrument for Measuring Self-Care Behaviors After Left Ventricular Assist Device Implantation. <i>Progress in Transplantation</i> , 2019 , 29, 335-343	1.1	3
48	NUP98-HBO1-fusion generates phenotypically and genetically relevant chronic myelomonocytic leukemia pathogenesis. <i>Blood Advances</i> , 2019 , 3, 1047-1060	7.8	4
47	A systematic review of heart failure dyadic self-care interventions focusing on intervention components, contexts, and outcomes. <i>International Journal of Nursing Studies</i> , 2018 , 77, 232-242	5.8	42
46	Quality of life of family caregivers of patients with a left ventricular assist device in Japan. <i>Journal of Cardiology</i> , 2018 , 71, 81-87	3	12
45	Novel working hypothesis for pathogenesis of hematological malignancies: combination of mutations-induced cellular phenotypes determines the disease (cMIP-DD). <i>Journal of Biochemistry</i> , 2016 , 159, 17-25	3.1	4
44	High Dose EBlocker Therapy Triggers Additional Reverse Remodeling in Patients With Idiopathic Non-Ischemic Cardiomyopathy. <i>International Heart Journal</i> , 2016 , 57, 717-724	1.8	11
43	How effective is an in-hospital heart failure self-care program in a Japanese setting? Lessons from a randomized controlled pilot study. <i>Patient Preference and Adherence</i> , 2016 , 10, 171-81	2.4	17
42	Development and Validation of a Heart Failure-Specific Health Literacy Scale. <i>Journal of Cardiovascular Nursing</i> , 2016 , 31, 131-9	2.1	19
41	Health literacy is independently associated with self-care behavior in patients with heart failure. <i>Patient Education and Counseling</i> , 2016 , 99, 1026-32	3.1	52
40	Quality of Life and Influential Factors in Patients Implanted With a Left Ventricular Assist Device. <i>Circulation Journal</i> , 2015 , 79, 2186-92	2.9	18
39	Parasympathetic reinnervation accompanied by improved post-exercise heart rate recovery and quality of life in heart transplant recipients. <i>International Heart Journal</i> , 2015 , 56, 180-5	1.8	17
38	Heart Failure Telemonitoring in Japan and Sweden: A Cross-Sectional Survey. <i>Journal of Medical Internet Research</i> , 2015 , 17, e258	7.6	12
37	Learning self-care after left ventricular assist device implantation. <i>Current Heart Failure Reports</i> , 2014 , 11, 290-8	2.8	53

36	Late-onset right ventricular failure in patients with preoperative small left ventricle after implantation of continuous flow left ventricular assist device. <i>Circulation Journal</i> , 2014 , 78, 625-33	2.9	41
35	Urine sodium excretion after tolvaptan administration is dependent upon baseline serum sodium levels: a possible explanation for the improvement of hyponatremia with scarce chance of hypernatremia by a vasopressin receptor antagonist. <i>International Heart Journal</i> , 2014 , 55, 131-7	1.8	18
34	Differential impacts of achieved heart rate and achieved dose of β blocker on clinical outcomes in heart failure with and without atrial fibrillation. <i>International Journal of Cardiology</i> , 2014 , 173, 331-3	3.2	9
33	Comparison of self-care behaviors of heart failure patients in 15 countries worldwide. <i>Patient Education and Counseling</i> , 2013 , 92, 114-20	3.1	153
32	Acute pulmonary vasoreactivity test with sildenafil or nitric monoxide before left ventricular assist device implantation. <i>Journal of Artificial Organs</i> , 2013 , 16, 389-92	1.8	8
31	A Randomized Controlled Pilot Study of the Japanese Heart Failure Self-Management Program: Impacts on Heart Failure Knowledge and Clinical Outcomes. <i>Journal of Cardiac Failure</i> , 2013 , 19, S91	3.3	2
30	Successful conversion from thiazide to tolvaptan in a patient with stage d heart failure and chronic kidney disease before heart transplantation. <i>International Heart Journal</i> , 2013 , 54, 48-50	1.8	11
29	Bosentan improved persistent pulmonary hypertension in a case after implantation of a left ventricular assist device. <i>Journal of Artificial Organs</i> , 2013 , 16, 101-4	1.8	13
28	Novel criteria of urine osmolality effectively predict response to tolvaptan in decompensated heart failure patients--association between non-responders and chronic kidney disease. <i>Circulation Journal</i> , 2013 , 77, 397-404	2.9	93
27	Trend of clinical outcome and surrogate markers during titration of β blocker in heart failure patients with reduced ejection fraction: relevance of achieved heart rate and β blocker dose. <i>Circulation Journal</i> , 2013 , 77, 1001-8	2.9	18
26	Urine osmolality estimated using urine urea nitrogen, sodium and creatinine can effectively predict response to tolvaptan in decompensated heart failure patients. <i>Circulation Journal</i> , 2013 , 77, 1208-13	2.9	34
25	Preoperative levels of bilirubin or creatinine adjusted by age can predict their reversibility after implantation of left ventricular assist device. <i>Circulation Journal</i> , 2013 , 77, 96-104	2.9	37
24	Insufficient self-care is an independent risk factor for adverse clinical outcomes in Japanese patients with heart failure. <i>International Heart Journal</i> , 2013 , 54, 382-9	1.8	34
23	Development and psychometric properties of the Japanese heart failure knowledge scale. <i>International Heart Journal</i> , 2013 , 54, 228-33	1.8	14
22	A case with recovery of response to tolvaptan associated with remission of acute kidney injury and increased urine osmolality. <i>International Heart Journal</i> , 2013 , 54, 115-8	1.8	8
21	Successful treatment of hemodynamic compromise caused by antibody-mediated and cellular rejection in a recipient 12 years after heart transplantation. <i>International Heart Journal</i> , 2013 , 54, 328-31	1.8	6
20	Development of self-care educational material for patients with heart failure in Japan: a pilot study. <i>Australian Journal of Cancer Nursing</i> , 2012 , 14, 156-64	1.9	7
19	Novel risk scoring system with preoperative objective parameters gives a good prediction of 1-year mortality in patients with a left ventricular assist device. <i>Circulation Journal</i> , 2012 , 76, 1895-903	2.9	36

18	Combination evaluation of preoperative risk indices predicts requirement of biventricular assist device. <i>Circulation Journal</i> , 2012 , 76, 2785-91	2.9	33
17	How to demonstrate the reversibility of end-organ function before implantation of left ventricular assist device in INTERMACS profile 2 patients?. <i>Journal of Artificial Organs</i> , 2012 , 15, 395-8	1.8	10
16	An elevated ratio of early to late diastolic filling velocity recovers after heart transplantation in a time-dependent manner. <i>Journal of Cardiology</i> , 2012 , 60, 295-300	3	8
15	Depressive symptoms are common and associated with adverse clinical outcomes in heart failure with reduced and preserved ejection fraction. <i>Journal of Cardiology</i> , 2012 , 60, 23-30	3	42
14	Correction of hyponatremia by tolvaptan before left ventricular assist device implantation. <i>International Heart Journal</i> , 2012 , 53, 391-3	1.8	10
13	Successful conversion to everolimus after cytomegalovirus infection in a heart transplant recipient. <i>International Heart Journal</i> , 2012 , 53, 199-201	1.8	9
12	Knowledge and Demand for Information about Islet Transplantation in Patients with Type 1 Diabetes. <i>Journal of Transplantation</i> , 2011 , 2011, 136298	2.3	3
11	Two types of C/EBP β mutations play distinct but collaborative roles in leukemogenesis: lessons from clinical data and BMT models. <i>Blood</i> , 2011 , 117, 221-33	2.2	49
10	Age and preoperative total bilirubin level can stratify prognosis after extracorporeal pulsatile left ventricular assist device implantation. <i>Circulation Journal</i> , 2011 , 75, 121-8	2.9	28
9	Less frequent opening of the aortic valve and a continuous flow pump are risk factors for postoperative onset of aortic insufficiency in patients with a left ventricular assist device. <i>Circulation Journal</i> , 2011 , 75, 1147-55	2.9	80
8	Quality of life as an independent predictor for cardiac events and death in patients with heart failure. <i>Circulation Journal</i> , 2011 , 75, 1661-9	2.9	61
7	Translation and validation study of the Japanese versions of the Coronary Revascularisation Outcome Questionnaire (CROQ-J). <i>European Journal of Cardiovascular Nursing</i> , 2011 , 10, 22-30	3.3	7
6	Validity and reliability of Seattle angina questionnaire Japanese version in patients with coronary artery disease. <i>Asian Nursing Research</i> , 2010 , 4, 57-63	2.5	4
5	Preventive advice given by patients with type 2 diabetes to their offspring. <i>British Journal of General Practice</i> , 2009 , 59, 37-42	1.6	9
4	Possible involvement of RasGRP4 in leukemogenesis. <i>International Journal of Hematology</i> , 2009 , 89, 470-481	2.5	17
3	Relationship of depressive symptoms with hospitalization and death in Japanese patients with heart failure. <i>Journal of Cardiac Failure</i> , 2009 , 15, 912-9	3.3	34
2	Adherence to self-care behavior and factors related to this behavior among patients with heart failure in Japan. <i>Heart and Lung: Journal of Acute and Critical Care</i> , 2009 , 38, 398-409	2.6	48
1	Validity and reliability of the Japanese version of the European Heart Failure Self-Care Behavior Scale. <i>European Journal of Cardiovascular Nursing</i> , 2008 , 7, 284-9	3.3	33

