

Yuichi Sato

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4521790/publications.pdf>

Version: 2024-02-01

81
papers

1,914
citations

236612

25
h-index

276539

41
g-index

84
all docs

84
docs citations

84
times ranked

2529
citing authors

#	ARTICLE	IF	CITATIONS
1	Expression of bcl-2 oncogene protein is prevalent in small cell lung carcinomas. <i>Journal of Pathology</i> , 1995, 177, 135-138.	2.1	167
2	Detection of Epstein-Barr Virus DNA in Reed-Sternberg Cells of Hodgkin's Disease Using the Polymerase Chain Reaction and in situ Hybridization. <i>Japanese Journal of Cancer Research</i> , 1990, 81, 272-278.	1.7	112
3	Comparison of the DNA Extraction Methods for Polymerase Chain Reaction Amplification from Formalin-Fixed and Paraffin-Embedded Tissues. <i>Diagnostic Molecular Pathology</i> , 2001, 10, 265-271.	2.1	102
4	hASH1 expression is closely correlated with endocrine phenotype and differentiation extent in pulmonary neuroendocrine tumors. <i>Modern Pathology</i> , 2004, 17, 222-229.	2.9	99
5	A new marker, insulinoma-associated protein 1 (INSM1), for high-grade neuroendocrine carcinoma of the uterine cervix: Analysis of 37 cases. <i>Gynecologic Oncology</i> , 2017, 144, 384-390.	0.6	77
6	Differential expression of N-cadherin and E-cadherin in normal human tissues. <i>Archives of Histology and Cytology</i> , 2006, 69, 135-145.	0.2	75
7	Expression of RACK1 is a novel biomarker in pulmonary adenocarcinomas. <i>Lung Cancer</i> , 2010, 69, 54-59.	0.9	61
8	Detection of Epstein-Barr Virus DNA in Formalin-Fixed Paraffin-Embedded Tissue of Nasopharyngeal Carcinoma Using Polymerase Chain Reaction and In Situ Hybridization. <i>Laryngoscope</i> , 1991, 101, 279-283.	1.1	53
9	Detection of tumor-specific autoantibodies in sera of patients with lung cancer. <i>Lung Cancer</i> , 2008, 62, 364-373.	0.9	46
10	Incidence of latent infection of Epstein-Barr virus in lung cancers—an analysis of EBEB1 expression in lung cancers by in situ hybridization. <i>Journal of Pathology</i> , 1994, 174, 257-265.	2.1	45
11	Prognostic Significance of Nestin Expression in Resected Non-small Cell Lung Cancer. <i>Chest</i> , 2011, 139, 862-869.	0.4	45
12	Diagnostic and Prognostic Significances of MUC5B and TTF-1 Expressions in Resected Non-Small Cell Lung Cancer. <i>Scientific Reports</i> , 2015, 5, 8649.	1.6	45
13	Prognostic Significance of MYH9 Expression in Resected Non-Small Cell Lung Cancer. <i>PLoS ONE</i> , 2015, 10, e0121460.	1.1	43
14	Proteomics of tumor-specific proteins in cerebrospinal fluid of patients with astrocytoma: Usefulness of gelsolin protein. <i>Pathology International</i> , 2009, 59, 797-803.	0.6	41
15	Clinicopathological features and the impact of the new TNM classification of malignant tumors in patients with pulmonary large cell neuroendocrine carcinoma. <i>Molecular and Clinical Oncology</i> , 2013, 1, 437-443.	0.4	39
16	Calnexin is a novel sero-diagnostic marker for lung cancer. <i>Lung Cancer</i> , 2015, 90, 342-345.	0.9	37
17	CAXII Is a Sero-Diagnostic Marker for Lung Cancer. <i>PLoS ONE</i> , 2012, 7, e33952.	1.1	37
18	Prognostic significance of G6PD expression and localization in lung adenocarcinoma. <i>Biochimica Et Biophysica Acta - Proteins and Proteomics</i> , 2019, 1867, 38-46.	1.1	34

#	ARTICLE	IF	CITATIONS
19	Occlusal disharmony attenuates glucocorticoid negative feedback in aged SAMP8 mice. <i>Neuroscience Letters</i> , 2007, 427, 71-76.	1.0	31
20	Cytoskeleton-Associated Protein 4 Is a Novel Serodiagnostic Marker for Lung Cancer. <i>American Journal of Pathology</i> , 2018, 188, 1328-1333.	1.9	31
21	Significant high expression of cytokeratins 7, 8, 18, 19 in pulmonary large cell neuroendocrine carcinomas, compared to small cell lung carcinomas. <i>Pathology International</i> , 2010, 60, 71-77.	0.6	28
22	S100A16 is a Prognostic Marker for Lung Adenocarcinomas. <i>Asian Pacific Journal of Cancer Prevention</i> , 2015, 16, 7039-7044.	0.5	28
23	Human cytomegalovirus infection in foci of Langerhans cell histiocytosis. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 1999, 434, 109-115.	1.4	27
24	Factors Contributing to Early Recovery of Urinary Continence Analyzed by Pre- and Postoperative Pelvic Anatomical Features at Robot-Assisted Laparoscopic Radical Prostatectomy. <i>Journal of Endourology</i> , 2015, 29, 683-690.	1.1	27
25	Angiogenic factors in normal endometrium and endometrial adenocarcinoma. <i>Pathology International</i> , 2007, 57, 140-147.	0.6	26
26	Proteomic study of sera from patients with bladder cancer: usefulness of S100A8 and S100A9 proteins. <i>Cancer Genomics and Proteomics</i> , 2010, 7, 181-9.	1.0	26
27	The clinicopathological significance of Lgr5 expression in lung adenocarcinoma. <i>Lung Cancer</i> , 2013, 82, 143-148.	0.9	24
28	Maternal chewing during prenatal stress ameliorates stress-induced hypomyelination, synaptic alterations, and learning impairment in mouse offspring. <i>Brain Research</i> , 2016, 1651, 36-43.	1.1	24
29	The balance between the expressions of hASH1 and HES1 differs between large cell neuroendocrine carcinoma and small cell carcinoma of the lung. <i>Lung Cancer</i> , 2011, 74, 405-410.	0.9	23
30	Clinicopathological Significance of S100A10 Expression in Lung Adenocarcinomas. <i>Asian Pacific Journal of Cancer Prevention</i> , 2016, 17, 289-294.	0.5	23
31	S100A16, a promising candidate as a prognostic marker for platinum-based adjuvant chemotherapy in resected lung adenocarcinoma. <i>OncoTargets and Therapy</i> , 2017, Volume 10, 5273-5279.	1.0	22
32	Loss of periplakin expression is associated with pathological stage and cancerspecific survival in patients with urothelial carcinoma of the urinary bladder. <i>Biomedical Research</i> , 2014, 35, 201-206.	0.3	21
33	Detection of tumor-associated antigens in culture supernatants using autoantibodies in sera from patients with bladder cancer. <i>Biomedical Research</i> , 2014, 35, 25-35.	0.3	20
34	Moesin and Stress-Induced Phosphoprotein-1 Are Possible Sero-Diagnostic Markers of Psoriasis. <i>PLoS ONE</i> , 2014, 9, e101773.	1.1	19
35	Prognostic significance of NAP1L1 expression in patients with early lung adenocarcinoma. <i>Biomedical Research</i> , 2020, 41, 149-159.	0.3	17
36	Prevalent and up-regulated vimentin expression in micropapillary components of lung adenocarcinomas and its adverse prognostic significance. <i>Pathology International</i> , 2015, 65, 183-192.	0.6	16

#	ARTICLE	IF	CITATIONS
37	Immunohistochemical examination of tumor-suppressor gene p53 product and pyrimidine dimer in solar keratosis. <i>Journal of Cancer Research and Clinical Oncology</i> , 1993, 119, 260-262.	1.2	15
38	Prognostic impact of nestin expression in resected large cell neuroendocrine carcinoma of the lung. <i>Lung Cancer</i> , 2012, 77, 415-420.	0.9	15
39	Nestin Is an Independent Predictor of Cancer-Specific Survival after Radical Cystectomy in Patients with Urothelial Carcinoma of the Bladder. <i>PLoS ONE</i> , 2014, 9, e91548.	1.1	15
40	pH-sensitive expression of calcium-sensing receptor (CaSR) in type-B intercalated cells of the cortical collecting ducts (CCD) in mouse kidney. <i>Clinical and Experimental Nephrology</i> , 2015, 19, 771-782.	0.7	15
41	Co-expression of N-cadherin and alpha-fetoprotein in stomach cancer. <i>Pathology International</i> , 2001, 51, 612-618.	0.6	14
42	A New Possible Lung Cancer Marker: VGF Detection from the Conditioned Medium of Pulmonary Large Cell Neuroendocrine Carcinoma-Derived Cells using Secretome Analysis. <i>International Journal of Biological Markers</i> , 2009, 24, 282-285.	0.7	14
43	Prognostic significance of S100A16 subcellular localization in lung adenocarcinoma. <i>Human Pathology</i> , 2018, 74, 148-155.	1.1	14
44	Erythropoietin production by the kidney and the liver in response to severe hypoxia evaluated by Western blotting with deglycosylation. <i>Physiological Reports</i> , 2020, 8, e14485.	0.7	14
45	Clinicopathological and prognostic significance of nuclear UGDH localization in lung adenocarcinoma. <i>Biomedical Research</i> , 2019, 40, 17-27.	0.3	12
46	Histiocytosis X: Successful Treatment with Recombinant Interferon- α . <i>Pediatrics International</i> , 1990, 32, 151-154.	0.2	11
47	Fludrocortisone stimulates erythropoietin production in the intercalated cells of the collecting ducts. <i>Biochemical and Biophysical Research Communications</i> , 2018, 503, 3121-3127.	1.0	11
48	Prognostic significance of nuclear DJ-1 expression in astrocytoma. <i>Anticancer Research</i> , 2010, 30, 265-9.	0.5	11
49	Upregulation of calbindin D28k in the late distal tubules in the potassium-loaded adrenalectomized mouse kidney. <i>Clinical and Experimental Nephrology</i> , 2011, 15, 355-362.	0.7	9
50	Impact of EGFR-Tyrosine Kinase Inhibitors on Postoperative Recurrent Non-Small-Cell Lung Cancer Harboring EGFR Mutations. <i>Oncology Research and Treatment</i> , 2017, 40, 7-13.	0.8	9
51	Late Somatic Gene 2 disrupts parental spheroids cooperatively with Volvox hatching enzyme A in Volvox. <i>Planta</i> , 2017, 245, 183-192.	1.6	9
52	Prognostic significance of nestin expression in patients with resected non-small cell lung cancer treated with platinum-based adjuvant chemotherapy; relationship between nestin expression and epithelial to mesenchymal transition related markers. <i>PLoS ONE</i> , 2017, 12, e0173886.	1.1	9
53	Serum Anti-Gal-3 Autoantibody is a Predictive Marker of the Efficacy of Platinum-Based Chemotherapy against Pulmonary Adenocarcinoma. <i>Asian Pacific Journal of Cancer Prevention</i> , 2015, 16, 7959-7965.	0.5	9
54	Acquisition of useful sero-diagnostic autoantibodies using the same patients' sera and tumor tissues. <i>Biomedical Research</i> , 2014, 35, 133-143.	0.3	8

#	ARTICLE	IF	CITATIONS
55	Novel stress increases hypothalamic-pituitary-adrenal activity in mice with a raised bite. <i>Archives of Oral Biology</i> , 2016, 68, 55-60.	0.8	8
56	Basigin expression as a prognostic indicator in stage I pulmonary adenocarcinoma. <i>Pathology International</i> , 2018, 68, 232-240.	0.6	8
57	Effects of Angiotensin II on Erythropoietin Production in the Kidney and Liver. <i>Molecules</i> , 2021, 26, 5399.	1.7	8
58	Peroxiredoxin 1 expression in active ulcerative colitis mucosa identified by proteome analysis and involvement of thioredoxin based on immunohistochemistry. <i>Oncology Letters</i> , 2017, 15, 2364-2372.	0.8	6
59	Prognostic significance of IMMT expression in surgically resected lung adenocarcinoma. <i>Thoracic Cancer</i> , 2019, 10, 2142-2151.	0.8	6
60	TRAP1 is a predictive biomarker of platinum-based adjuvant chemotherapy benefits in patients with resected lung adenocarcinoma. <i>Biomedical Research</i> , 2020, 41, 53-65.	0.3	6
61	PD-L1 expression in tumor-infiltrating lymphocytes (TILs) as an independent predictor of prognosis in patients with pN0 bladder cancer undergoing radical cystectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2021, 39, 195.e15-195.e23.	0.8	6
62	Prognostic Impact of AHNK2 Expression in Patients Treated with Radical Cystectomy. <i>Cancers</i> , 2021, 13, 1748.	1.7	6
63	Analysis of Autoantibodies Related to Tumor Progression in Sera from Patients with High-grade Non-muscle-invasive Bladder Cancer. <i>Anticancer Research</i> , 2017, 37, 6705-6714.	0.5	6
64	Differentiation of endogenous erythropoietin and exogenous ESAs by Western blotting. <i>Heliyon</i> , 2020, 6, e05389.	1.4	5
65	High HNRNPA3 expression is associated with lymph node metastasis and poor prognosis in patients treated with radical cystectomy. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2021, 39, 196.e1-196.e7.	0.8	5
66	Prognostic significance of galectin-3 expression in patients with resected NSCLC treated with platinum-based adjuvant chemotherapy. <i>Thoracic Cancer</i> , 2021, 12, 1570-1578.	0.8	5
67	Immunohistochemical and Semiquantitative Immunoblot Analyses of Nm23-H1 and H2 Isoforms in Normal Human Tissues.. <i>Acta Histochemica Et Cytochemica</i> , 1998, 31, 411-418.	0.8	4
68	Qualitative and Quantitative Assessment of Modified In Situ Reverse Transcription-Polymerase Chain Reaction (In Situ RT-PCR) Method.. <i>Acta Histochemica Et Cytochemica</i> , 1999, 32, 423-430.	0.8	4
69	Semiquantitative immunoblot analysis of nm23-H1 and -H2 isoforms in adenocarcinomas of the lung: Prognostic significance. <i>Pathology International</i> , 2000, 50, 200-205.	0.6	4
70	Effects of Roxadustat on Erythropoietin Production in the Rat Body. <i>Molecules</i> , 2022, 27, 1119.	1.7	4
71	Vulnerability to stress in mouse offspring is ameliorated when pregnant dams are provided a chewing stick during prenatal stress. <i>Archives of Oral Biology</i> , 2019, 97, 150-155.	0.8	3
72	TROY expression is associated with pathological stage and poor prognosis in patients treated with radical cystectomy. <i>Cancer Biomarkers</i> , 2019, 24, 91-96.	0.8	3

#	ARTICLE	IF	CITATIONS
73	Serum Epiplakin Might Be a Potential Serodiagnostic Biomarker for Bladder Cancer. <i>Cancers</i> , 2021, 13, 5150.	1.7	3
74	Expression of Membranous CD155 Is Associated with Aggressive Phenotypes and a Poor Prognosis in Patients with Bladder Cancer. <i>Cancers</i> , 2022, 14, 1576.	1.7	3
75	Exploring disease-specific methylated CpGs in human male genital abnormalities by using methylated-site display-amplified fragment length polymorphism (MSD-AFLP). <i>Journal of Reproduction and Development</i> , 2019, 65, 491-497.	0.5	2
76	DJ-1 Expression Might Serve as a Biologic Marker in Patients with Bladder Cancer. <i>Cancers</i> , 2022, 14, 2535.	1.7	2
77	Phase I study of Nedaplatin, a platinum based antineoplastic drug, combined with nab-paclitaxel in patients with advanced squamous non-small cell lung cancer. <i>Investigational New Drugs</i> , 2018, 36, 45-52.	1.2	1
78	Proteome analysis using the cancer stem cell-like micropapillary adenocarcinoma cell line. <i>Denki Eido</i> , 2017, 61, 39-44.	0.0	1
79	Relationship between the Laterality of Retinopathy and the Development of Ischemic Stroke in Diabetes Mellitus. <i>The Journal of Japan Atherosclerosis Society</i> , 1995, 22, 735-739.	0.0	0
80	Search of sero-diagnostic marker using the Reverse-phase protein array. <i>Denki Eido</i> , 2017, 61, 23-28.	0.0	0
81	Application of markers obtained by various proteomic methods to resected non-small cell lung cancer treated with platinum-based adjuvant chemotherapy. <i>Denki Eido</i> , 2017, 61, 128-131.	0.0	0