

# JÃ¼rgen Machann

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4530176/publications.pdf>

Version: 2024-02-01

166  
papers

10,296  
citations

28274

55  
h-index

38395

95  
g-index

172  
all docs

172  
docs citations

172  
times ranked

14062  
citing authors

#	ARTICLE	IF	CITATIONS
1	Editorial for "Concentration of Gallbladder Phosphatidylcholine in Cholangiopathies: A <sup>31</sup> P MR Spectroscopy Pilot Study". Journal of Magnetic Resonance Imaging, 2022, 55, 541-542.	3.4	0
2	A comparison of emulsifiers for the formation of oil-in-water emulsions: stability of the emulsions within 9h after production and MR signal properties. Magnetic Resonance Materials in Physics, Biology, and Medicine, 2022, 35, 401-410.	2.0	5
3	Metabolic implications of pancreatic fat accumulation. Nature Reviews Endocrinology, 2022, 18, 43-54.	9.6	46
4	Empagliflozin Improves Insulin Sensitivity of the Hypothalamus in Humans With Prediabetes: A Randomized, Double-Blind, Placebo-Controlled, Phase 2 Trial. Diabetes Care, 2022, 45, 398-406.	8.6	43
5	Short-Term Variability of Proton Density Fat Fraction in Pancreas and Liver Assessed by Multiecho Chemical-Shift Encoding-Based <sup>1</sup> H MRI at 3T. Journal of Magnetic Resonance Imaging, 2022, 56, 1018-1026.	3.4	8
6	Population-based cohort imaging: skeletal muscle mass by magnetic resonance imaging in correlation to bioelectrical impedance analysis. Journal of Cachexia, Sarcopenia and Muscle, 2022, 13, 976-986.	7.3	8
7	Impaired Metabolic Health and Low Cardiorespiratory Fitness Independently Associate With Subclinical Atherosclerosis in Obesity. Journal of Clinical Endocrinology and Metabolism, 2022, , .	3.6	3
8	The German Gestational Diabetes Study (PREG), a prospective multicentre cohort study: rationale, methodology and design. BMJ Open, 2022, 12, e058268.	1.9	5
9	Eight weeks of empagliflozin does not affect pancreatic fat content and insulin secretion in people with prediabetes. Diabetes, Obesity and Metabolism, 2022, 24, 1661-1666.	4.4	4
10	Fat Distribution Patterns and Future Type 2 Diabetes. Diabetes, 2022, 71, 1937-1945.	0.6	20
11	Proton magnetic resonance spectroscopy in skeletal muscle: Experts' consensus recommendations. NMR in Biomedicine, 2021, 34, e4266.	2.8	39
12	Elevated Circulating Glutamate Is Associated With Subclinical Atherosclerosis Independently of Established Risk Markers: A Cross-Sectional Study. Journal of Clinical Endocrinology and Metabolism, 2021, 106, e982-e989.	3.6	17
13	Noninvasive, longitudinal imaging-based analysis of body adipose tissue and water composition in a melanoma mouse model and in immune checkpoint inhibitor-treated metastatic melanoma patients. Cancer Immunology, Immunotherapy, 2021, 70, 1263-1275.	4.2	8
14	Long distance running "Can bioprofiling predict success in endurance athletes?". Medical Hypotheses, 2021, 146, 110474.	1.5	3
15	Pathophysiology-based subphenotyping of individuals at elevated risk for type 2 diabetes. Nature Medicine, 2021, 27, 49-57.	30.7	203
16	Lifestyle Intervention Improves Prothrombotic Coagulation Profile in Individuals at High Risk for Type 2 Diabetes. Journal of Clinical Endocrinology and Metabolism, 2021, 106, e3198-e3207.	3.6	8
17	Liver fat scores do not reflect interventional changes in liver fat content induced by high-protein diets. Scientific Reports, 2021, 11, 8843.	3.3	3
18	Hemostatic alterations linked to body fat distribution, fatty liver, and insulin resistance. Molecular Metabolism, 2021, 53, 101262.	6.5	9

#	ARTICLE	IF	CITATIONS
19	Distribution patterns of intramyocellular and extramyocellular fat by magnetic resonance imaging in subjects with diabetes, prediabetes and normoglycaemic controls. <i>Diabetes, Obesity and Metabolism</i> , 2021, 23, 1868-1878.	4.4	14
20	Pancreatic fat cells of humans with type 2 diabetes display reduced adipogenic and lipolytic activity. <i>American Journal of Physiology - Cell Physiology</i> , 2021, 320, C1000-C1012.	4.6	10
21	Determinants of hepatic insulin clearance – Results from a Mendelian Randomization study. <i>Metabolism: Clinical and Experimental</i> , 2021, 119, 154776.	3.4	2
22	Free fatty acids, glicentin and glucose-dependent insulinotropic polypeptide as potential major determinants of fasting substrate oxidation. <i>Scientific Reports</i> , 2021, 11, 16642.	3.3	4
23	Quantification of liver and muscular fat using contrast-enhanced Dual Source Dual Energy Computed Tomography compared to an established multi-echo Dixon MRI sequence. <i>European Journal of Radiology</i> , 2021, 142, 109845.	2.6	9
24	Different Effects of Lifestyle Intervention in High- and Low-Risk Prediabetes: Results of the Randomized Controlled Prediabetes Lifestyle Intervention Study (PLIS). <i>Diabetes</i> , 2021, 70, 2785-2795.	0.6	35
25	Detection of diabetes from whole-body MRI using deep learning. <i>JCI Insight</i> , 2021, 6, .	5.0	10
26	Resolution of severe hepatosteatosis in a cystic fibrosis patient with multifactorial choline deficiency: A case report. <i>Nutrition</i> , 2021, 89, 111348.	2.4	2
27	Elevated circulating follistatin associates with an increased risk of type 2 diabetes. <i>Nature Communications</i> , 2021, 12, 6486.	12.8	31
28	Metabolomic Characteristics of Fatty Pancreas. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2020, 128, 804-810.	1.2	14
29	Empagliflozin Effectively Lowers Liver Fat Content in Well-Controlled Type 2 Diabetes: A Randomized, Double-Blind, Phase 4, Placebo-Controlled Trial. <i>Diabetes Care</i> , 2020, 43, 298-305.	8.6	185
30	Predictive effect of GIPR SNP rs10423928 on glucose metabolism liver fat and adiposity in prediabetic and diabetic subjects. <i>Peptides</i> , 2020, 125, 170237.	2.4	5
31	Magnetic resonance imaging of obesity and metabolic disorders: Summary from the 2019 ISMRM Workshop. <i>Magnetic Resonance in Medicine</i> , 2020, 83, 1565-1576.	3.0	24
32	Normalized Indices Derived from Visceral Adipose Mass Assessed by Magnetic Resonance Imaging and Their Correlation with Markers for Insulin Resistance and Prediabetes. <i>Nutrients</i> , 2020, 12, 2064.	4.1	17
33	High-protein diet more effectively reduces hepatic fat than low-protein diet despite lower autophagy and FGF21 levels. <i>Liver International</i> , 2020, 40, 2982-2997.	3.9	42
34	Lipodystrophic Nonalcoholic Fatty Liver Disease Induced by Immune Checkpoint Blockade. <i>Annals of Internal Medicine</i> , 2020, 172, 836-837.	3.9	44
35	Pancreatic Steatosis Associates With Impaired Insulin Secretion in Genetically Predisposed Individuals. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020, 105, 3518-3525.	3.6	37
36	Investigating obesity-associated brain inflammation using quantitative water content mapping. <i>Journal of Neuroendocrinology</i> , 2020, 32, e12907.	2.6	22

#	ARTICLE	IF	CITATIONS
37	Dietary Rapeseed Oil Supplementation Reduces Hepatic Steatosis in Obese Men—A Randomized Controlled Trial. <i>Molecular Nutrition and Food Research</i> , 2020, 64, e2000419.	3.3	16
38	Response of Mitochondrial Respiration in Adipose Tissue and Muscle to 8 Weeks of Endurance Exercise in Obese Subjects. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020, 105, e4023-e4037.	3.6	23
39	Lack of $GLI2$ proteins in adipocytes attenuates diet-induced obesity. <i>Molecular Metabolism</i> , 2020, 40, 101029.	6.5	10
40	Vertebral Bone Marrow Fat Is Independently Associated to VAT but Not to SAT: KORA FF4—Whole-Body MR Imaging in a Population-Based Cohort. <i>Nutrients</i> , 2020, 12, 1527.	4.1	7
41	Bone marrow fat fraction assessment in regard to physical activity: KORA FF4—3-T MR imaging in a population-based cohort. <i>European Radiology</i> , 2020, 30, 3417-3428.	4.5	19
42	Brain insulin sensitivity is linked to adiposity and body fat distribution. <i>Nature Communications</i> , 2020, 11, 1841.	12.8	81
43	Long-term effects of a food pattern on cardiovascular risk factors and age-related changes of muscular and cognitive function. <i>Medicine (United States)</i> , 2020, 99, e22381.	1.0	2
44	Fully Automated and Standardized Segmentation of Adipose Tissue Compartments via Deep Learning in 3D Whole-Body MRI of Epidemiologic Cohort Studies. <i>Radiology: Artificial Intelligence</i> , 2020, 2, e200010.	5.8	30
45	Applications of Fat Mapping. <i>Advances in Magnetic Resonance Technology and Applications</i> , 2020, 1, 735-777.	0.1	1
46	The Gly385(388)Arg Polymorphism of the FGFR4 Receptor Regulates Hepatic Lipogenesis Under Healthy Diet. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019, 104, 2041-2053.	3.6	8
47	Choline Supplementation in Cystic Fibrosis—The Metabolic and Clinical Impact. <i>Nutrients</i> , 2019, 11, 656.	4.1	16
48	Potential effects of reduced red meat compared with increased fiber intake on glucose metabolism and liver fat content: a randomized and controlled dietary intervention study. <i>American Journal of Clinical Nutrition</i> , 2019, 109, 288-296.	4.7	15
49	Visceral Adiposity Index as an Independent Marker of Subclinical Atherosclerosis in Individuals Prone to Diabetes Mellitus. <i>Journal of Atherosclerosis and Thrombosis</i> , 2019, 26, 821-834.	2.0	36
50	A Polygenic Risk Score of Lipolysis-Increasing Alleles Determines Visceral Fat Mass and Proinsulin Conversion. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019, 104, 1090-1098.	3.6	10
51	Characteristics and associated risk factors of diverticular disease assessed by magnetic resonance imaging in subjects from a Western general population. <i>European Radiology</i> , 2019, 29, 1094-1103.	4.5	10
52	MRI-based assessment and characterization of epicardial and paracardial fat depots in the context of impaired glucose metabolism and subclinical left-ventricular alterations. <i>British Journal of Radiology</i> , 2019, 92, 20180562.	2.2	16
53	Genome-Wide and Abdominal MRI Data Provide Evidence That a Genetically Determined Favorable Adiposity Phenotype Is Characterized by Lower Ectopic Liver Fat and Lower Risk of Type 2 Diabetes, Heart Disease, and Hypertension. <i>Diabetes</i> , 2019, 68, 207-219.	0.6	72
54	Periaortic Adipose Tissue Compared With Peribrachial Adipose Tissue Mass as Markers and Possible Modulators of Cardiometabolic Risk. <i>Angiology</i> , 2018, 69, 854-860.	1.8	11

#	ARTICLE	IF	CITATIONS
55	Effects of resveratrol supplementation on liver fat content in overweight and insulin-resistant subjects: A randomized, double-blind, placebo-controlled clinical trial. <i>Diabetes, Obesity and Metabolism</i> , 2018, 20, 1793-1797.	4.4	66
56	Phenotypic Multiorgan Involvement of Subclinical Disease as Quantified by Magnetic Resonance Imaging in Subjects With Prediabetes, Diabetes, and Normal Glucose Tolerance. <i>Investigative Radiology</i> , 2018, 53, 357-364.	6.2	8
57	Inter- and intra-observer variability of an anatomical landmark-based, manual segmentation method by MRI for the assessment of skeletal muscle fat content and area in subjects from the general population. <i>British Journal of Radiology</i> , 2018, 91, 20180019.	2.2	14
58	The role of visceral and subcutaneous adipose tissue measurements and their ratio by magnetic resonance imaging in subjects with prediabetes, diabetes and healthy controls from a general population without cardiovascular disease. <i>British Journal of Radiology</i> , 2018, 91, 20170808.	2.2	31
59	Leptin Replacement Reestablishes Brain Insulin Action in the Hypothalamus in Congenital Leptin Deficiency. <i>Diabetes Care</i> , 2018, 41, 907-910.	8.6	11
60	Acute Endothelial Benefits of Fat Restriction over Carbohydrate Restriction in Type 2 Diabetes Mellitus: Beyond Carbs and Fats. <i>Nutrients</i> , 2018, 10, 1859.	4.1	9
61	Single Nucleotide Polymorphisms in the G-Protein Coupled Receptor Kinase 5 (GRK5) Gene are associated with Plasma LDL-Cholesterol Levels in Humans. <i>Scientific Reports</i> , 2018, 8, 7745.	3.3	3
62	Assessment of the degree of abdominal myosteatosis by magnetic resonance imaging in subjects with diabetes, prediabetes and healthy controls from the general population. <i>European Journal of Radiology</i> , 2018, 105, 261-268.	2.6	20
63	Association between abdominal adiposity and subclinical measures of left-ventricular remodeling in diabetics, prediabetics and normal controls without history of cardiovascular disease as measured by magnetic resonance imaging: results from the KORA-FF4 Study. <i>Cardiovascular Diabetology</i> , 2018, 17, 88.	6.8	21
64	Genetic determination of body fat distribution and the attributive influence on metabolism. <i>Obesity</i> , 2017, 25, 1277-1283.	3.0	15
65	Intra- and interindividual variability of fatty acid unsaturation in six different human adipose tissue compartments assessed by <sup>1</sup> H-MRS <i>in vivo</i> at 3T. <i>NMR in Biomedicine</i> , 2017, 30, e3744.	2.8	29
66	Body and liver fat content and adipokines in schizophrenia: a magnetic resonance imaging and spectroscopy study. <i>Psychopharmacology</i> , 2017, 234, 1923-1932.	3.1	14
67	Nonsuppressed Glucagon After Glucose Challenge as a Potential Predictor for Glucose Tolerance. <i>Diabetes</i> , 2017, 66, 1373-1379.	0.6	25
68	Metabolic crosstalk between fatty pancreas and fatty liver: effects on local inflammation and insulin secretion. <i>Diabetologia</i> , 2017, 60, 2240-2251.	6.3	100
69	Intranasal insulin enhances brain functional connectivity mediating the relationship between adiposity and subjective feeling of hunger. <i>Scientific Reports</i> , 2017, 7, 1627.	3.3	63
70	Comparison of T1-weighted 2D TSE, 3D SPGR, and two-point 3D Dixon MRI for automated segmentation of visceral adipose tissue at 3 Tesla. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2017, 30, 139-151.	2.0	18
71	Iso-caloric Diets High in Animal or Plant Protein Reduce Liver Fat and Inflammation in Individuals With Type 2 Diabetes. <i>Gastroenterology</i> , 2017, 152, 571-585.e8.	1.3	194
72	Subclinical Disease Burden as Assessed by Whole-Body MRI in Subjects With Prediabetes, Subjects With Diabetes, and Normal Control Subjects From the General Population: The KORA-MRI Study. <i>Diabetes</i> , 2017, 66, 158-169.	0.6	102

#	ARTICLE	IF	CITATIONS
73	Pancreatic fat content by magnetic resonance imaging in subjects with prediabetes, diabetes, and controls from a general population without cardiovascular disease. PLoS ONE, 2017, 12, e0177154.	2.5	54
74	Obesity and renal disease: not all fat is created equal and not all obesity is harmful to the kidneys. Nephrology Dialysis Transplantation, 2016, 31, 726-730.	0.7	40
75	TGF-Î² Contributes to Impaired Exercise Response by Suppression of Mitochondrial Key Regulators in Skeletal Muscle. Diabetes, 2016, 65, 2849-2861.	0.6	62
76	Genetic Variation in the 11Î²-hydroxysteroid-dehydrogenase 1 Gene Determines NAFLD and Visceral Obesity. Journal of Clinical Endocrinology and Metabolism, 2016, 101, 4743-4751.	3.6	20
77	A fully automatic reference deconvolution strategy to increase the accuracy of in vivo lipid signal quantification. Magnetic Resonance in Medicine, 2016, 75, 1391-1401.	3.0	0
78	Glucose-Raising Polymorphisms in the Human Clock Gene Cryptochrome 2 (CRY2) Affect Hepatic Lipid Content. PLoS ONE, 2016, 11, e0145563.	2.5	27
79	Variation in the Phosphoinositide 3-Kinase Gamma Gene Affects Plasma HDL-Cholesterol without Modification of Metabolic or Inflammatory Markers. PLoS ONE, 2015, 10, e0144494.	2.5	22
80	Selective Insulin Resistance in Homeostatic and Cognitive Control Brain Areas in Overweight and Obese Adults. Diabetes Care, 2015, 38, 1044-1050.	8.6	126
81	A high-risk phenotype associates with reduced improvement in glycaemia during a lifestyle intervention in prediabetes. Diabetologia, 2015, 58, 2877-2884.	6.3	56
82	Cinnamon Extract Improves Insulin Sensitivity in the Brain and Lowers Liver Fat in Mouse Models of Obesity. PLoS ONE, 2014, 9, e92358.	2.5	80
83	Common Genetic Variation in the Human CTF1 Locus, Encoding Cardiotrophin-1, Determines Insulin Sensitivity. PLoS ONE, 2014, 9, e100391.	2.5	4
84	Reduced cortical thickness associated with visceral fat and BMI. NeuroImage: Clinical, 2014, 6, 307-311.	2.7	96
85	Modulation of Amino Acid Metabolic Signatures by Supplemented Isoenergetic Diets Differing in Protein and Cereal Fiber Content. Journal of Clinical Endocrinology and Metabolism, 2014, 99, E2599-E2609.	3.6	32
86	Relationships of body composition and liver fat content with insulin resistance in obesity-matched adolescents and adults. Obesity, 2014, 22, 1325-1331.	3.0	35
87	Inhibition of 11Î²-HSD1 with RO5093151 for non-alcoholic fatty liver disease: a multicentre, randomised, double-blind, placebo-controlled trial. Lancet Diabetes and Endocrinology, 2014, 2, 406-416.	11.4	98
88	Impact of the Adipokine Adiponectin and the Hepatokine Fetuin-A on the Development of Type 2 Diabetes: Prospective Cohort- and Cross-Sectional Phenotyping Studies. PLoS ONE, 2014, 9, e92238.	2.5	63
89	Characteristics, changes and influence of body composition during a 4486-km transcontinental ultramarathon: results from the Transeurope Footrace mobile whole body MRI-project. BMC Medicine, 2013, 11, 122.	5.5	28
90	Circulating Lysophosphatidylcholines Are Markers of a Metabolically Benign Nonalcoholic Fatty Liver. Diabetes Care, 2013, 36, 2331-2338.	8.6	100

#	ARTICLE	IF	CITATIONS
91	A moderate weight reduction through dietary intervention decreases hepatic fat content in patients with non-alcoholic fatty liver disease (NAFLD): a pilot study. <i>European Journal of Nutrition</i> , 2013, 52, 527-535.	3.9	71
92	Diagnostic imaging in obesity. <i>Best Practice and Research in Clinical Endocrinology and Metabolism</i> , 2013, 27, 261-277.	4.7	35
93	Fraction of unsaturated fatty acids in visceral adipose tissue (VAT) is lower in subjects with high total VAT volume – a combined <sup>1</sup> H MRS and volumetric MRI study in male subjects. <i>NMR in Biomedicine</i> , 2013, 26, 232-236.	2.8	28
94	Genetic Variation in <i>NR1H4</i> Encoding the Bile Acid Receptor FXR Determines Fasting Glucose and Free Fatty Acid Levels in Humans. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013, 98, E1224-E1229.	3.6	24
95	Common Genetic Variation in the Human <i>FNDC5</i> Locus, Encoding the Novel Muscle-Derived “Browning” Factor Irisin, Determines Insulin Sensitivity. <i>PLoS ONE</i> , 2013, 8, e61903.	2.5	83
96	Monounsaturated Fatty Acids Prevent the Aversive Effects of Obesity on Locomotion, Brain Activity, and Sleep Behavior. <i>Diabetes</i> , 2012, 61, 1669-1679.	0.6	48
97	Predicting volumes of metabolically important whole-body adipose tissue compartments in overweight and obese adolescents by different MRI approaches and anthropometry. <i>European Journal of Radiology</i> , 2012, 81, 1488-1494.	2.6	20
98	The Transeurope Footrace Project: longitudinal data acquisition in a cluster randomized mobile MRI observational cohort study on 44 endurance runners at a 64-stage 4,486km transcontinental ultramarathon. <i>BMC Medicine</i> , 2012, 10, 78.	5.5	47
99	High Hepatic SCD1 Activity Is Associated with Low Liver Fat Content in Healthy Subjects under a Lipogenic Diet. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012, 97, E2288-E2292.	3.6	66
100	Common Genetic Variation in the <i>SERPINF1</i> Locus Determines Overall Adiposity, Obesity-Related Insulin Resistance, and Circulating Leptin Levels. <i>PLoS ONE</i> , 2012, 7, e34035.	2.5	28
101	Quantifying the Improvement of Surrogate Indices of Hepatic Insulin Resistance Using Complex Measurement Techniques. <i>PLoS ONE</i> , 2012, 7, e39029.	2.5	16
102	Magnetic resonance techniques for mapping fat deposits and directing therapy. <i>Clinical Lipidology</i> , 2011, 6, 93-107.	0.4	1
103	Assessment of relevant hepatic steatosis in obese adolescents by rapid fat-selective GRE imaging with spatial-spectral excitation: a quantitative comparison with spectroscopic findings. <i>European Radiology</i> , 2011, 21, 816-822.	4.5	9
104	Morning to evening changes of intramyocellular lipid content in dependence on nutrition and physical activity during one single day: a volume selective <sup>1</sup> H-MRS study. <i>Magnetic Resonance Materials in Physics, Biology, and Medicine</i> , 2011, 24, 29-33.	2.0	23
105	Insulin Sensitivity and Liver Fat: Role of Iron Load. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011, 96, E958-E961.	3.6	40
106	Effects of supplemented isoenergetic diets differing in cereal fiber and protein content on insulin sensitivity in overweight humans. <i>American Journal of Clinical Nutrition</i> , 2011, 94, 459-471.	4.7	148
107	Genetic Ablation of cGMP-Dependent Protein Kinase Type I Causes Liver Inflammation and Fasting Hyperglycemia. <i>Diabetes</i> , 2011, 60, 1566-1576.	0.6	34
108	Quantitative Assessment of Intrahepatic Lipids Using Fat-Selective Imaging With Spectral-Spatial Excitation and In-/Opposed-Phase Gradient Echo Imaging Techniques Within a Study Population of Extremely Obese Patients. <i>Investigative Radiology</i> , 2010, 45, 484-490.	6.2	14



#	ARTICLE	IF	CITATIONS
109	Quantitative Analysis of Adipose Tissue in Single Transverse Slices for Estimation of Volumes of Relevant Fat Tissue Compartments. <i>Investigative Radiology</i> , 2010, 45, 788-794.	6.2	53
110	Quantitative Assessment of Visceral Fat in Morbidly Obese Patients by Means of Wide-Bore MRI and its Relation to Lower Esophageal Sphincter Pressure and Signs of Gastroesophageal Reflux. <i>Obesity Surgery</i> , 2010, 20, 749-756.	2.1	9
111	Enforced expression of protein kinase C in skeletal muscle causes physical inactivity, fatty liver and insulin resistance in the brain. <i>Journal of Cellular and Molecular Medicine</i> , 2010, 14, 903-913.	3.6	16
112	Pancreatic fat is negatively associated with insulin secretion in individuals with impaired fasting glucose and/or impaired glucose tolerance: a nuclear magnetic resonance study. <i>Diabetes/Metabolism Research and Reviews</i> , 2010, 26, 200-205.	4.0	212
113	Topography mapping of whole body adipose tissue using A fully automated and standardized procedure. <i>Journal of Magnetic Resonance Imaging</i> , 2010, 31, 430-439.	3.4	82
114	Novel Obesity Risk Loci Do Not Determine Distribution of Body Fat Depots: A Whole-body MRI/MRS study. <i>Obesity</i> , 2010, 18, 1212-1217.	3.0	30
115	Medium Chain Acylcarnitines Dominate the Metabolite Pattern in Humans under Moderate Intensity Exercise and Support Lipid Oxidation. <i>PLoS ONE</i> , 2010, 5, e11519.	2.5	118
116	Circulating Palmitoleate Strongly and Independently Predicts Insulin Sensitivity in Humans. <i>Diabetes Care</i> , 2010, 33, 405-407.	8.6	130
117	Exercise-induced normalization of decreased BDNF serum concentration in elderly women with remitted major depression. <i>International Journal of Neuropsychopharmacology</i> , 2010, 13, 595-602.	2.1	142
118	Relationships of Circulating Sex Hormone-binding Globulin With Metabolic Traits in Humans. <i>Diabetes</i> , 2010, 59, 3167-3173.	0.6	130
119	Normal-Weight 14-Year-Old Girl with Acanthosis Nigricans and Markedly Increased Hepatic Steatosis: Evidence for the Important Role of Ectopic Fat Deposition in the Pathogenesis of Insulin Resistance in Childhood and Adolescence. <i>Hormone Research in Paediatrics</i> , 2010, 74, 376-380.	1.8	4
120	Interscapular Fat Is Strongly Associated with Insulin Resistance. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2010, 95, 4736-4742.	3.6	21
121	Follow-up Whole-Body Assessment of Adipose Tissue Compartments during a Lifestyle Intervention in a Large Cohort at Increased Risk for Type 2 Diabetes. <i>Radiology</i> , 2010, 257, 353-363.	7.3	105
122	Gene Variants of <i>TCF7L2</i> Influence Weight Loss and Body Composition During Lifestyle Intervention in a Population at Risk for Type 2 Diabetes. <i>Diabetes</i> , 2010, 59, 747-750.	0.6	69
123	The D299G/T399I Toll-Like Receptor 4 Variant Associates with Body and Liver Fat: Results from the TULIP and METSIM Studies. <i>PLoS ONE</i> , 2010, 5, e13980.	2.5	27
124	Liver fat content determined by magnetic resonance imaging and spectroscopy. <i>World Journal of Gastroenterology</i> , 2010, 16, 1560.	3.3	107
125	Muscle-Derived Angiotensin-Like Protein 4 Is Induced by Fatty Acids via Peroxisome Proliferator-Activated Receptor (PPAR)- $\delta$ and Is of Metabolic Relevance in Humans. <i>Diabetes</i> , 2009, 58, 579-589.	0.6	166
126	Dissociation Between Fatty Liver and Insulin Resistance in Humans Carrying a Variant of the Patatin-Like Phospholipase 3 Gene. <i>Diabetes</i> , 2009, 58, 2616-2623.	0.6	291



#	ARTICLE	IF	CITATIONS
127	Individual Stearoyl-CoA Desaturase 1 Expression Modulates Endoplasmic Reticulum Stress and Inflammation in Human Myotubes and Is Associated With Skeletal Muscle Lipid Storage and Insulin Sensitivity In Vivo. <i>Diabetes</i> , 2009, 58, 1757-1765.	0.6	134
128	The Insulin Effect on Cerebrocortical Theta Activity Is Associated with Serum Concentrations of Saturated Nonesterified Fatty Acids. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2009, 94, 4600-4607.	3.6	40
129	RARRES2, encoding the novel adipokine chemerin, is a genetic determinant of disproportionate regional body fat distribution: a comparative magnetic resonance imaging study. <i>Metabolism: Clinical and Experimental</i> , 2009, 58, 519-524.	3.4	53
130	Preliminary report: genetic variation within the GPBAR1 gene is not associated with metabolic traits in white subjects at an increased risk for type 2 diabetes mellitus. <i>Metabolism: Clinical and Experimental</i> , 2009, 58, 1809-1811.	3.4	18
131	Intermuscular adipose tissue (IMAT): Association with other adipose tissue compartments and insulin sensitivity. <i>Journal of Magnetic Resonance Imaging</i> , 2009, 29, 1340-1345.	3.4	160
132	Aging effects on human calf muscle properties assessed by MRI at 3 Tesla. <i>Journal of Magnetic Resonance Imaging</i> , 2009, 29, 1346-1354.	3.4	65
133	Impact of Variation Near <i>MC4R</i> on Whole-body Fat Distribution, Liver Fat, and Weight Loss. <i>Obesity</i> , 2009, 17, 1942-1945.	3.0	48
134	Non-invasive assessment and quantification of liver steatosis by ultrasound, computed tomography and magnetic resonance. <i>Journal of Hepatology</i> , 2009, 51, 433-445.	3.7	667
135	High plasma fetuin-A is associated with increased carotid intima-media thickness in a middle-aged population. <i>Atherosclerosis</i> , 2009, 207, 341-342.	0.8	58
136	Technical challenges and opportunities of whole-body magnetic resonance imaging at 3T. <i>Physica Medica</i> , 2008, 24, 63-70.	0.7	28
137	Impact of Variation in the <i>FTO</i> Gene on Whole Body Fat Distribution, Ectopic Fat, and Weight Loss. <i>Obesity</i> , 2008, 16, 1969-1972.	3.0	102
138	1H MR spectroscopy of skeletal muscle, liver and bone marrow. <i>European Journal of Radiology</i> , 2008, 67, 275-284.	2.6	97
139	Variations in <i>PPAR<math>\gamma</math></i> Determine the Change in Body Composition during Lifestyle Intervention: A Whole-Body Magnetic Resonance Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008, 93, 1497-1500.	3.6	71
140	Identification and Characterization of Metabolically Benign Obesity in Humans. <i>Archives of Internal Medicine</i> , 2008, 168, 1609.	3.8	869
141	Fatty Liver Is Independently Associated With Alterations in Circulating HDL2 and HDL3 Subfractions. <i>Diabetes Care</i> , 2008, 31, 366-368.	8.6	55
142	T2* Relaxometry in Liver, Pancreas, and Spleen in a Healthy Cohort of One Hundred Twenty-Nine Subjectsâ€“Correlation With Age, Gender, and Serum Ferritin. <i>Investigative Radiology</i> , 2008, 43, 854-860.	6.2	89
143	Quantification of Pancreatic Lipomatosis and Liver Steatosis by MRI: Comparison of In/Opposed-Phase and Spectral-Spatial Excitation Techniques. <i>Investigative Radiology</i> , 2008, 43, 330-337.	6.2	104
144	High Circulating Retinol-Binding Protein 4 Is Associated With Elevated Liver Fat but Not With Total, Subcutaneous, Visceral, or Intramyocellular Fat in Humans. <i>Diabetes Care</i> , 2007, 30, 1173-1178.	8.6	203

#	ARTICLE	IF	CITATIONS
145	Genetic Variations in <i>PPARD</i> and <i>PPARGC1A</i> Determine Mitochondrial Function and Change in Aerobic Physical Fitness and Insulin Sensitivity during Lifestyle Intervention. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2007, 92, 1827-1833.	3.6	123
146	Impact of Different Fat Depots on Insulin Sensitivity: Predominant Role of Liver Fat. <i>Journal of Diabetes Science and Technology</i> , 2007, 1, 753-759.	2.2	18
147	Upstream transcription factor 1 gene polymorphisms are associated with high antilipolytic insulin sensitivity and show gene-gene interactions. <i>Journal of Molecular Medicine</i> , 2006, 85, 55-61.	3.9	24
148	Hepatic lipid accumulation in healthy subjects: A comparative study using spectral fat-selective MRI and volume-localized <sup>1</sup> H-MR spectroscopy. <i>Magnetic Resonance in Medicine</i> , 2006, 55, 913-917.	3.0	146
149	Magnetic Resonance Techniques for Assessment of Body Components. <i>Hormone Research in Paediatrics</i> , 2006, 66, 65-72.	1.8	3
150	Î²2-Heremans-Schmid Glycoprotein/ Fetuin-A Is Associated With Insulin Resistance and Fat Accumulation in the Liver in Humans. <i>Diabetes Care</i> , 2006, 29, 853-857.	8.6	440
151	3D proton MR spectroscopic imaging of prostate cancer using a standard spine coil at 1.5i½T in clinical routine: a feasibility study. <i>European Radiology</i> , 2005, 15, 653-660.	4.5	21
152	Standardized assessment of whole body adipose tissue topography by MRI. <i>Journal of Magnetic Resonance Imaging</i> , 2005, 21, 455-462.	3.4	216
153	New Imaging Techniques of Fat, Muscle and Liver within the Context of Determining Insulin Sensitivity. <i>Hormone Research in Paediatrics</i> , 2005, 64, 38-44.	1.8	37
154	Intrahepatic Lipids Are Predicted by Visceral Adipose Tissue Mass in Healthy Subjects. <i>Diabetes Care</i> , 2004, 27, 2726-2729.	8.6	69
155	Multiple Symmetric Lipomatosis. <i>Diabetes Care</i> , 2004, 27, 794-795.	8.6	27
156	Intramyoellular lipids and insulin resistance. <i>Diabetes, Obesity and Metabolism</i> , 2004, 6, 239-248.	4.4	153
157	Lipid content in the musculature of the lower leg assessed by fat selective MRI: Intra- and interindividual differences and correlation with anthropometric and metabolic data. <i>Journal of Magnetic Resonance Imaging</i> , 2003, 17, 350-357.	3.4	67
158	Relationship of Serum Adiponectin and Leptin Concentrations with Body Fat Distribution in Humans. <i>Obesity</i> , 2003, 11, 368-376.	4.0	195
159	Intramyoellular Lipids: Anthropometric Determinants and Relationships with Maximal Aerobic Capacity and Insulin Sensitivity. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2003, 88, 1785-1791.	3.6	210
160	In Vivo Proton NMR Studies in Skeletal Musculature. <i>Annual Reports on NMR Spectroscopy</i> , 2003, 50, 1-74.	1.5	8
161	MRI of muscular fat. <i>Magnetic Resonance in Medicine</i> , 2002, 47, 720-727.	3.0	93
162	Osteodensitometry of human heel bones by MR spin-echo imaging: Comparison with MR gradient-echo imaging and quantitative computed tomography. <i>Journal of Magnetic Resonance Imaging</i> , 2001, 14, 147-155.	3.4	14

#	ARTICLE	IF	CITATIONS
163	Magnetic Resonance Osteodensitometry in Human Heel Bones. Investigative Radiology, 2000, 35, 393-400.	6.2	11
164	Improved clinical echo-planar MRI using spatial-spectral excitation. Journal of Magnetic Resonance Imaging, 1998, 8, 960-967.	3.4	28
165	Highly selective water and fat imaging applying multislice sequences without sensitivity to B1 field inhomogeneities. Magnetic Resonance in Medicine, 1997, 38, 269-274.	3.0	63
166	Sequence parameters of double spin-echo sequences affect quantification of citrate. Magnetic Resonance Imaging, 1996, 14, 663-672.	1.8	5