
## Hyunil Ha

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/453577/publications.pdf Version: 2024-02-01



Ηγιινίι Ηλ

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Haptoglobin Acts as a TLR4 Ligand to Suppress Osteoclastogenesis via the TLR4–IFN-β Axis. Journal of<br>Immunology, 2019, 202, 3359-3369.   | 0.8 | 8         |
| 2  | JN-2, a C-X-C motif chemokine receptor 3 antagonist, ameliorates arthritis progression in an animal model. European Journal of Pharmacology, 2018, 823, 1-10.   | 3.5 | 14        |
| 3  | Data on the expression of CXCR3 ligands and pro-inflammatory cytokines in macrophages and CD4+ T<br>cells after stimuli of CXCR3 ligands. Data in Brief, 2018, 18, 518-522.                                 | 1.0 | 0         |
| 4  | Trapidil induces osteogenesis by upregulating the signaling of bone morphogenetic proteins. Cellular<br>Signalling, 2018, 49, 68-78.  | 3.6 | 4         |
| 5  | Pathogenic roles of CXCL10 signaling through CXCR3 and TLR4 in macrophages and T cells: relevance for arthritis. Arthritis Research and Therapy, 2017, 19, 163.   | 3.5 | 104       |
| 6  | A water extract of Malva verticillata seeds suppresses osteoclastogenesis and bone resorption stimulated by RANK ligand. BMC Complementary and Alternative Medicine, 2016, 16, 332.                         | 3.7 | 10        |
| 7  | Trolox inhibits osteolytic bone metastasis of breast cancer through both PGE2-dependent and independent mechanisms. Biochemical Pharmacology, 2014, 91, 51-60.  | 4.4 | 25        |
| 8  | Trapidil, a platelet-derived growth factor antagonist, inhibits osteoclastogenesis by down-regulating NFATc1 and suppresses bone loss in mice. Biochemical Pharmacology, 2013, 86, 782-790.                 | 4.4 | 19        |
| 9  | Water extract of Spatholobus suberectus inhibits osteoclast differentiation and bone resorption.<br>BMC Complementary and Alternative Medicine, 2013, 13, 112.  | 3.7 | 26        |
| 10 | The anti-osteoporotic effect of Yijung-tang in an ovariectomized rat model mediated by inhibition of osteoclast differentiation. Journal of Ethnopharmacology, 2013, 146, 83-89.                            | 4.1 | 16        |
| 11 | Hwangryun-Haedok-Tang Fermented withLactobacillus caseiSuppresses Ovariectomy-Induced Bone<br>Loss. Evidence-based Complementary and Alternative Medicine, 2012, 2012, 1-12.                                | 1.2 | 11        |
| 12 | <i>In Vitro</i> and <i>In Vivo</i> Genotoxicity Assessment of <i>Aristolochia manshuriensis</i> Kom<br>Evidence-based Complementary and Alternative Medicine, 2012, 2012, 1-9.                              | 1.2 | 15        |
| 13 | CXCL10 Promotes Osteolytic Bone Metastasis by Enhancing Cancer Outgrowth and Osteoclastogenesis. Cancer Research, 2012, 72, 3175-3186.  | 0.9 | 97        |
| 14 | MS-275, a benzamide histone deacetylase inhibitor, prevents osteoclastogenesis by down-regulating<br>c-Fos expression and suppresses bone loss in mice. European Journal of Pharmacology, 2012, 691, 69-76. | 3.5 | 21        |
| 15 | Aristolochia Manshuriensis Kom Inhibits Adipocyte Differentiation by Regulation of ERK1/2 and Akt<br>Pathway. PLoS ONE, 2012, 7, e49530.  | 2.5 | 27        |
| 16 | α-Tocotrienol inhibits osteoclastic bone resorption by suppressing RANKL expression and signaling and bone resorbing activity. Biochemical and Biophysical Research Communications, 2011, 406, 546-551.     | 2.1 | 43        |
| 17 | Histone deacetylase inhibitor MS-275 stimulates bone formation in part by enhancing Dhx36-mediated TNAP transcription. Journal of Bone and Mineral Research, 2011, 26, 2161-2173.                           | 2.8 | 48        |
| 18 | Epigallocatechin-3-gallate Inhibits Osteoclastogenesis by Down-Regulating c-Fos Expression and<br>Suppressing the Nuclear Factor-κB Signal. Molecular Pharmacology, 2010, 77, 17-25.                        | 2.3 | 117       |
|    |   |     |           |

Hyunil Ha

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | cAMP-response-element-binding protein positively regulates breast cancer metastasis and subsequent bone destruction. Biochemical and Biophysical Research Communications, 2010, 398, 309-314.   | 2.1 | 43        |
| 20 | ATP6v0d2 deficiency increases bone mass, but does not influence ovariectomy-induced bone loss.<br>Biochemical and Biophysical Research Communications, 2010, 403, 73-78.  | 2.1 | 24        |
| 21 | Trolox Prevents Osteoclastogenesis by Suppressing RANKL Expression and Signaling. Journal of<br>Biological Chemistry, 2009, 284, 13725-13734.   | 3.4 | 60        |
| 22 | Trichostatin A inhibits osteoclastogenesis and bone resorption by suppressing the induction of c-Fos by RANKL. European Journal of Pharmacology, 2009, 623, 22-29.  | 3.5 | 33        |
| 23 | Pyridone 6, A Pan-Janus-Activated Kinase Inhibitor, Suppresses Osteoclast Formation and Bone<br>Resorption through Down-Regulation of Receptor Activator of Nuclear FactorKAPPA.B (NFKAPPA.B)<br>Ligand (RANKL)-Induced c-Fos and Nuclear Factor of Activated T Cells (NFAT) c1 Expression. Biological<br>and Pharmaceutical Bulletin. 2009. 32. 45-50. | 1.4 | 17        |
| 24 | Reciprocal cross-talk between RANKL and interferon-γ–inducible protein 10 is responsible for bone-erosive experimental arthritis. Arthritis and Rheumatism, 2008, 58, 1332-1342.  | 6.7 | 105       |
| 25 | Tanshinone IIA suppresses inflammatory bone loss by inhibiting the synthesis of prostaglandin E2 in osteoblasts. European Journal of Pharmacology, 2008, 601, 30-37.  | 3.5 | 31        |
| 26 | Stimulation by TLR5 Modulates Osteoclast Differentiation through STAT1/IFN-β. Journal of Immunology, 2008, 180, 1382-1389.  | 0.8 | 47        |
| 27 | α-Lipoic Acid Inhibits Inflammatory Bone Resorption by Suppressing Prostaglandin E2 Synthesis. Journal of Immunology, 2006, 176, 111-117.   | 0.8 | 83        |
| 28 | Tanshinone IIA inhibits osteoclast differentiation through down-regulation of c-Fos and NFATc1.<br>Experimental and Molecular Medicine, 2006, 38, 256-264.  | 7.7 | 64        |
| 29 | Monokine induced by interferon-Â is induced by receptor activator of nuclear factor ÂB ligand and is involved in osteoclast adhesion and migration. Blood, 2005, 105, 2963-2969.  | 1.4 | 60        |
| 30 | Modulation of the caveolin-3 and Akt status in caveolae by insulin resistance in H9c2 cardiomyoblasts.<br>Experimental and Molecular Medicine, 2005, 37, 169-178.   | 7.7 | 26        |
| 31 | Tumor necrosis factor.α induces differentiation of human peripheral blood mononuclear cells into osteoclasts through the induction of p21(WAF1/Cip1). Biochemical and Biophysical Research Communications, 2005, 330, 1080-1086.  | 2.1 | 26        |
| 32 | Inhibition of osteoclast differentiation and bone resorption by tanshinone IIA isolated from Salvia miltiorrhiza Bunge. Biochemical Pharmacology, 2004, 67, 1647-1656.  | 4.4 | 72        |
| 33 | Reactive oxygen species mediate RANK signaling in osteoclasts. Experimental Cell Research, 2004, 301, 119-127.  | 2.6 | 289       |
| 34 | Membrane Rafts Play a Crucial Role in Receptor Activator of Nuclear Factor κB Signaling and<br>Osteoclast Function. Journal of Biological Chemistry, 2003, 278, 18573-18580.  | 3.4 | 80        |
| 35 | Lipid rafts are important for the association of RANK and TRAF6. Experimental and Molecular Medicine, 2003, 35, 279-284.  | 7.7 | 32        |