

Nadia Duarte

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4539237/publications.pdf>

Version: 2024-02-01

18
papers

638
citations

932766

10
h-index

839053

18
g-index

24
all docs

24
docs citations

24
times ranked

1119
citing authors

#	ARTICLE	IF	CITATIONS
1	Skin and Peripheral Lymph Node Invariant NKT Cells Are Mainly Retinoic Acid Receptor-Related Orphan Receptor I^3t^+ and Respond Preferentially under Inflammatory Conditions. <i>Journal of Immunology</i> , 2009, 183, 2142-2149.	0.4	140
2	How Inflammation Impinges on NAFLD: A Role for Kupffer Cells. <i>BioMed Research International</i> , 2015, 2015, 1-11.	0.9	100
3	Prevention of Diabetes in Nonobese Diabetic Mice Mediated by CD1d-Restricted Nonclassical NKT Cells. <i>Journal of Immunology</i> , 2004, 173, 3112-3118.	0.4	98
4	iNKT cell development is orchestrated by different branches of TGF- β^2 signaling. <i>Journal of Experimental Medicine</i> , 2009, 206, 1365-1378.	4.2	81
5	Trem-2 Promotes Emergence of Restorative Macrophages and Endothelial Cells During Recovery From Hepatic Tissue Damage. <i>Frontiers in Immunology</i> , 2020, 11, 616044.	2.2	34
6	Diabetes Protection and Restoration of Thymocyte Apoptosis in NOD Idd6 Congenic Strains. <i>Diabetes</i> , 2003, 52, 1677-1682.	0.3	29
7	Production of high-quality SARS-CoV-2 antigens: Impact of bioprocess and storage on glycosylation, biophysical attributes, and ELISA serologic tests performance. <i>Biotechnology and Bioengineering</i> , 2021, 118, 2202-2219.	1.7	27
8	Population homogeneity for the antibody response to COVID-19 BNT162b2/Comirnaty vaccine is only reached after the second dose across all adult age ranges. <i>Nature Communications</i> , 2022, 13, 140.	5.8	22
9	Defective Induction of CTLA-4 in the NOD Mouse Is Controlled by the NOD Allele of Idd3/IL-2 and a Novel Locus (Ctex) Telomeric on Chromosome 1. <i>Diabetes</i> , 2006, 55, 538-544.	0.3	21
10	A New Mouse Model That Spontaneously Develops Chronic Liver Inflammation and Fibrosis. <i>PLoS ONE</i> , 2016, 11, e0159850.	1.1	11
11	Autoimmunity Triggers in the NOD Mouse. <i>Annals of the New York Academy of Sciences</i> , 2009, 1173, 442-448.	1.8	10
12	Dipeptidyl Peptidase-4 Is a Pro-Recovery Mediator During Acute Hepatotoxic Damage and Mirrors Severe Shifts in Kupffer Cells. <i>Hepatology Communications</i> , 2018, 2, 1080-1094.	2.0	10
13	The MHC locus controls size variations in the CD4 compartment of the mouse thymus. <i>Immunogenetics</i> , 2001, 53, 662-668.	1.2	8
14	The Idd6.2 diabetes susceptibility region controls defective expression of the Lrmp gene in nonobese diabetic (NOD) mice. <i>Immunogenetics</i> , 2007, 59, 407-416.	1.2	8
15	Insights into Macrophage/Monocyte-Endothelial Cell Crosstalk in the Liver: A Role for Trem-2. <i>Journal of Clinical Medicine</i> , 2021, 10, 1248.	1.0	7
16	Immunoglobulin M gene association with autoantibody reactivity and type 1 diabetes. <i>Immunogenetics</i> , 2017, 69, 429-437.	1.2	6
17	Longitudinal Analysis of Antibody Responses to the mRNA BNT162b2 Vaccine in Patients Undergoing Maintenance Hemodialysis: A 6-Month Follow-Up. <i>Frontiers in Medicine</i> , 2021, 8, 796676.	1.2	6
18	Prediabetes blunts DPP4 genetic control of postprandial glycaemia and insulin secretion. <i>Diabetologia</i> , 2022, 65, 861-871.	2.9	3