

Adriano Soares RÃ^ãgo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/454544/publications.pdf>

Version: 2024-02-01

12
papers

48
citations

1937685

4
h-index

1720034

7
g-index

12
all docs

12
docs citations

12
times ranked

53
citing authors

#	ARTICLE	IF	CITATIONS
1	BIOLOGICAL ASPECTS OF <i>Ceraeochrysa everes</i> (NEUROPTERA: CHRYSOPIDAE) FED ON PINK HIBISCUS MEALYBUG. <i>Revista Caatinga</i> , 2022, 35, 363-370.	0.7	0
2	Efficacy of <i>Amblyseius largoensis</i> (Muma) as a biocontrol agent of the red palm mite <i>Raoiella indica</i> (Acari: Tenuipalpidae). <i>Phytoparasitica</i> , 2021, 49, 103-111.	1.2	4
3	Compatibility of degummed soybean and babassu oils with the generalist predatory mite <i>Typhlodromus</i> (<i>Anthoseius</i>) <i>ornatus</i> (Acari: Phytoseiidae) preying on <i>Aceria guerreronis</i> (Acari: Eriophyidae). <i>International Journal of Acarology</i> , 2021, 47, 242-247.	0.7	0
4	The potential of <i>Beauveria bassiana</i> to control <i>Raoiella indica</i> (Acari: Tenuipalpidae) and its compatibility with predatory mites. <i>Crop Protection</i> , 2021, 149, 105776.	2.1	2
5	Lethal and sublethal effects of babassu and degummed soybean oils on the predatory mite <i>Typhlodromus ornatus</i> (Acari: Phytoseiidae). <i>International Journal of Acarology</i> , 2020, 46, 180-184.	0.7	2
6	Resistance elicitors and defense response enhancers of maize to <i>Spodoptera frugiperda</i> (J.E.Smith) (Lepidoptera: Noctuidae). <i>Australian Journal of Crop Science</i> , 2019, , 1001-1008.	0.3	2
7	The lacewing <i>Ceraeochrysa caligata</i> as a potential biological agent for controlling the red palm mite <i>Raoiella indica</i> . <i>PeerJ</i> , 2019, 7, e7123.	2.0	8
8	DoenÃ§as da vinagreira no Estado do MaranhÃ£o. <i>Summa Phytopathologica</i> , 2014, 40, 378-380.	0.1	3
9	Functional response of <i>Euseius concordis</i> to densities of different developmental stages of the cassava green mite. <i>Experimental and Applied Acarology</i> , 2014, 64, 277-286.	1.6	11
10	Relative contribution of biotic and abiotic factors to the population density of the cassava green mite, <i>Mononychellus tanajoa</i> (Acari: Tetranychidae). <i>Experimental and Applied Acarology</i> , 2013, 60, 479-484.	1.6	11
11	Toxicity of neem oil to the cassava green mite <i>Mononychellus tanajoa</i> (Bondar) (Acari: Tetranychidae). <i>Chilean Journal of Agricultural Research</i> , 2013, 73, 315-319.	1.1	5
12	Natural Biological Control of Pest Mites in Brazilian Sun Coffee Agroecosystems. <i>Recent Patents on Food, Nutrition & Agriculture</i> , 2010, 2, 160-165.	0.9	0