

Magda Babina

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4550316/publications.pdf>

Version: 2024-02-01

81
papers

5,511
citations

172386

29
h-index

91828

69
g-index

82
all docs

82
docs citations

82
times ranked

10183
citing authors

#	ARTICLE	IF	CITATIONS
1	Mast cells instruct keratinocytes to produce thymic stromal lymphopoietin: Relevance of the tryptase/protease-activated receptor 2 axis. <i>Journal of Allergy and Clinical Immunology</i> , 2022, 149, 2053-2061.e6.	1.5	14
2	Fc μ RI ϵ and MRGPRX2 ϵ evoked acute degranulation responses are fully additive in human skin mast cells. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2022, 77, 1906-1909.	2.7	14
3	MRGPRX2-Mediated Degranulation of Human Skin Mast Cells Requires the Operation of G \pm i, G \pm q, Ca $^{++}$ Channels, ERK1/2 and PI3K ϵ Interconnection between Early and Late Signaling. <i>Cells</i> , 2022, 11, 953.	1.8	17
4	Tolerance induction through non-avoidance to prevent persistent food allergy (TINA) in children and adults with peanut or tree nut allergy: rationale, study design and methods of a randomized controlled trial and observational cohort study. <i>Trials</i> , 2022, 23, 236.	0.7	7
5	Tolerance induction through early feeding to prevent food allergy in infants with eczema (TEFFA): rationale, study design, and methods of a randomized controlled trial. <i>Trials</i> , 2022, 23, 210.	0.7	8
6	Serological profiling reveals hsa-miR-451a as a possible biomarker of anaphylaxis. <i>JCI Insight</i> , 2022, 7, .	2.3	9
7	The <sc>SCF</sc>/<sc>KIT</sc> axis in human mast cells: Capicua acts as potent <sc>KIT</sc> repressor and <sc>ERK</sc> predominates <sc>PI3K</sc>. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2022, 77, 3337-3349.	2.7	8
8	Diversities of allergic pathologies and their modifiers: Report from the second DGAKI-JSA meeting. <i>Allergology International</i> , 2022, 71, 310-317.	1.4	1
9	Mas-related G protein ϵ coupled receptor X2 and its activators in dermatologic allergies. <i>Journal of Allergy and Clinical Immunology</i> , 2021, 147, 456-469.	1.5	70
10	MRGPRX2 Is the Codeine Receptor of Human Skin Mast Cells: Desensitization through β -Arrestin and Lack of Correlation with the Fc μ RI Pathway. <i>Journal of Investigative Dermatology</i> , 2021, 141, 1286-1296.e4.	0.3	39
11	Thymic Stromal Lymphopoietin Promotes MRGPRX2-Triggered Degranulation of Skin Mast Cells in a STAT5-Dependent Manner with Further Support from JNK. <i>Cells</i> , 2021, 10, 102.	1.8	24
12	Cytokines Stimulated by IL-33 in Human Skin Mast Cells: Involvement of NF- κ B and p38 at Distinct Levels and Potent Co-Operation with Fc μ RI and MRGPRX2. <i>International Journal of Molecular Sciences</i> , 2021, 22, 3580.	1.8	33
13	Cytokine Stimulation via MRGPRX2 Occurs with Lower Potency than by Fc μ RI-aggregation but with Similar Dependence on the ERK1/2 Module in Human Skin Mast Cells. <i>Journal of Investigative Dermatology</i> , 2021, , .	0.3	8
14	Thymic stromal lymphopoietin production induced by skin irritation results from concomitant activation of protease ϵ activated receptor 2 and interleukin 1 pathways. <i>British Journal of Dermatology</i> , 2020, 182, 119-129.	1.4	12
15	MRGPRX2 signals its importance in cutaneous mast cell biology: Does MRGPRX2 connect mast cells and atopic dermatitis?. <i>Experimental Dermatology</i> , 2020, 29, 1104-1111.	1.4	35
16	PGE2 deficiency predisposes to anaphylaxis by causing mast cell hyperresponsiveness. <i>Journal of Allergy and Clinical Immunology</i> , 2020, 146, 1387-1396.e13.	1.5	31
17	Mechanisms governing the pioneering and redistribution capabilities of the non-classical pioneer PU.1. <i>Nature Communications</i> , 2020, 11, 402.	5.8	76
18	The pseudo-allergic/neurogenic route of mast cell activation via MRGPRX2: discovery, functional programs, regulation, relevance to disease, and relation with allergic stimulation. <i>Itch (Philadelphia, Pa)</i> 2020, 10, 10.	0.0	0

#	ARTICLE	IF	CITATIONS
19	Integration of the Human Dermal Mast Cell into the Organotypic Co-culture Skin Model. <i>Methods in Molecular Biology</i> , 2020, 2163, 91-107.	0.4	2
20	Olive Oil-Derived Polyphenols Effectively Attenuate Inflammatory Responses of Human Keratinocytes by Interfering with the NF- κ B Pathway. <i>Molecular Nutrition and Food Research</i> , 2019, 63, 1900019.	1.5	20
21	Thymic Stromal Lymphopoietin Interferes with the Apoptosis of Human Skin Mast Cells by a Dual Strategy Involving STAT5/Mcl-1 and JNK/Bcl-xL. <i>Cells</i> , 2019, 8, 829.	1.8	24
22	An Allosteric Anti-tryptase Antibody for the Treatment of Mast Cell-Mediated Severe Asthma. <i>Cell</i> , 2019, 179, 417-431.e19.	13.5	76
23	Yin-Yang of IL-33 in Human Skin Mast Cells: Reduced Degranulation, but Augmented Histamine Synthesis through p38 Activation. <i>Journal of Investigative Dermatology</i> , 2019, 139, 1516-1525.e3.	0.3	39
24	IL-33 and MRGPRX2-Triggered Activation of Human Skin Mast Cells—Elimination of Receptor Expression on Chronic Exposure, but Reinforced Degranulation on Acute Priming. <i>Cells</i> , 2019, 8, 341.	1.8	42
25	Allergic Fc γ R1 and pseudo-allergic MRGPRX2-triggered mast cell activation routes are independent and inversely regulated by SCF. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2018, 73, 256-260.	2.7	68
26	MRGPRX2 is negatively targeted by SCF and IL-4 to diminish pseudo-allergic stimulation of skin mast cells in culture. <i>Experimental Dermatology</i> , 2018, 27, 1298-1303.	1.4	29
27	The Impact on Allergy-Related Cells of a Birch Pollen Allergoid, with and without Monophosphoryl Lipid A, in Comparison with the Native Equivalent. <i>International Archives of Allergy and Immunology</i> , 2017, 172, 20-26.	0.9	10
28	An atlas of human long non-coding RNAs with accurate 5' ends. <i>Nature</i> , 2017, 543, 199-204.	13.7	898
29	An efficient method for gene knockdown by RNA interference in human skin mast cells. <i>Experimental Dermatology</i> , 2017, 26, 1136-1139.	1.4	21
30	Serum levels of 9 α ,11 β -PGF2 and apolipoprotein A1 achieve high predictive power as biomarkers of anaphylaxis. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2017, 72, 1801-1805.	2.7	12
31	Apoptotic resistance of human skin mast cells is mediated by Mcl-1. <i>Cell Death Discovery</i> , 2017, 3, 17048.	2.0	16
32	FANTOM5 CAGE profiles of human and mouse samples. <i>Scientific Data</i> , 2017, 4, 170112.	2.4	195
33	Skin mast cell phenotypes between two highly divergent cohorts are more pronounced variability within than between groups. <i>Experimental Dermatology</i> , 2017, 26, 446-449.	1.4	11
34	Retinoic Acid Negatively Impacts Proliferation and MCTC Specific Attributes of Human Skin Derived Mast Cells, but Reinforces Allergic Stimulability. <i>International Journal of Molecular Sciences</i> , 2017, 18, 525.	1.8	30
35	Phenotypic variability in human skin mast cells. <i>Experimental Dermatology</i> , 2016, 25, 434-439.	1.4	37
36	Bortezomib treatment diminishes hazelnut-induced intestinal anaphylaxis in mice. <i>European Journal of Immunology</i> , 2016, 46, 1727-1736.	1.6	8

#	ARTICLE	IF	CITATIONS
37	IL-4 and human skin mast cells revisited: reinforcement of a pro-allergic phenotype upon prolonged exposure. <i>Archives of Dermatological Research</i> , 2016, 308, 665-670.	1.1	22
38	Serum levels of $9\beta,11\beta$ -PGF 2 and cysteinyl leukotrienes are useful biomarkers of anaphylaxis. <i>Journal of Allergy and Clinical Immunology</i> , 2016, 137, 312-314.e7.	1.5	32
39	Ramipril and metoprolol intake aggravate human and murine anaphylaxis: Evidence for direct mast cell priming. <i>Journal of Allergy and Clinical Immunology</i> , 2015, 135, 491-499.	1.5	98
40	Retinoic acid potentiates inflammatory cytokines in human mast cells: Identification of mast cells as prominent constituents of the skin retinoid network. <i>Molecular and Cellular Endocrinology</i> , 2015, 406, 49-59.	1.6	25
41	Technical Advance: Transcription factor, promoter, and enhancer utilization in human myeloid cells. <i>Journal of Leukocyte Biology</i> , 2015, 97, 985-995.	1.5	23
42	Impact of sex on anaphylaxis severity—data from the Anaphylaxis Registry. <i>Journal of Allergy and Clinical Immunology</i> , 2015, 136, 1425-1426.	1.5	24
43	Thymic stromal lymphopoietin induction by skin irritation is independent of tumour necrosis factor- β , but supported by interleukin-1. <i>British Journal of Dermatology</i> , 2015, 172, 951-960.	1.4	22
44	Mast cell transcriptome elucidation: what are the implications for allergic disease in the clinic and where do we go next?. <i>Expert Review of Clinical Immunology</i> , 2014, 10, 977-980.	1.3	10
45	A promoter-level mammalian expression atlas. <i>Nature</i> , 2014, 507, 462-470.	13.7	1,838
46	Redefinition of the human mast cell transcriptome by deep-CAGE sequencing. <i>Blood</i> , 2014, 123, e58-e67.	0.6	175
47	Skin mast cells develop non-synchronized changes in typical lineage characteristics upon culture. <i>Experimental Dermatology</i> , 2014, 23, 933-935.	1.4	32
48	Integration of the Human Dermal Mast Cell into the Organotypic Co-culture Skin Model. <i>Methods in Molecular Biology</i> , 2014, 1192, 69-85.	0.4	2
49	Opposing effects on immune function and skin barrier regulation by the proteasome inhibitor bortezomib in an allergen-induced eczema model. <i>Experimental Dermatology</i> , 2013, 22, 742-747.	1.4	11
50	Causes and risk factors for anaphylaxis. <i>JDDG - Journal of the German Society of Dermatology</i> , 2013, 11, 44-50.	0.4	23
51	Auslöser und Risikofaktoren der Anaphylaxie. <i>JDDG - Journal of the German Society of Dermatology</i> , 2013, 11, 44-51.	0.4	24
52	Vitamin D Receptor Activation Improves Allergen-Triggered Eczema in Mice. <i>Journal of Investigative Dermatology</i> , 2012, 132, 330-336.	0.3	40
53	Testosterone exerts selective anti-inflammatory effects on human skin mast cells in a cell subset dependent manner. <i>Experimental Dermatology</i> , 2012, 21, 878-880.	1.4	19
54	Targeting the vitamin D receptor inhibits the B cell-dependent allergic immune response. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2011, 66, 540-548.	2.7	104

#	ARTICLE	IF	CITATIONS
55	Mast cell-derived TNF- α and histamine modify IL-6 and IL-8 expression and release from cutaneous tumor cells. <i>Experimental Dermatology</i> , 2011, 20, 1020-1022.	1.4	13
56	Long-Term Cultured Human Skin Mast Cells Are Suitable for Pharmacological Studies of Anti-Allergic Drugs Due to High Responsiveness to Fc μ RI Cross-Linking. <i>Bioscience, Biotechnology and Biochemistry</i> , 2011, 75, 382-384.	0.6	29
57	Mast cell lines HMC-1 and LAD2 in comparison with mature human skin mast cells “drastically reduced levels of tryptase and chymase in mast cell lines. <i>Experimental Dermatology</i> , 2010, 19, 845-847.	1.4	91
58	Tamoxifen counteracts the allergic immune response and improves allergen-induced dermatitis in mice. <i>Clinical and Experimental Allergy</i> , 2010, 40, 1256-1265.	1.4	22
59	Infection of in vivo differentiated human mast cells with hantaviruses. <i>Journal of General Virology</i> , 2010, 91, 1256-1261.	1.3	23
60	Interleukin-17A Promotes IgE Production in Human B Cells. <i>Journal of Investigative Dermatology</i> , 2010, 130, 2621-2628.	0.3	91
61	Baseline and stimulated turnover of cell surface c-Kit expression in different types of human mast cells. <i>Experimental Dermatology</i> , 2006, 15, 530-537.	1.4	17
62	Bivalent Effect of UV Light on Human Skin Mast Cells—Low-Level Mediator Release at Baseline but Potent Suppression Upon Mast Cell Triggering. <i>Journal of Investigative Dermatology</i> , 2005, 124, 453-456.	0.3	43
63	Evidence for a restricted rather than generalized stimulatory response of skin-derived human mast cells to substance P. <i>Journal of Neuroimmunology</i> , 2005, 163, 92-101.	1.1	58
64	The transcription factor profile of human mast cells in comparison with monocytes and granulocytes. <i>Cellular and Molecular Life Sciences</i> , 2005, 62, 214-226.	2.4	32
65	Comparative cytokine profile of human skin mast cells from two compartments—strong resemblance with monocytes at baseline but induction of IL-5 by IL-4 priming. <i>Journal of Leukocyte Biology</i> , 2004, 75, 244-252.	1.5	65
66	Human Skin Mast Cells Express H2 and H4, but not H3 Receptors. <i>Journal of Investigative Dermatology</i> , 2004, 123, 116-123.	0.3	123
67	Regulation of mast cell characteristics by cytokines: divergent effects of interleukin-4 on immature mast cell lines versus mature human skin mast cells. <i>Archives of Dermatological Research</i> , 2004, 296, 134-8.	1.1	25
68	All-trans retinoic acid down-regulates expression and function of β 2 integrins by human monocytes: opposite effects on monocytic cell lines. <i>European Journal of Immunology</i> , 2003, 33, 616-625.	1.6	12
69	Mast cells as initiators of immunity and host defense. <i>Experimental Dermatology</i> , 2001, 10, 1-10.	1.4	159
70	Retinoic acid up-regulates myeloid ICAM-3 expression and function in a cell-specific fashion—evidence for retinoid signaling pathways in the mast cell lineage. <i>Journal of Leukocyte Biology</i> , 2001, 69, 361-72.	1.5	13
71	Human Leukemic (HMC-1) Mast Cells Are Responsive to $1\alpha,25$ -Dihydroxyvitamin D3: Selective Promotion of ICAM-3 Expression and Constitutive Presence of Vitamin D3 Receptor. <i>Biochemical and Biophysical Research Communications</i> , 2000, 273, 1104-1110.	1.0	21
72	ICAM-3 (CD50) is expressed by human mast cells: Induction of homotypic mast cell aggregation via ICAM-3. <i>Cell Adhesion and Communication</i> , 1999, 7, 195-209.	1.7	18

#	ARTICLE	IF	CITATIONS
73	Signal Transduction via CD43 (Leukosialin, Sialophorin) and Associated Biological Effects in a Human Mast Cell Line (HMC-1). <i>Biochemical and Biophysical Research Communications</i> , 1998, 243, 163-169.	1.0	16
74	Leukosialin (CD43) is proteolytically cleaved from stimulated HMC-1 cells. <i>Immunobiology</i> , 1997, 197, 82-96.	0.8	12
75	Retinoic acids and dexamethasone alter cell-surface density of β_2 -integrins and ICAM-1 on human leukemic (HMC-1) mast cells. <i>Archives of Dermatological Research</i> , 1997, 289, 111-115.	1.1	12
76	Human Leukaemic (HMC-1) and Normal Skin Mast Cells Express β_2 -Integrins: Characterization of β_2 -Integrins and ICAM-1 on HMC-1 Cells. <i>Scandinavian Journal of Immunology</i> , 1997, 45, 471-481.	1.3	45
77	CD43 (leukosialin, sialophorin) expression is differentially regulated by retinoic acids. <i>European Journal of Immunology</i> , 1997, 27, 1147-1151.	1.6	19
78	A subclone (5C6) of the human mast cell line HMC-1 represents a more differentiated phenotype than the original cell line. <i>Archives of Dermatological Research</i> , 1996, 288, 778-782.	1.1	38
79	A subclone (5C6) of the human mast cell line HMC-1 represents a more differentiated phenotype than the original cell line. <i>Archives of Dermatological Research</i> , 1996, 288, 778-782.	1.1	6
80	An Allosteric Anti-Tryptase Antibody for the Treatment of Mast Cell-Mediated Severe Asthma. <i>SSRN Electronic Journal</i> , 0, , .	0.4	0
81	β_2 -arrestin-1 and β_2 -arrestin-2 Restrain MRGPRX2-Triggered Degranulation and ERK1/2 Activation in Human Skin Mast Cells. <i>Frontiers in Allergy</i> , 0, 3, .	1.2	11