Jan Hjort

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/4560136/jan-hjort-publications-by-year.pdf

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

72 2,414 30 46 g-index

83 2,995 5.3 5.31 ext. papers ext. citations avg, IF L-index

#	Paper	IF	Citations
72	Impacts of permafrost degradation on infrastructure. <i>Nature Reviews Earth & Environment</i> , 2022 , 3, 24-	·3 8 0.2	18
71	New high-resolution estimates of the permafrost thermal state and hydrothermal conditions over the Northern Hemisphere. <i>Earth System Science Data</i> , 2022 , 14, 865-884	10.5	4
70	Environmental Controls of InSAR-Based Periglacial Ground Dynamics in a Sub-Arctic Landscape. Journal of Geophysical Research F: Earth Surface, 2021 , 126, e2021JF006175	3.8	1
69	Current climate overrides historical effects on species richness and range size of freshwater plants in Europe and North America. <i>Journal of Ecology</i> , 2020 , 108, 1262-1275	6	11
68	Deriving a Frozen Area Fraction From Metop ASCAT Backscatter Based on Sentinel-1. <i>IEEE Transactions on Geoscience and Remote Sensing</i> , 2020 , 58, 6008-6019	8.1	2
67	High potential for loss of permafrost landforms in a changing climate. <i>Environmental Research Letters</i> , 2020 , 15, 104065	6.2	10
66	Changes in the functional features of macrophyte communities and driving factors across a 70-year period. <i>Hydrobiologia</i> , 2020 , 847, 3811-3827	2.4	9
65	Geodiversity-biodiversity relationship needs more empirical evidence. <i>Nature Ecology and Evolution</i> , 2020 , 4, 2-3	12.3	17
64	Does catchment geodiversity foster stream biodiversity?. <i>Landscape Ecology</i> , 2019 , 34, 2469-2485	4.3	18
63	New insights into the environmental factors controlling the ground thermal regime across the Northern Hemisphere: a comparison between permafrost and non-permafrost areas. <i>Cryosphere</i> , 2019 , 13, 693-707	5.5	21
62	Snow to Precipitation Ratio Controls Catchment Storage and Summer Flows in Boreal Headwater Catchments. <i>Water Resources Research</i> , 2019 , 55, 4096-4109	5.4	10
61	Mapping supply and demand of a provisioning ecosystem service across Europe. <i>Ecological Indicators</i> , 2019 , 103, 520-529	5.8	26
60	Landforms contribute to plant biodiversity at alpha, beta and gamma levels. <i>Journal of Biogeography</i> , 2019 , 46, 1699-1710	4.1	21
59	Opinion: To advance sustainable stewardship, we must document not only biodiversity but geodiversity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2019 , 116, 16155-16158	11.5	48
58	Is catchment geodiversity a useful surrogate of aquatic plant species richness?. <i>Journal of Biogeography</i> , 2019 , 46, 1711-1722	4.1	17
57	Accessibility analysis in evaluating exposure risk to an ecosystem disservice. <i>Applied Geography</i> , 2019 , 113, 102098	4.4	2
56	Correlates of different facets and components of beta diversity in stream organisms. <i>Oecologia</i> , 2019 , 191, 919-929	2.9	20

(2017-2019)

55	Circumpolar permafrost maps and geohazard indices for near-future infrastructure risk assessments. <i>Scientific Data</i> , 2019 , 6, 190037	8.2	31
54	The role of geodiversity in providing ecosystem services at broad scales. <i>Ecological Indicators</i> , 2018 , 91, 47-56	5.8	36
53	Different species trait groups of stream diatoms show divergent responses to spatial and environmental factors in a subarctic drainage basin. <i>Hydrobiologia</i> , 2018 , 816, 213-230	2.4	11
52	Predicting occupancy and abundance by niche position, niche breadth and body size in stream organisms. <i>Oecologia</i> , 2018 , 186, 205-216	2.9	23
51	Highly variable species distribution models in a subarctic stream metacommunity: Patterns, mechanisms and implications. <i>Freshwater Biology</i> , 2018 , 63, 33-47	3.1	7
50	The added value of geodiversity indices in explaining variation of stream macroinvertebrate diversity. <i>Ecological Indicators</i> , 2018 , 94, 420-429	5.8	14
49	Statistical Forecasting of Current and Future Circum-Arctic Ground Temperatures and Active Layer Thickness. <i>Geophysical Research Letters</i> , 2018 , 45, 4889-4898	4.9	48
48	Degrading permafrost puts Arctic infrastructure at risk by mid-century. <i>Nature Communications</i> , 2018 , 9, 5147	17.4	181
47	Local environment and space drive multiple facets of stream macroinvertebrate beta diversity. <i>Journal of Biogeography</i> , 2018 , 45, 2744-2754	4.1	55
46	Global variation in the beta diversity of lake macrophytes is driven by environmental heterogeneity rather than latitude. <i>Journal of Biogeography</i> , 2017 , 44, 1758-1769	4.1	82
45	Modelling native and alien vascular plant species richness: At which scales is geodiversity most relevant?. <i>Global Ecology and Biogeography</i> , 2017 , 26, 763-776	6.1	52
44	Integrating dispersal proxies in ecological and environmental research in the freshwater realm. <i>Environmental Reviews</i> , 2017 , 25, 334-349	4.5	55
43	Species richness and taxonomic distinctness of lake macrophytes along environmental gradients in two continents. <i>Freshwater Biology</i> , 2017 , 62, 1194-1206	3.1	13
42	Spatial relationship between biodiversity and geodiversity across a gradient of land-use intensity in high-latitude landscapes. <i>Landscape Ecology</i> , 2017 , 32, 1049-1063	4.3	25
41	Average niche breadths of species in lake macrophyte communities respond to ecological gradients variably in four regions on two continents. <i>Oecologia</i> , 2017 , 184, 219-235	2.9	12
40	Spatio-temporal aspects of the environmental factors affecting water quality in boreal rivers. <i>Environmental Earth Sciences</i> , 2017 , 76, 1	2.9	7
39	Urbanity as a determinant of exposure to grass pollen in Helsinki Metropolitan area, Finland. <i>PLoS ONE</i> , 2017 , 12, e0186348	3.7	13
38	Combining geodiversity with climate and topography to account for threatened species richness. <i>Conservation Biology</i> , 2017 , 31, 364-375	6	49

37	Geography of global change and species richness in the North. Environmental Reviews, 2017, 25, 184-19	2 4.5	21
36	Permafrost Map for Norway, Sweden and Finland. Permafrost and Periglacial Processes, 2017, 28, 359-37	78 1.2	64
35	Hierarchical decomposition of trait patterns of macroinvertebrate communities in subarctic streams. <i>Freshwater Science</i> , 2016 , 35, 1032-1048	2	19
34	Introducing accessibility analysis in mapping cultural ecosystem services. <i>Ecological Indicators</i> , 2016 , 66, 416-427	5.8	56
33	Importance of spatial scale in structuring emergent lake vegetation across environmental gradients and scales: GIS-based approach. <i>Ecological Indicators</i> , 2016 , 60, 1164-1172	5.8	6
32	Fine-Scale Exposure to Allergenic Pollen in the Urban Environment: Evaluation of Land Use Regression Approach. <i>Environmental Health Perspectives</i> , 2016 , 124, 619-26	8.4	35
31	Extreme urban-rural temperatures in the coastal city of Turku, Finland: Quantification and visualization based on a generalized additive model. <i>Science of the Total Environment</i> , 2016 , 569-570, 507-517	10.2	20
30	Why geodiversity matters in valuing nature& stage. Conservation Biology, 2015, 29, 630-9	6	123
29	A review of selection-based tests of abiotic surrogates for species representation. <i>Conservation Biology</i> , 2015 , 29, 668-79	6	32
28	Geomorphological factors predict water quality in boreal rivers. <i>Earth Surface Processes and Landforms</i> , 2015 , 40, 1989-1999	3.7	31
27	Inferring the effects of potential dispersal routes on the metacommunity structure of stream insects: as the crow flies, as the fish swims or as the fox runs?. <i>Journal of Animal Ecology</i> , 2015 , 84, 1342	- \$ 3	58
26	Which Environmental Factors Determine Recent Cryoturbation and Solifluction Activity in a Subarctic Landscape? A Comparison between Active and Inactive Features. <i>Permafrost and Periglacial Processes</i> , 2014 , 25, 136-143	4.2	11
25	Transferability of geomorphological distribution models: Evaluation using solifluction features in subarctic and Arctic regions. <i>Geomorphology</i> , 2014 , 204, 165-176	4.3	13
24	Land use impacts on trace metal concentrations of suburban stream sediments in the Helsinki region, Finland. <i>Science of the Total Environment</i> , 2013 , 456-457, 222-30	10.2	37
23	2.6 Statistical Methods for Geomorphic Distribution Modeling 2013 , 59-73		19
22	Inclusion of explicit measures of geodiversity improve biodiversity models in a boreal landscape. <i>Biodiversity and Conservation</i> , 2012 , 21, 3487-3506	3.4	65
21	Can geodiversity be predicted from space?. <i>Geomorphology</i> , 2012 , 153-154, 74-80	4.3	32
20	Effects of scale on modelling the urban heat island in Turku, SW Finland. <i>Climate Research</i> , 2012 , 55, 10:	511618	16

(2004-2011)

19	Spatial prediction of urbanflural temperatures using statistical methods. <i>Theoretical and Applied Climatology</i> , 2011 , 106, 139-152	3	25
18	Novel theoretical insights into geomorphic processenvironment relationships using simulated response curves. <i>Earth Surface Processes and Landforms</i> , 2011 , 36, 363-371	3.7	17
17	Recent vegetation changes at the high-latitude tree line ecotone are controlled by geomorphological disturbance, productivity and diversity. <i>Global Ecology and Biogeography</i> , 2010 , 19, 810-821	6.1	101
16	Geodiversity of high-latitude landscapes in northern Finland. <i>Geomorphology</i> , 2010 , 115, 109-116	4.3	81
15	The thermal state of permafrost in the nordic area during the international polar year 2007 2009. <i>Permafrost and Periglacial Processes</i> , 2010 , 21, 156-181	4.2	210
14	Effects of scale and data source in periglacial distribution modelling in a high arctic environment, western Svalbard. <i>Permafrost and Periglacial Processes</i> , 2010 , 21, 345-354	4.2	10
13	Assessing spatial uncertainty in predictive geomorphological mapping: A multi-modelling approach. <i>Computers and Geosciences</i> , 2010 , 36, 355-361	4.5	14
12	Statistical consensus methods for improving predictive geomorphology maps. <i>Computers and Geosciences</i> , 2009 , 35, 615-625	4.5	36
11	Periglacial distribution modelling with a boosting method. <i>Permafrost and Periglacial Processes</i> , 2009 , 20, 15-25	4.2	11
10	Interaction of geomorphic and ecologic features across altitudinal zones in a subarctic landscape. <i>Geomorphology</i> , 2009 , 112, 324-333	4.3	41
9	Effects of sample size on the accuracy of geomorphological models. <i>Geomorphology</i> , 2008 , 102, 341-350	04.3	35
8	Downscaling of coarse-grained geomorphological data. <i>Earth Surface Processes and Landforms</i> , 2008 , 33, 75-89	3.7	17
7	Can abundance of geomorphological features be predicted using presence besence data?. <i>Earth Surface Processes and Landforms</i> , 2008 , 33, 741-750	3.7	6
6	A comparison of predictive methods in modelling the distribution of periglacial landforms in Finnish Lapland. <i>Earth Surface Processes and Landforms</i> , 2008 , 33, 2241-2254	3.7	41
5	Landscape scale determinants of periglacial features in subarctic Finland: a grid-based modelling approach. <i>Permafrost and Periglacial Processes</i> , 2007 , 18, 115-127	4.2	31
4	Scale matters multi-resolution study of the determinants of patterned ground activity in subarctic Finland. <i>Geomorphology</i> , 2006 , 80, 282-294	4.3	30
3	Modelling patterned ground distribution in finnish lapland: an integration of topographical, ground and remote sensing information. <i>Geografiska Annaler, Series A: Physical Geography</i> , 2006 , 88, 19-29	1.1	33
2	Generalized linear modelling in periglacial studies: terrain parameters and patterned ground. Permafrost and Periglacial Processes, 2004, 15, 327-338	4.2	38

New high-resolution estimates of the permafrost thermal state and hydrothermal conditions over the Northern Hemisphere

6