
Manu Soto

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4569976/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|--|------------|----------------|
| 1 | Impacts of sewage sludges deposition on agricultural soils: Effects upon model soil organisms. Comparative Biochemistry and Physiology Part - C: Toxicology and Pharmacology, 2022, 255, 109276. | 2.6 | 2 |
| 2 | Infection Rate in Seabasses Fed with Viscera Parasitised by Anisakid Larvae. Acta Parasitologica, 2022, 67, 835-841. | 1.1 | 3 |
| 3 | Variability and distribution of parasites, pathologies and their effect on wild mussels (Mytilus sp) in different environments along a wide latitudinal span in the Northern Atlantic and Arctic Oceans. Marine Environmental Research, 2022, 176, 105585. | 2.5 | 5 |
| 4 | Predicting environmental concentrations and the potential risk of Plant Protection Products (PPP) on non-target soil organisms accounting for regional and landscape ecological variability in european soils. Chemosphere, 2022, 303, 135045. | 8.2 | 5 |
| 5 | Reviewing <i>Pseudoloma neurophilia</i> infections in the popular zebrafish model. Reviews in Aquaculture, 2021, 13, 1816-1827. | 9.0 | 6 |
| 6 | Application of in situ bioremediation strategies in soils amended with sewage sludges. Science of the Total Environment, 2021, 766, 144099. | 8.0 | 22 |
| 7 | Sea Bass Primary Cultures versus RTgill-W1 Cell Line: Influence of Cell Model on the Sensitivity to Nanoparticles. Nanomaterials, 2021, 11, 3136. | 4.1 | 3 |
| 8 | Organotropism and biomarker response in oyster Crassostrea gigas exposed to platinum in seawater. Environmental Science and Pollution Research, 2020, 27, 3584-3599. | 5.3 | 6 |
| 9 | Gentle remediation options for soil with mixed chromium (VI) and lindane pollution: biostimulation, bioaugmentation, phytoremediation and vermiremediation. Heliyon, 2020, 6, e04550. | 3.2 | 37 |
| 10 | Effects of elevated temperatures and cadmium exposure on stress biomarkers at different biological complexity levels in Eisenia fetida earthworms. Comparative Biochemistry and Physiology Part - C: Toxicology and Pharmacology, 2020, 231, 108735. | 2.6 | 5 |
| 11 | Morphological Characterization and Hydrodynamic Behavior of Shortfin Mako Shark (Isurus) Tj ETQq1 1 0.7843 | 14 rgBT /(| Overlock 10 Ti |
| 12 | Influence of season-depending ecological variables on biomarker baseline levels in mussels (Mytilus) Tj ETQq0 0 | 0 rgBT /0 | verlock 10 Tf |
| 13 | Selection of an optimal culture medium and the most responsive viability assay to assess AgNPs toxicity with primary cultures of Eisenia fetida coelomocytes. Ecotoxicology and Environmental Safety, 2019, 183, 109545. | 6.0 | 14 |
| 14 | The EU Horizon 2020 project GRACE: integrated oil spill response actions and environmental effects. Environmental Sciences Europe, 2019, 31, . | 5.5 | 23 |
| 15 | Evaluation of the effectiveness of a bioremediation process in experimental soils polluted with chromium and lindane. Ecotoxicology and Environmental Safety, 2019, 181, 255-263. | 6.0 | 32 |
| 16 | Enhanced discrimination of basophilic cells on mussel digestive gland tissue sections by means of toluidine-eosin staining. Journal of Invertebrate Pathology, 2019, 161, 29-39. | 3.2 | 6 |
| 17 | Bioaccumulation, tissue and cell distribution, biomarkers and toxicopathic effects of CdS quantum dots in mussels, Mytilus galloprovincialis. Ecotoxicology and Environmental Safety, 2019, 167, 288-300. | 6.0 | 18 |
| 18 | Nanoparticles Under the Spotlight. , 2019, , 169-188. | | 1 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Changes in metallothionein transcription levels in the mussel Mytilus galloprovincialis exposed to CdTe quantum dots. Ecotoxicology, 2018, 27, 402-410. | 2.4 | 13 |
| 20 | Chemical contamination assessment in mangrove-lined Caribbean coastal systems using the oyster Crassostrea rhizophorae as biomonitor species. Environmental Science and Pollution Research, 2018, 25, 13396-13415. | 5.3 | 28 |
| 21 | Tracing platinum accumulation kinetics in oyster Crassostrea gigas, a sentinel species in coastal marine environments. Science of the Total Environment, 2018, 615, 652-663. | 8.0 | 15 |
| 22 | Digestive cell lysosomes as main targets for Ag accumulation and toxicity in marine mussels, <i>Mytilus galloprovincialis</i> , exposed to maltose-stabilised Ag nanoparticles of different sizes. Nanotoxicology, 2017, 11, 168-183. | 3.0 | 38 |
| 23 | Lysosomal responses to different gold forms (nanoparticles, aqueous, bulk) in mussel digestive cells: a trade-off between the toxicity of the capping agent and form, size and exposure concentration. Nanotoxicology, 2017, 11, 658-670. | 3.0 | 9 |
| 24 | Integrative assessment of the effects produced by Ag nanoparticles at different levels of biological complexity in Eisenia fetida maintained in two standard soils (OECD and LUFA 2.3). Chemosphere, 2017, 181, 747-758. | 8.2 | 33 |
| 25 | Assessment of health status of oysters (Crassostrea gigas) exposed to environmentally relevant concentrations of Ag and Cu in brackish waters. Journal of Sea Research, 2017, 130, 229-238. | 1.6 | 6 |
| 26 | Responses to silver nanoparticles and silver nitrate in a battery of biomarkers measured in coelomocytes and in target tissues of Eisenia fetida earthworms. Ecotoxicology and Environmental Safety, 2017, 141, 57-63. | 6.0 | 43 |
| 27 | Successive Onset of Molecular, Cellular and Tissue-Specific Responses in Midgut Gland of Littorina littorea Exposed to Sub-Lethal Cadmium Concentrations. International Journal of Molecular Sciences, 2017, 18, 1815. | 4.1 | 25 |
| 28 | Assessing the variability of telomere length measures by means of Telomeric Restriction Fragments (TRF) in different tissues of cod Gadus morhua. Gene Reports, 2016, 5, 117-125. | 0.8 | 2 |
| 29 | Assessment of the effects of Cu and Ag in oysters Crassostrea gigas (Thunberg, 1793) using a battery of cell and tissue level biomarkers. Marine Environmental Research, 2016, 122, 11-22. | 2.5 | 20 |
| 30 | Uptake route and resulting toxicity of silver nanoparticles in Eisenia fetida earthworm exposed through Standard OECD Tests. Ecotoxicology, 2016, 25, 1543-1555. | 2.4 | 44 |
| 31 | Nanoparticle size and combined toxicity of TiO ₂ and DSLS (surfactant) contribute to lysosomal responses in digestive cells of mussels exposed to TiO ₂ nanoparticles. Nanotoxicology, 2016, 10, 1168-1176. | 3.0 | 43 |
| 32 | Silver and copper bioaccumulation kinetics in oyster Crassostrea gigas tissues at environmentally relevant exposure levels using stable isotope spikes. Estuarine, Coastal and Shelf Science, 2016, 179, 135-144. | 2.1 | 10 |
| 33 | Dynamic Quality Index for agricultural soils based on fuzzy logic. Ecological Indicators, 2016, 60, 678-692. | 6.3 | 28 |
| 34 | Biological characterization of the skin of shortfin mako shark <i>Isurus oxyrinchus</i> and preliminary study of the hydrodynamic behaviour through computational fluid dynamics. Journal of Fish Biology, 2015, 87, 123-137. | 1.6 | 33 |
| 35 | Application of ecological risk assessment based on a novel TRIAD-tiered approach to contaminated soil surrounding a closed non-sealed landfill. Science of the Total Environment, 2015, 514, 49-59. | 8.0 | 20 |
| 36 | Establishment of toxicity thresholds in subpopulations of coelomocytes (amoebocytes vs. eleocytes) of Eisenia fetida exposed in vitro to a variety of metals: implications for biomarker measurements. Ecotoxicology, 2015, 24, 1004-1013. | 2.4 | 36 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | STEM-in-SEM high resolution imaging of gold nanoparticles and bivalve tissues in bioaccumulation experiments. Analyst, The, 2015, 140, 3082-3089. | 3.5 | 24 |
| 38 | Combining chemical and biological endpoints, a major challenge for twenty-first century's environmental specimen banks. Environmental Science and Pollution Research, 2015, 22, 1631-1634. | 5.3 | 5 |
| 39 | Effects of Soil Organic Matter Content on Cadmium Toxicity in Eisenia Fetida: Implications for the Use of Biomarkers and Standard Toxicity Tests. Archives of Environmental Contamination and Toxicology, 2015, 68, 181-192. | 4.1 | 36 |
| 40 | Zonation in the digestive tract of Eisenia fetida: Implications in biomarker measurements for toxicity assessment. Comparative Biochemistry and Physiology Part - C: Toxicology and Pharmacology, 2014, 160, 42-53. | 2.6 | 5 |
| 41 | Lysosomal and tissue-level biomarkers in mussels cross-transplanted among four estuaries with different pollution levels. Science of the Total Environment, 2014, 472, 36-48. | 8.0 | 27 |
| 42 | Optimization of NRU assay in primary cultures of Eisenia fetida for metal toxicity assessment. Ecotoxicology, 2014, 23, 1326-1335. | 2.4 | 20 |
| 43 | Toxicity assessment through multiple endpoint bioassays in soils posing environmental risk according to regulatory screening values. Environmental Science and Pollution Research, 2014, 21, 9689-9708. | 5.3 | 22 |
| 44 | Molecular cloning and measurement of telomerase reverse transcriptase (TERT) transcription patterns in tissues of European hake (Merluccius merluccius) and Atlantic cod (Gadus morhua) during aging. Gene, 2014, 541, 8-18. | 2.2 | 10 |
| 45 | Health status assessment through an integrative biomarker approach in mussels of different ages with a different history of exposure to the Prestige oil spill. Science of the Total Environment, 2014, 493, 65-78. | 8.0 | 24 |
| 46 | Combined use of native and caged mussels to assess biological effects of pollution through the integrative biomarker approach. Aquatic Toxicology, 2013, 136-137, 32-48. | 4.0 | 97 |
| 47 | Behaviour of Au-citrate nanoparticles in seawater and accumulation in bivalves at environmentally relevant concentrations. Environmental Pollution, 2013, 174, 134-141. | 7.5 | 79 |
| 48 | Towards an integrative soil health assessment strategy: A three tier (integrative biomarker response) approach with Eisenia fetida applied to soils subjected to chronic metal pollution. Science of the Total Environment, 2013, 442, 344-365. | 8.0 | 44 |
| 49 | Marine ecosystem health status assessment through integrative biomarker indices: a comparative study after the Prestige oil spill "Mussel Watch― Ecotoxicology, 2013, 22, 486-505. | 2.4 | 135 |
| 50 | Silver accumulation in oysters from the Gironde Estuary (France): Distribution between different organs and histopathological alterations based on microscopical observations. Comparative Biochemistry and Physiology Part A, Molecular & Integrative Physiology, 2012, 163, S20-S21. | 1.8 | 1 |
| 51 | Metal pollution assessment in different seasons of the year in the Oka river estuary using cell and tissue level biomarkers in oysters. Comparative Biochemistry and Physiology Part A, Molecular & Integrative Physiology, 2012, 163, S21. | 1.8 | 0 |
| 52 | Integrated coastal monitoring of a gas processing plant using native and caged mussels. Science of the Total Environment, 2012, 426, 375-386. | 8.0 | 26 |
| 53 | Application of a battery of biomarkers in mussel digestive gland to assess long-term effects of the Prestige oil spill in Galicia and Bay of Biscay: Tissue-level biomarkers and histopathology. Journal of Environmental Monitoring, 2011, 13, 915. | 2.1 | 79 |
| 54 | Application of a battery of biomarkers in mussel digestive gland to assess long-term effects of the Prestige oil spill in Galicia and Bay of Biscay: Correlation and multivariate analysis. Journal of Environmental Monitoring, 2011, 13, 933. | 2.1 | 34 |

Μανύ δοτο

| # | Article | IF | CITATIONS |
|----|--|------------|---------------|
| 55 | Occurrence and Distribution of Metals in Mussels from the Cantabrian Coast. Archives of Environmental Contamination and Toxicology, 2010, 59, 235-243. | 4.1 | 38 |
| 56 | Seasonality in cell and tissue-level biomarkers in Mytilus galloprovincialis: relevance for long-term pollution monitoring. Aquatic Biology, 2010, 9, 203-219. | 1.4 | 37 |
| 57 | Neutral red uptake assay with coelomocytes of Eisenia fetida as an effective and reliable screening method for soil health assessment. Comparative Biochemistry and Physiology Part A, Molecular & Integrative Physiology, 2009, 153, S88. | 1.8 | Ο |
| 58 | Assessment of biological effects of environmental pollution along the NW Mediterranean Sea using red mullets as sentinel organisms. Environmental Pollution, 2008, 153, 157-168. | 7.5 | 49 |
| 59 | Changes in cell-type composition in digestive gland of slugs and its influence in biomarkers following transplantation between a relatively unpolluted and a chronically metal-polluted site. Environmental Pollution, 2008, 156, 367-379. | 7.5 | 32 |
| 60 | Metallothionein expression and Neutral Red uptake as biomarkers of metal exposure and effect in Eisenia fetida and Lumbricus terrestris exposed to Cd. European Journal of Soil Biology, 2007, 43, S233-S238. | 3.2 | 51 |
| 61 | Assessment of biological effects of environmental pollution along the NW Mediterranean Sea using mussels as sentinel organisms. Environmental Pollution, 2007, 148, 236-250. | 7.5 | 142 |
| 62 | Digestive cell turnover in digestive gland epithelium of slugs experimentally exposed to a mixture of cadmium and kerosene. Chemosphere, 2007, 70, 144-154. | 8.2 | 30 |
| 63 | Tissue- and cell-specific expression of metallothionein genes in cadmium- and copper-exposed mussels analyzed by in situ hybridization and RT–PCR. Toxicology and Applied Pharmacology, 2007, 220, 186-196. | 2.8 | 66 |
| 64 | Immunolocalization of metallothioneins in different tissues of turbot (Scophthalmus maximus) exposed to Cd. Histology and Histopathology, 2007, 22, 719-28. | 0.7 | 10 |
| 65 | Quantitative changes in metallothionein expression in target cell-types in the gills of turbot (Scophthalmus maximus) exposed to Cd, Cu, Zn and after a depuration treatment. Aquatic Toxicology, 2006, 77, 64-77. | 4.0 | 45 |
| 66 | Biomarkers in mussels from a copper site gradient (Visnes, Norway): An integrated biochemical, histochemical and histological study. Aquatic Toxicology, 2006, 78, S109-S116. | 4.0 | 105 |
| 67 | Bioavailability and cellular effects of metals on Lumbricus terrestris inhabiting volcanic soils. Environmental Pollution, 2006, 142, 103-108. | 7.5 | 31 |
| 68 | Signs of recovery of mussels health two years after the Prestige oil spill. Marine Environmental Research, 2006, 62, S337-S341. | 2.5 | 44 |
| 69 | Freshwater molluscs from volcanic areas as model organisms to assess adaptation to metal chronic pollution. Science of the Total Environment, 2006, 371, 168-175. | 8.0 | 15 |
| 70 | Cell and tissue biomarkers in mussel, and histopathology in hake and anchovy from Bay of Biscay after the Prestige oil spill (Monitoring Campaign 2003). Marine Pollution Bulletin, 2006, 53, 287-304. | 5.0 | 125 |
| 71 | Application of two SH-based methods for metallothionein determination in mussels and intercalibration of the spectrophotometric method: laboratory and field studies in the Mediterranean Sea. Biomarkers, 2005, 10, 342-359. | 1.9 | 31 |
| 72 | Cellular biomarkers of exposure and biological effect in hepatocytes of turbot (Scophthalmus) Tj ETQq0 0 0 rgB | T /Overloc | k 10 Tf 50 62 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Biomonitoring of environmental pollution along the Basque coast, using molecular, cellular and tissue-level biomarkers: an integrative approach. Elsevier Oceanography Series, 2004, 70, 335-364. | 0.1 | 7 |
| 74 | Autometallography and metallothionein immunohistochemistry in hepatocytes of turbot (Scophthalmus maximusL.) after exposure to cadmium and depuration treatment. Biomarkers, 2002, 7, 491-500. | 1.9 | 24 |
| 75 | Introduction: Studies on metal localization in animal cells. Microscopy Research and Technique, 2002, 56, 315-317. | 2.2 | 0 |
| 76 | Cellular and subcellular distribution of metals in molluscs. Microscopy Research and Technique, 2002, 56, 358-392. | 2.2 | 401 |
| 77 | Subcellular distribution of cadmium and its cellular ligands in mussel digestive gland cells as revealed by combined autometallography and X-ray microprobe analysis. The Histochemical Journal, 2002, 34, 273-280. | 0.6 | 54 |
| 78 | Chemical, biochemical and cellular responses in the digestive gland of the musselMytilus galloprovincialisfrom the Spanish Mediterranean coast. Biomarkers, 2001, 6, 335-350. | 1.9 | 61 |
| 79 | Biomarkers and trace metals in the digestive gland of indigenous and transplanted mussels, Mytilus galloprovincialis, in Venice Lagoon, Italy. Marine Environmental Research, 2000, 50, 417-423. | 2.5 | 58 |
| 80 | Changes in mussel biometry on exposure to metals: implications in estimation of metal bioavailability in â€Mussel-Watch' programmes. Science of the Total Environment, 2000, 247, 175-187. | 8.0 | 70 |
| 81 | Autometallographical localisation of Cu and Zn within target cell compartments of winkles following exposure to Cu&Zn mixtures. European Journal of Histochemistry, 1999, 43, 323-34. | 1.5 | 9 |
| 82 | Autometallographed metal content in digestive cells of winkles: a cost-effective screening tool for monitoring Cu and Zn pollution. Aquatic Toxicology, 1998, 40, 123-140. | 4.0 | 11 |
| 83 | Autometallographical Procedure for the Localization of Metal Traces in Molluscan Tissues by Light Microscopy. Journal of Histotechnology, 1998, 21, 123-127. | 0.5 | 31 |
| 84 | Autometallographical Procedure for the Localization of Metal Traces in Molluscan Tissues by Light Microscopy. Journal of Histotechnology, 1998, 21, 123-127. | 0.5 | 2 |
| 85 | BSD Extent, an Index for Metal Pollution Screening Based on the Metal Content within Digestive Cell Lysosomes of Mussels as Determined by Autometallography. Ecotoxicology and Environmental Safety, 1997, 37, 141-151. | 6.0 | 22 |
| 86 | The contribution of metal/shell-weight index in target-tissues to metal body burden in sentinel marine molluscs. 2. Mytilus galloprovincialis. Science of the Total Environment, 1997, 198, 149-160. | 8.0 | 28 |
| 87 | The contribution of metal/shell-weight index in target-tissues to metal body burden in sentinel marine molluscs. 1. Littorina littorea. Science of the Total Environment, 1997, 198, 135-147. | 8.0 | 29 |
| 88 | Metal bioavailability assessment in 'mussel-watch' programmes by automated image analysis of autometallographical black silver deposits (BSD) in digestive cell lysosomes. Marine Ecology - Progress Series, 1997, 156, 141-150. | 1.9 | 38 |
| 89 | Tissue and cell distribution of copper, zinc and cadmium in the mussel, Mytilus galloprovincialis, determined by autometallography. Tissue and Cell, 1996, 28, 557-568. | 2.2 | 81 |
| 90 | Autometallographic localization of protein-bound copper and zinc in the common winkle,Littorina littorea: A light microscopical study. The Histochemical Journal, 1996, 28, 689-701. | 0.6 | 29 |

Manu Soto

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 91 | Tissue-level biomarkers and biological effect of mercury on sentinel slugs, Arion ater. Archives of Environmental Contamination and Toxicology, 1996, 31, 54-62. | 4.1 | 47 |
| 92 | Tissue-Level Biomarkers and Biological Effect of Mercury on Sentinel Slugs, Arion ater. Archives of Environmental Contamination and Toxicology, 1996, 31, 54-62. | 4.1 | 2 |
| 93 | Bioavailable heavy metals in estuarine waters as assessed by metal/shell-weight indices in sentinel mussels Mytilus galloprovincialis. Marine Ecology - Progress Series, 1995, 125, 127-136. | 1.9 | 68 |
| 94 | Seasonal variability in the quantitative structure of the digestive tubules ofLittorina littorea. Aquatic Living Resources, 1992, 5, 299-305. | 1.2 | 12 |
| 95 | A PLANIMETRIC STUDY OF MORPHOLOGICAL VARIABILITY IN THE DIGESTIVE DIVERTICULA OF LITTORINA LITTOREA (LINNAEUS) AND MYTILUS EDULIS LINNAEUS. Journal of Molluscan Studies, 1990, 56, 339-344. | 1.2 | 2 |
| 96 | Histochemistry and elemental composition of the stomach cells in Littorina littorea (L.). Folia Histochemica Et Cytobiologica, 1990, 28, 239-48. | 1.5 | 5 |
| 97 | Lurzoru kutsatuen karakterizazio intentsiboa in vivo eta in silico fokatzeak erabiliz. Ekaia (journal), 0, , . | 0.0 | 0 |
| 98 | Araztegi lokatzak jasotako lurzoruaren analisi toxikologikoa zizare eta landareak erabiliz. Ekaia (journal), 0, , . | 0.0 | 0 |