## Erkki Isometsä

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/457575/publications.pdf

Version: 2024-02-01

156 papers 8,477 citations

50170 46 h-index 49773 87 g-index

164 all docs 164 docs citations

164 times ranked 9267 citing authors

#	Article	IF	Citations
1	Lifetime Prevalence of Psychotic and Bipolar I Disorders in a General Population. Archives of General Psychiatry, 2007, 64, 19.	13.8	1,112
2	DSM-IV mood-, anxiety- and alcohol use disorders and their comorbidity in the Finnish general population. Social Psychiatry and Psychiatric Epidemiology, 2005, 40, 1-10.	1.6	318
3	Suicidal Ideation and Attempts Among Psychiatric Patients With Major Depressive Disorder. Journal of Clinical Psychiatry, 2003, 64, 1094-1100.	1.1	282
4	International Society for Bipolar Disorders Task Force on Suicide: metaâ€analyses and metaâ€regression of correlates of suicide attempts and suicide deaths in bipolar disorder. Bipolar Disorders, 2015, 17, 1-16.	1.1	265
5	The relationship of neuroticism and extraversion to symptoms of anxiety and depression in the general population. Depression and Anxiety, 2006, 23, 281-289.	2.0	260
6	Suicide attempts preceding completed suicide. British Journal of Psychiatry, 1998, 173, 531-535.	1.7	243
7	Suicidal Behaviour in Mood Disorders—Who, When, and Why?. Canadian Journal of Psychiatry, 2014, 59, 120-130.	0.9	231
8	Suicidal Ideation and Attempts in Bipolar I and II Disorders. Journal of Clinical Psychiatry, 2005, 66, 1456-1462.	1.1	204
9	Current Comorbidity of Psychiatric Disorders Among DSM-IV Major Depressive Disorder Patients in Psychiatric Care in the Vantaa Depression Study. Journal of Clinical Psychiatry, 2002, 63, 126-134.	1.1	202
10	Incidence and Predictors of Suicide Attempts in DSM–IV Major Depressive Disorder: A Five-Year Prospective Study. American Journal of Psychiatry, 2010, 167, 801-808.	4.0	195
11	Prospective study of risk factors for attempted suicide among patients with DSM–IV major depressive disorder. British Journal of Psychiatry, 2005, 186, 314-318.	1.7	161
12	Major depressive disorder and white matter abnormalities: A diffusion tensor imaging study with tract-based spatial statistics. Journal of Affective Disorders, 2010, 120, 240-244.	2.0	160
13	Differences in incidence of suicide attempts between bipolar I and II disorders and major depressive disorder. Bipolar Disorders, 2014, 16, 652-661.	1.1	141
14	Severity and Comorbidity Predict Episode Duration and Recurrence of DSM-IV Major Depressive Disorder. Journal of Clinical Psychiatry, 2004, 65, 810-819.	1.1	135
15	The 12-month prevalence and risk factors for major depressive episode in Finland: representative sample of 5993 adults. Acta Psychiatrica Scandinavica, 2000, 102, 178-184.	2.2	134
16	The clinical characteristics of DSM-IV bipolar I and II disorders: baseline findings from the Jorvi Bipolar Study (JoBS). Bipolar Disorders, 2004, 6, 395-405.	1.1	130
17	Childhood adversities as risk factors for adult mental disorders. Social Psychiatry and Psychiatric Epidemiology, 2005, 40, 769-777.	1.6	130
18	Epidemiology, neurobiology and pharmacological interventions related to suicide deaths and suicide attempts in bipolar disorder: Part I of a report of the International Society for Bipolar Disorders Task Force on Suicide in Bipolar Disorder. Australian and New Zealand Journal of Psychiatry, 2015, 49, 785-802.	1.3	122

#	Article	IF	CITATIONS
19	Prospective study of risk factors for attempted suicide among patients with bipolar disorder. Bipolar Disorders, 2006, 8, 576-585.	1.1	112
20	Differences in incidence of suicide attempts during phases of bipolar I and II disorders. Bipolar Disorders, 2008, 10, 588-596.	1.1	107
21	Early age at onset of bipolar disorder is associated with more severe clinical features but delayed treatment seeking. Bipolar Disorders, 2007, 9, 698-705.	1.1	103
22	Long-Term Outcome of Major Depressive Disorder in Psychiatric Patients Is Variable. Journal of Clinical Psychiatry, 2008, 69, 196-205.	1.1	95
23	Suicidal behaviour during different phases of bipolar disorder. Journal of Affective Disorders, 2007, 97, 101-107.	2.0	93
24	Continuity Is the Main Challenge in Treating Major Depressive Disorder in Psychiatric Care. Journal of Clinical Psychiatry, 2005, 66, 220-227.	1.1	90
25	A review of factors associated with greater likelihood of suicide attempts and suicide deaths in bipolar disorder: Part II of a report of the International Society for Bipolar Disorders Task Force on Suicide in Bipolar Disorder. Australian and New Zealand Journal of Psychiatry, 2015, 49, 1006-1020.	1.3	87
26	Differences in Axis I and II Comorbidity Between Bipolar I and II Disorders and Major Depressive Disorder. Journal of Clinical Psychiatry, 2006, 67, 584-593.	1.1	85
27	The Mood Disorder Questionnaire improves recognition of bipolar disorder in psychiatric care. BMC Psychiatry, 2003, 3, 8.	1.1	84
28	Depressive disorders in primary care: recurrent, chronic, and co-morbid. Psychological Medicine, 2005, 35, 673-682.	2.7	84
29	<i>P2RX7</i> gene is associated consistently with mood disorders and predicts clinical outcome in three clinical cohorts. American Journal of Medical Genetics Part B: Neuropsychiatric Genetics, 2011, 156, 435-447.	1.1	79
30	Differences in outcome of DSMâ€N bipolar I and II disorders. Bipolar Disorders, 2008, 10, 413-425.	1.1	75
31	Machine learning and big data analytics in bipolar disorder: A position paper from the International Society for Bipolar Disorders Big Data Task Force. Bipolar Disorders, 2019, 21, 582-594.	1.1	74
32	The Effect of Personality on Daily Life Emotional Processes. PLoS ONE, 2014, 9, e110907.	1.1	71
33	Antidepressant utilisation patterns and determinants of short-term and non-psychiatric use in the Finnish general adult population. Journal of Affective Disorders, 2008, 110, 94-105.	2.0	70
34	Differences and similarities of risk factors for suicidal ideation and attempts among patients with depressive or bipolar disorders. Journal of Affective Disorders, 2016, 193, 318-330.	2.0	70
35	Differences in Recent Life Events between Alcoholic and Depressive Nonalcoholic Suicides. Alcoholism: Clinical and Experimental Research, 1994, 18, 1143-1149.	1.4	62
36	Functional and Work Disability in Major Depressive Disorder. Journal of Nervous and Mental Disease, 2005, 193, 189-195.	0.5	62

#	Article	IF	CITATIONS
37	Use of health services for major depressive episode in Finland. Journal of Affective Disorders, 2004, 79, 105-112.	2.0	60
38	Neuroticism, introversion, and major depressive disorder-traits, states, or scars?. Depression and Anxiety, 2009, 26, 325-334.	2.0	60
39	Treatment of major depressive disorder in the Finnish general population. Depression and Anxiety, 2009, 26, 1049-1059.	2.0	59
40	Decline in suicidal ideation among patients with MDD is preceded by decline in depression and hopelessness. Journal of Affective Disorders, 2006, 95, 95-102.	2.0	58
41	Cigarette smoking, alcohol intoxication and major depressive episode in a representative population sample. Journal of Epidemiology and Community Health, 2001, 55, 573-576.	2.0	57
42	Co-morbidity and stability of melancholic features in DSM-IV major depressive disorder. Psychological Medicine, 2004, 34, 1443-1452.	2.7	57
43	Gender differences in bipolar disorder type I and II. Acta Psychiatrica Scandinavica, 2009, 120, 464-473.	2.2	51
44	Long-term personality changes and predictive adaptive responses after depressive episodes. Evolution and Human Behavior, 2015, 36, 337-344.	1.4	51
45	Short-term escitalopram treatment normalizes aberrant self-referential processing in major depressive disorder. Journal of Affective Disorders, 2018, 236, 222-229.	2.0	50
46	Predictors for Switch From Unipolar Major Depressive Disorder to Bipolar Disorder Type I or II. Journal of Clinical Psychiatry, 2008, 69, 1267-1275.	1.1	49
47	A prospective latent analyses study of psychiatric comorbidity of DSMâ€N bipolar I and II disorders. Bipolar Disorders, 2010, 12, 271-284.	1.1	48
48	Temperament and character traits predict future burden of depression. Journal of Affective Disorders, 2014, 158, 139-147.	2.0	46
49	Differences in neuroticism and extraversion between patients with bipolar I or II and general population subjects or major depressive disorder patients. Journal of Affective Disorders, 2010, 125, 42-52.	2.0	45
50	Determinants of Functional Disability and Social Adjustment in Major Depressive Disorder. Journal of Nervous and Mental Disease, 2006, 194, 570-576.	0.5	44
51	Relationships of neuroticism and extraversion with axis I and II comorbidity among patients with DSM-IV major depressive disorder. Journal of Affective Disorders, 2009, 114, 110-121.	2.0	44
52	Incidence and predictors of suicide attempts among primary-care patients with depressive disorders: a 5-year prospective study. Psychological Medicine, 2014, 44, 291-302.	2.7	44
53	Do Means Matter?. Journal of Nervous and Mental Disease, 2003, 191, 745-750.	0.5	43
54	Relationship between sunlight and the age of onset of bipolar disorder: An international multisite study. Journal of Affective Disorders, 2014, 167, 104-111.	2.0	43

#	Article	IF	CITATIONS
55	Incidence and predictors of suicide attempts in bipolar I and II disorders: A 5â€year followâ€up study. Bipolar Disorders, 2017, 19, 13-22.	1.1	43
56	Predictors of adherence to psychopharmacological and psychosocial treatment in bipolar I or II disorders – an 18-month prospective study. Journal of Affective Disorders, 2014, 155, 110-117.	2.0	42
57	Fiveâ€year outcome of bipolar I and II disorders: findings of the Jorvi Bipolar Study. Bipolar Disorders, 2015, 17, 363-374.	1.1	42
58	Influence of light exposure during early life on the age of onset of bipolar disorder. Journal of Psychiatric Research, 2015, 64, 1-8.	1.5	39
59	Prevalence of current anxiety disorders in people with bipolar disorder during euthymia: a meta-analysis. Psychological Medicine, 2017, 47, 1107-1115.	2.7	39
60	Relationships between self-reported childhood traumatic experiences, attachment style, neuroticism and features of borderline personality disorders in patients with mood disorders. Journal of Affective Disorders, 2017, 210, 82-89.	2.0	38
61	Treatment attitudes and adherence of psychiatric patients with major depressive disorder: A five-year prospective study. Journal of Affective Disorders, 2010, 127, 102-112.	2.0	37
62	White matter hyperintensities and cognitive performance in adult patients with bipolar I, bipolar II, and major depressive disorders. European Psychiatry, 2014, 29, 226-232.	0.1	37
63	Life Events, Social Support, and Onset of Major Depressive Episode in Finnish Patients. Journal of Nervous and Mental Disease, 2004, 192, 373-381.	0.5	36
64	Screening of psychiatric outpatients for borderline personality disorder with the McLean Screening Instrument for Borderline Personality Disorder (MSI-BPD). Nordic Journal of Psychiatry, 2009, 63, 475-479.	0.7	36
65	Completed suicide and recent lithium treatment. Journal of Affective Disorders, 1992, 26, 101-103.	2.0	35
66	Adequacy of Treatment Received by Diagnosed and Undiagnosed Patients With Bipolar I and II Disorders. Journal of Clinical Psychiatry, 2007, 68, 102-110.	1.1	35
67	Postâ€discharge suicides of inpatients with bipolar disorder in Finland. Bipolar Disorders, 2014, 16, 867-874.	1.1	34
68	Risk factors for suicide in depression in Finland: firstâ€hospitalized patients followed up to 24Âyears. Acta Psychiatrica Scandinavica, 2019, 139, 154-163.	2.2	33
69	Differences in temperament and character dimensions in patients with bipolar I or II or major depressive disorder and general population subjects. Psychological Medicine, 2011, 41, 1579-1591.	2.7	32
70	Do characteristics of patients with major depressive disorder differ between primary and psychiatric care?. Psychological Medicine, 2007, 37, 893-904.	2.7	31
71	Only half of bipolar I and II patients report prodromal symptoms. Journal of Affective Disorders, 2008, 111, 366-371.	2.0	31
72	Disability Pension for Major Depression in Finland. American Journal of Psychiatry, 2000, 157, 1869-1872.	4.0	30

#	Article	IF	CITATIONS
73	Health-related quality of life 6 months after burns among hospitalized patients: Predictive importance of mental disorders and burn severity. Burns, 2015, 41, 742-748.	1.1	30
74	Anxiety symptoms in a major mood and schizophrenia spectrum disorders. European Psychiatry, 2016, 37, 1-7.	0.1	30
75	Predictors of longâ€ŧerm work disability among patients with type I and <scp>II</scp> bipolar disorder: a prospective 18â€month followâ€up study. Bipolar Disorders, 2015, 17, 821-835.	1.1	29
76	Temperament, Character and Suicide Attempts in Unipolar and Bipolar Mood Disorders. Journal of Clinical Psychiatry, 2016, 77, 252-260.	1.1	29
77	DEPRESSION AND SMOKING: A 5-YEAR PROSPECTIVE STUDY OF PATIENTS WITH MAJOR DEPRESSIVE DISORDER. Depression and Anxiety, 2013, 30, 580-588.	2.0	28
78	Predominant polarity in bipolar I and II disorders: A five-year follow-up study. Journal of Affective Disorders, 2019, 246, 806-813.	2.0	28
79	Categorical and Dimensional Stability of Comorbid Personality Disorder Symptoms in <i>DSM-IV</i> Major Depressive Disorder. Journal of Clinical Psychiatry, 2010, 71, 287-295.	1.1	26
80	Association between solar insolation and a history of suicide attempts in bipolar I disorder. Journal of Psychiatric Research, 2019, 113, 1-9.	1.5	25
81	Borderline Personality Disorder With Depression Confers Significant Risk of Suicidal Behavior in Mood Disorder Patientsâ€"A Comparative Study. Frontiers in Psychiatry, 2020, 11, 290.	1.3	25
82	Data Collection for Mental Health Studies Through Digital Platforms: Requirements and Design of a Prototype. JMIR Research Protocols, 2017, 6, e110.	0.5	25
83	Borderline personality disorder among primary care depressive patients: A five-year study. Journal of Affective Disorders, 2014, 155, 303-306.	2.0	23
84	Pharmacological and psychosocial treatment of depression in primary care: Low intensity and poor adherence and continuity. Journal of Affective Disorders, 2016, 202, 145-152.	2.0	23
85	How suicidal bipolar patients are depends on how suicidal ideation is defined. Journal of Affective Disorders, 2009, 118, 48-54.	2.0	22
86	A 5-year prospective study of predictors for functional and work disability among primary care patients with depressive disorders. European Psychiatry, 2015, 30, 51-57.	0.1	22
87	Temperament, character and personality disorders. European Psychiatry, 2013, 28, 483-491.	0.1	21
88	Return to work six months after burn: A prospective study at the Helsinki Burn Center. Burns, 2015, 41, 1152-1160.	1.1	21
89	Personality disorders and suicide attempts in unipolar and bipolar mood disorders. Journal of Affective Disorders, 2016, 190, 632-639.	2.0	21
90	Level of functioning, perceived work ability, and work status among psychiatric patients with major mental disorders. European Psychiatry, 2017, 44, 83-89.	0.1	21

#	Article	IF	Citations
91	Alcohol use and smoking in burn patients at the Helsinki Burn Center. Burns, 2018, 44, 158-167.	1.1	21
92	Clinical predictors of unrecognized bipolar I and II disorders. Bipolar Disorders, 2008, 10, 238-244.	1.1	20
93	Reduced visual contrast suppression during major depressive episodes. Journal of Psychiatry and Neuroscience, 2021, 46, E222-E231.	1.4	20
94	Role of Hopelessness in Suicidal Ideation Among Patients With Depressive Disorders. Journal of Clinical Psychiatry, 2020, 81, .	1.1	20
95	State and trait hopelessness in a prospective five-year study of patients with depressive disorders. Journal of Affective Disorders, 2018, 239, 107-114.	2.0	19
96	Employment Status Influences the Weekly Patterns of Suicide among Alcohol Misusers. Alcoholism: Clinical and Experimental Research, 1997, 21, 1704-1706.	1.4	18
97	Health-related quality of life of primary care patients with depressive disorders. European Psychiatry, 2016, 37, 28-34.	0.1	18
98	Obstetric and perinatal health outcomes related to schizophrenia: A national register-based follow-up study among Finnish women born between 1965 and 1980 and their offspring. European Psychiatry, 2018, 52, 68-75.	0.1	18
99	Differences and overlap in self-reported symptoms of bipolar disorder and borderline personality disorder. European Psychiatry, 2015, 30, 914-919.	0.1	17
100	Hopelessness across phases of bipolar I or II disorder: A prospective study. Journal of Affective Disorders, 2009, 115, 11-17.	2.0	15
101	Typology of clinical course in bipolar disorder based on 18-month naturalistic follow-up. Psychological Medicine, 2013, 43, 789-799.	2.7	15
102	Diagnostic conversion from unipolar depression to bipolar disorder, schizophrenia, or schizoaffective disorder: A nationwide prospective 15â€year register study on 43Â495 inpatients. Bipolar Disorders, 2020, 22, 582-592.	1.1	15
103	NEUROTICISM MEDIATES THE EFFECT OF P2RX7 ON OUTCOMES OF MOOD DISORDERS. Depression and Anxiety, 2012, 29, 816-823.	2.0	14
104	Mediating role of borderline personality disorder traits in the effects of childhood maltreatment on suicidal behaviour among mood disorder patients. European Psychiatry, 2017, 44, 53-60.	0.1	14
105	Decline in suicide mortality after psychiatric hospitalization for depression in Finland between 1991 and 2014. World Psychiatry, 2018, 17, 110-112.	4.8	14
106	Schizophrenia and induced abortions: A national register-based follow-up study among Finnish women born between 1965–1980 with schizophrenia or schizoaffective disorder. Schizophrenia Research, 2018, 192, 142-147.	1.1	14
107	Maintenance pharmacotherapy for recurrent major depressive disorder: 5-year follow-up study. British Journal of Psychiatry, 2008, 193, 163-164.	1.7	13
108	Do antidepressants change personality?—A five-year observational study. Journal of Affective Disorders, 2012, 142, 200-207.	2.0	13

#	Article	IF	Citations
109	A single dose of mirtazapine attenuates neural responses to self-referential processing. Journal of Psychopharmacology, 2016, 30, 23-32.	2.0	13
110	Schizophrenia and pregnancy: a national register-based follow-up study among Finnish women born between 1965 and 1980. Archives of Women's Mental Health, 2020, 23, 91-100.	1.2	13
111	Suicides in Mood Disorders in Psychiatric Settings in Nordic National Register–Based Studies. Frontiers in Psychiatry, 2020, 11, 721.	1.3	13
112	Maintenance treatment received by patients with bipolar I and II disorders â€" A naturalistic prospective study. Journal of Affective Disorders, 2010, 121, 116-126.	2.0	12
113	Self-reported symptoms of schizotypal and borderline personality disorder in patients with mood disorders. European Psychiatry, 2016, 33, 37-44.	0.1	12
114	Maternal schizophrenia and out-of-home placements of offspring: A national follow-up study among Finnish women born 1965–1980 and their children. Psychiatry Research, 2019, 273, 9-14.	1.7	12
115	Comorbid alcohol use disorder in psychiatric MDD patients: A five-year prospective study. Journal of Affective Disorders, 2020, 267, 283-288.	2.0	12
116	Family history of psychiatric disorders and the outcome of psychiatric patients with DSM-IV major depressive disorder. Journal of Affective Disorders, 2011, 131, 251-259.	2.0	11
117	Presence and Overlap of Bipolar Symptoms and Borderline Features during Major Depressive Episodes. Journal of Affective Disorders, 2021, 280, 467-477.	2.0	11
118	Validity and reliability of the Finnish version of the Functioning Assessment Short Test (FAST) in bipolar disorder. International Journal of Bipolar Disorders, 2015, 3, 10.	0.8	9
119	Escitalopram enhances synchrony of brain responses during emotional narratives in patients with major depressive disorder. Neurolmage, 2021, 237, 118110.	2.1	9
120	Variants in regulatory elements of PDE4D associate with major mental illness in the Finnish population. Molecular Psychiatry, 2021, 26, 816-824.	4.1	8
121	Concomitants of Family Histories of Mood Disorders and Alcoholism in a Clinical Cohort of Patients With Bipolar I and II Disorder. Journal of Nervous and Mental Disease, 2012, 200, 388-394.	0.5	7
122	Single dose of mirtazapine modulates whole-brain functional connectivity during emotional narrative processing. Psychiatry Research - Neuroimaging, 2017, 263, 61-69.	0.9	7
123	Psychoactive substance use in specialized psychiatric care patients. International Journal of Psychiatry in Medicine, 2017, 52, 399-415.	0.8	7
124	Self-reported psychosis-like experiences in patients with mood disorders. European Psychiatry, 2018, 51, 90-97.	0.1	7
125	Self-reported treatment adherence among psychiatric in- and outpatients. Nordic Journal of Psychiatry, 2018, 72, 526-533.	0.7	7
126	Clinical course predicts long-term outcomes in bipolar disorder. Psychological Medicine, 2019, 49, 1109-1117.	2.7	6

#	Article	IF	CITATIONS
127	Variations in seasonal solar insolation are associated with a history of suicide attempts in bipolar I disorder. International Journal of Bipolar Disorders, 2021, 9, 26.	0.8	6
128	Implementation of CYP2D6 copy-number imputation panel and frequency of key pharmacogenetic variants in Finnish individuals with a psychotic disorder. Pharmacogenomics Journal, 2022, 22, 166-172.	0.9	6
129	Response style and severity and chronicity of depressive disorders in primary health care. European Psychiatry, 2016, 33, 1-8.	0.1	5
130	Do Suicide Attempts of Mood Disorder Patients Directly Increase the Risk for a Reattempt?. Frontiers in Psychiatry, 2020, 11, 547791.	1.3	5
131	A Randomized Clinical Trial of Attempted Suicide Short Intervention Program versus Crisis Counseling in Preventing Repeat Suicide Attempts: A Two-Year Follow-Up Study. Psychotherapy and Psychosomatics, 2022, 91, 190-199.	4.0	5
132	Long-term work disability due to type I and II bipolar disorder: findings of a six-year prospective study. International Journal of Bipolar Disorders, 2022, 10, .	0.8	5
133	The Relationship Between Smoking and Suicidal Behavior in Psychiatric Patients with Major Depressive Disorder. Archives of Suicide Research, 2019, 23, 590-604.	1.2	4
134	Effectiveness of brief schema group therapy for borderline personality disorder symptoms: a randomized pilot study. Nordic Journal of Psychiatry, 2021, 75, 176-185.	0.7	4
135	A prospective latent analysis study of Axis I psychiatric co-morbidity of DSM-IV major depressive disorder. Psychological Medicine, 2014, 44, 949-959.	2.7	3
136	Maintenance Pharmacotherapy for Recurrent Major Depressive Disorder in Primary Care: A 5-year Follow-up Study. European Psychiatry, 2017, 41, 111-114.	0.1	3
137	Features of borderline personality disorder as a mediator of the relation between childhood traumatic experiences and psychosis-like experiences in patients with mood disorder. European Psychiatry, 2018, 49, 9-15.	0.1	3
138	Reaction Time and Visual Memory in Connection with Alcohol Use in Schizophrenia and Schizoaffective Disorder. Brain Sciences, 2021, 11, 688.	1.1	3
139	Pregnancy, delivery and postpartum in women with schizophrenia or schizoaffective disorder in Finland: a national register-based comparative study. Psychiatry Research, 2020, 294, 113504.	1.7	3
140	Mood Zoom could be a promising tool for daily mood variability monitoring, potentially differentiating bipolar from borderline patients. Evidence-Based Mental Health, 2017, 20, e9-e9.	2.2	2
141	Sleep in Psychotic Disorders: Results From Nationwide SUPER Finland Study. Schizophrenia Bulletin Open, 2022, 3, .	0.9	2
142	Rumination And Primary Care Depression. European Psychiatry, 2016, 33, S422-S422.	0.1	1
143	White Matter Hyperintensities after Five-Year Follow-Up and a Cross-Sectional FA Decrease in Bipolar I and Major Depressive Patients. Neuropsychobiology, 2022, 81, 39-50.	0.9	1
144	Reaction Time and Visual Memory in Connection to Alcohol Use in Persons with Bipolar Disorder. Brain Sciences, 2021, 11, 1154.	1.1	1

#	Article	IF	CITATIONS
145	SNP Variants at $16p13.11$ Clarify the Role of the NDE1/miR-484 Locus in Major Mental Illness in Finland. Schizophrenia Bulletin Open, 2020, 1, .	0.9	1
146	A randomized clinical trial of the ASSIP vs. crisis counselling in preventing suicide attempt repetition: a two-year follow-up study. Nordic Journal of Psychiatry, 2022, , 1-1.	0.7	1
147	Differences in psychosocial functioning between psychotic disorders in the Finnish SUPER study. Schizophrenia Research, 2022, 244, 10-17.	1.1	1
148	Author's response. Brain Stimulation, 2014, 7, 338.	0.7	0
149	Large-scale suicide prevention by pharmacological treatment of mood disorders. European Psychiatry, 2016, 33, S53-S53.	0.1	0
150	Dangerously mixed. Bipolar Disorders, 2017, 19, 314-315.	1.1	0
151	Incidence and Predictors of Suicide Attempts in Bipolar I and II Disorders: A Five-year Follow-up. European Psychiatry, 2017, 41, S87-S87.	0.1	0
152	O5.5. SLEEP IN MAJOR PSYCHIATRIC DISORDERS: RESULTS FROM NATIONWIDE SUPER FINLAND STUDY. Schizophrenia Bulletin, 2018, 44, S88-S88.	2.3	0
153	T49. SOCIODEMOGRAPHIC AND LIFESTYLE FACTORS ASSOCIATED WITH GOOD PERFORMANCE IN PAIRED ASSOCIATES LEARNING TEST IN PATIENTS WITH SCHIZOPHRENIA. Schizophrenia Bulletin, 2020, 46, \$250-\$250.	2.3	0
154	The evidence does not support the premises of the environmental mismatch hypothesis. Response to â∈œBipolar disorder: An evolutionary psychoneuroimmunological approach― Neuroscience and Biobehavioral Reviews, 2021, 131, 539-540.	2.9	0
155	Reaction Time and Visual Memory in Connection to Hazardous Drinking Polygenic Scores in Schizophrenia, Schizoaffective Disorder and Bipolar Disorder. Brain Sciences, 2021, 11, 1422.	1.1	0
156	SYMPOSIUM 1: suicide prevention: Nordic Perspectives. Nordic Journal of Psychiatry, 2022, , 1-1.	0.7	0