

# Erkki Isometsä

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/457575/publications.pdf>

Version: 2024-02-01

156  
papers

8,477  
citations

50170

46  
h-index

49773

87  
g-index

164  
all docs

164  
docs citations

164  
times ranked

9267  
citing authors

#	ARTICLE	IF	CITATIONS
1	White Matter Hyperintensities after Five-Year Follow-Up and a Cross-Sectional FA Decrease in Bipolar I and Major Depressive Patients. <i>Neuropsychobiology</i> , 2022, 81, 39-50.	0.9	1
2	A Randomized Clinical Trial of Attempted Suicide Short Intervention Program versus Crisis Counseling in Preventing Repeat Suicide Attempts: A Two-Year Follow-Up Study. <i>Psychotherapy and Psychosomatics</i> , 2022, 91, 190-199.	4.0	5
3	Sleep in Psychotic Disorders: Results From Nationwide SUPER Finland Study. <i>Schizophrenia Bulletin Open</i> , 2022, 3, .	0.9	2
4	Implementation of CYP2D6 copy-number imputation panel and frequency of key pharmacogenetic variants in Finnish individuals with a psychotic disorder. <i>Pharmacogenomics Journal</i> , 2022, 22, 166-172.	0.9	6
5	A randomized clinical trial of the ASSIP vs. crisis counselling in preventing suicide attempt repetition: a two-year follow-up study. <i>Nordic Journal of Psychiatry</i> , 2022, , 1-1.	0.7	1
6	SYMPOSIUM 1: suicide prevention: Nordic Perspectives. <i>Nordic Journal of Psychiatry</i> , 2022, , 1-1.	0.7	0
7	Differences in psychosocial functioning between psychotic disorders in the Finnish SUPER study. <i>Schizophrenia Research</i> , 2022, 244, 10-17.	1.1	1
8	Long-term work disability due to type I and II bipolar disorder: findings of a six-year prospective study. <i>International Journal of Bipolar Disorders</i> , 2022, 10, .	0.8	5
9	Variants in regulatory elements of PDE4D associate with major mental illness in the Finnish population. <i>Molecular Psychiatry</i> , 2021, 26, 816-824.	4.1	8
10	Presence and Overlap of Bipolar Symptoms and Borderline Features during Major Depressive Episodes. <i>Journal of Affective Disorders</i> , 2021, 280, 467-477.	2.0	11
11	Effectiveness of brief schema group therapy for borderline personality disorder symptoms: a randomized pilot study. <i>Nordic Journal of Psychiatry</i> , 2021, 75, 176-185.	0.7	4
12	Reduced visual contrast suppression during major depressive episodes. <i>Journal of Psychiatry and Neuroscience</i> , 2021, 46, E222-E231.	1.4	20
13	Reaction Time and Visual Memory in Connection with Alcohol Use in Schizophrenia and Schizoaffective Disorder. <i>Brain Sciences</i> , 2021, 11, 688.	1.1	3
14	Reaction Time and Visual Memory in Connection to Alcohol Use in Persons with Bipolar Disorder. <i>Brain Sciences</i> , 2021, 11, 1154.	1.1	1
15	Escitalopram enhances synchrony of brain responses during emotional narratives in patients with major depressive disorder. <i>NeuroImage</i> , 2021, 237, 118110.	2.1	9
16	The evidence does not support the premises of the environmental mismatch hypothesis. Response to "Bipolar disorder: An evolutionary psychoneuroimmunological approach". <i>Neuroscience and Biobehavioral Reviews</i> , 2021, 131, 539-540.	2.9	0
17	Variations in seasonal solar insolation are associated with a history of suicide attempts in bipolar I disorder. <i>International Journal of Bipolar Disorders</i> , 2021, 9, 26.	0.8	6
18	Reaction Time and Visual Memory in Connection to Hazardous Drinking Polygenic Scores in Schizophrenia, Schizoaffective Disorder and Bipolar Disorder. <i>Brain Sciences</i> , 2021, 11, 1422.	1.1	0

#	ARTICLE	IF	CITATIONS
19	Schizophrenia and pregnancy: a national register-based follow-up study among Finnish women born between 1965 and 1980. Archives of Women's Mental Health, 2020, 23, 91-100.	1.2	13
20	T49. SOCIODEMOGRAPHIC AND LIFESTYLE FACTORS ASSOCIATED WITH GOOD PERFORMANCE IN PAIRED ASSOCIATES LEARNING TEST IN PATIENTS WITH SCHIZOPHRENIA. Schizophrenia Bulletin, 2020, 46, S250-S250.	2.3	0
21	Suicides in Mood Disorders in Psychiatric Settings in Nordic National Register-Based Studies. Frontiers in Psychiatry, 2020, 11, 721.	1.3	13
22	Do Suicide Attempts of Mood Disorder Patients Directly Increase the Risk for a Reattempt?. Frontiers in Psychiatry, 2020, 11, 547791.	1.3	5
23	Diagnostic conversion from unipolar depression to bipolar disorder, schizophrenia, or schizoaffective disorder: A nationwide prospective 15-year register study on 43,495 inpatients. Bipolar Disorders, 2020, 22, 582-592.	1.1	15
24	Comorbid alcohol use disorder in psychiatric MDD patients: A five-year prospective study. Journal of Affective Disorders, 2020, 267, 283-288.	2.0	12
25	Borderline Personality Disorder With Depression Confers Significant Risk of Suicidal Behavior in Mood Disorder Patients—A Comparative Study. Frontiers in Psychiatry, 2020, 11, 290.	1.3	25
26	Pregnancy, delivery and postpartum in women with schizophrenia or schizoaffective disorder in Finland: a national register-based comparative study. Psychiatry Research, 2020, 294, 113504.	1.7	3
27	Role of Hopelessness in Suicidal Ideation Among Patients With Depressive Disorders. Journal of Clinical Psychiatry, 2020, 81, .	1.1	20
28	SNP Variants at 16p13.11 Clarify the Role of the NDE1/miR-484 Locus in Major Mental Illness in Finland. Schizophrenia Bulletin Open, 2020, 1, .	0.9	1
29	The Relationship Between Smoking and Suicidal Behavior in Psychiatric Patients with Major Depressive Disorder. Archives of Suicide Research, 2019, 23, 590-604.	1.2	4
30	Clinical course predicts long-term outcomes in bipolar disorder. Psychological Medicine, 2019, 49, 1109-1117.	2.7	6
31	Machine learning and big data analytics in bipolar disorder: A position paper from the International Society for Bipolar Disorders Big Data Task Force. Bipolar Disorders, 2019, 21, 582-594.	1.1	74
32	Association between solar insolation and a history of suicide attempts in bipolar I disorder. Journal of Psychiatric Research, 2019, 113, 1-9.	1.5	25
33	Risk factors for suicide in depression in Finland: first-hospitalized patients followed up to 24 years. Acta Psychiatrica Scandinavica, 2019, 139, 154-163.	2.2	33
34	Predominant polarity in bipolar I and II disorders: A five-year follow-up study. Journal of Affective Disorders, 2019, 246, 806-813.	2.0	28
35	Maternal schizophrenia and out-of-home placements of offspring: A national follow-up study among Finnish women born 1965–1980 and their children. Psychiatry Research, 2019, 273, 9-14.	1.7	12
36	Short-term escitalopram treatment normalizes aberrant self-referential processing in major depressive disorder. Journal of Affective Disorders, 2018, 236, 222-229.	2.0	50

#	ARTICLE	IF	CITATIONS
37	O5.5. SLEEP IN MAJOR PSYCHIATRIC DISORDERS: RESULTS FROM NATIONWIDE SUPER FINLAND STUDY. Schizophrenia Bulletin, 2018, 44, S88-S88.	2.3	0
38	Features of borderline personality disorder as a mediator of the relation between childhood traumatic experiences and psychosis-like experiences in patients with mood disorder. European Psychiatry, 2018, 49, 9-15.	0.1	3
39	Decline in suicide mortality after psychiatric hospitalization for depression in Finland between 1991 and 2014. World Psychiatry, 2018, 17, 110-112.	4.8	14
40	Self-reported psychosis-like experiences in patients with mood disorders. European Psychiatry, 2018, 51, 90-97.	0.1	7
41	Schizophrenia and induced abortions: A national register-based follow-up study among Finnish women born between 1965 and 1980 with schizophrenia or schizoaffective disorder. Schizophrenia Research, 2018, 192, 142-147.	1.1	14
42	Alcohol use and smoking in burn patients at the Helsinki Burn Center. Burns, 2018, 44, 158-167.	1.1	21
43	Self-reported treatment adherence among psychiatric in- and outpatients. Nordic Journal of Psychiatry, 2018, 72, 526-533.	0.7	7
44	State and trait hopelessness in a prospective five-year study of patients with depressive disorders. Journal of Affective Disorders, 2018, 239, 107-114.	2.0	19
45	Obstetric and perinatal health outcomes related to schizophrenia: A national register-based follow-up study among Finnish women born between 1965 and 1980 and their offspring. European Psychiatry, 2018, 52, 68-75.	0.1	18
46	Maintenance Pharmacotherapy for Recurrent Major Depressive Disorder in Primary Care: A 5-year Follow-up Study. European Psychiatry, 2017, 41, 111-114.	0.1	3
47	Incidence and predictors of suicide attempts in bipolar I and II disorders: A 5-year follow-up study. Bipolar Disorders, 2017, 19, 13-22.	1.1	43
48	Level of functioning, perceived work ability, and work status among psychiatric patients with major mental disorders. European Psychiatry, 2017, 44, 83-89.	0.1	21
49	Mood Zoom could be a promising tool for daily mood variability monitoring, potentially differentiating bipolar from borderline patients. Evidence-Based Mental Health, 2017, 20, e9-e9.	2.2	2
50	Dangerously mixed. Bipolar Disorders, 2017, 19, 314-315.	1.1	0
51	Mediating role of borderline personality disorder traits in the effects of childhood maltreatment on suicidal behaviour among mood disorder patients. European Psychiatry, 2017, 44, 53-60.	0.1	14
52	Single dose of mirtazapine modulates whole-brain functional connectivity during emotional narrative processing. Psychiatry Research - Neuroimaging, 2017, 263, 61-69.	0.9	7
53	Prevalence of current anxiety disorders in people with bipolar disorder during euthymia: a meta-analysis. Psychological Medicine, 2017, 47, 1107-1115.	2.7	39
54	Relationships between self-reported childhood traumatic experiences, attachment style, neuroticism and features of borderline personality disorders in patients with mood disorders. Journal of Affective Disorders, 2017, 210, 82-89.	2.0	38

#	ARTICLE	IF	CITATIONS
55	Incidence and Predictors of Suicide Attempts in Bipolar I and II Disorders: A Five-year Follow-up. <i>European Psychiatry</i> , 2017, 41, S87-S87.	0.1	0
56	Psychoactive substance use in specialized psychiatric care patients. <i>International Journal of Psychiatry in Medicine</i> , 2017, 52, 399-415.	0.8	7
57	Data Collection for Mental Health Studies Through Digital Platforms: Requirements and Design of a Prototype. <i>JMIR Research Protocols</i> , 2017, 6, e110.	0.5	25
58	Rumination And Primary Care Depression. <i>European Psychiatry</i> , 2016, 33, S422-S422.	0.1	1
59	Anxiety symptoms in a major mood and schizophrenia spectrum disorders. <i>European Psychiatry</i> , 2016, 37, 1-7.	0.1	30
60	Health-related quality of life of primary care patients with depressive disorders. <i>European Psychiatry</i> , 2016, 37, 28-34.	0.1	18
61	Pharmacological and psychosocial treatment of depression in primary care: Low intensity and poor adherence and continuity. <i>Journal of Affective Disorders</i> , 2016, 202, 145-152.	2.0	23
62	Large-scale suicide prevention by pharmacological treatment of mood disorders. <i>European Psychiatry</i> , 2016, 33, S53-S53.	0.1	0
63	Differences and similarities of risk factors for suicidal ideation and attempts among patients with depressive or bipolar disorders. <i>Journal of Affective Disorders</i> , 2016, 193, 318-330.	2.0	70
64	A single dose of mirtazapine attenuates neural responses to self-referential processing. <i>Journal of Psychopharmacology</i> , 2016, 30, 23-32.	2.0	13
65	Self-reported symptoms of schizotypal and borderline personality disorder in patients with mood disorders. <i>European Psychiatry</i> , 2016, 33, 37-44.	0.1	12
66	Response style and severity and chronicity of depressive disorders in primary health care. <i>European Psychiatry</i> , 2016, 33, 1-8.	0.1	5
67	Personality disorders and suicide attempts in unipolar and bipolar mood disorders. <i>Journal of Affective Disorders</i> , 2016, 190, 632-639.	2.0	21
68	Temperament, Character and Suicide Attempts in Unipolar and Bipolar Mood Disorders. <i>Journal of Clinical Psychiatry</i> , 2016, 77, 252-260.	1.1	29
69	Predictors of long-term work disability among patients with type I and <sc>II</sc> bipolar disorder: a prospective 18-month follow-up study. <i>Bipolar Disorders</i> , 2015, 17, 821-835.	1.1	29
70	Five-year outcome of bipolar I and II disorders: findings of the Jorvi Bipolar Study. <i>Bipolar Disorders</i> , 2015, 17, 363-374.	1.1	42
71	Health-related quality of life 6 months after burns among hospitalized patients: Predictive importance of mental disorders and burn severity. <i>Burns</i> , 2015, 41, 742-748.	1.1	30
72	Validity and reliability of the Finnish version of the Functioning Assessment Short Test (FAST) in bipolar disorder. <i>International Journal of Bipolar Disorders</i> , 2015, 3, 10.	0.8	9

#	ARTICLE	IF	CITATIONS
73	Long-term personality changes and predictive adaptive responses after depressive episodes. <i>Evolution and Human Behavior</i> , 2015, 36, 337-344.	1.4	51
74	A review of factors associated with greater likelihood of suicide attempts and suicide deaths in bipolar disorder: Part II of a report of the International Society for Bipolar Disorders Task Force on Suicide in Bipolar Disorder. <i>Australian and New Zealand Journal of Psychiatry</i> , 2015, 49, 1006-1020.	1.3	87
75	Epidemiology, neurobiology and pharmacological interventions related to suicide deaths and suicide attempts in bipolar disorder: Part I of a report of the International Society for Bipolar Disorders Task Force on Suicide in Bipolar Disorder. <i>Australian and New Zealand Journal of Psychiatry</i> , 2015, 49, 785-802.	1.3	122
76	Influence of light exposure during early life on the age of onset of bipolar disorder. <i>Journal of Psychiatric Research</i> , 2015, 64, 1-8.	1.5	39
77	Differences and overlap in self-reported symptoms of bipolar disorder and borderline personality disorder. <i>European Psychiatry</i> , 2015, 30, 914-919.	0.1	17
78	Return to work six months after burn: A prospective study at the Helsinki Burn Center. <i>Burns</i> , 2015, 41, 1152-1160.	1.1	21
79	A 5-year prospective study of predictors for functional and work disability among primary care patients with depressive disorders. <i>European Psychiatry</i> , 2015, 30, 51-57.	0.1	22
80	International Society for Bipolar Disorders Task Force on Suicide: meta-analyses and meta-regression of correlates of suicide attempts and suicide deaths in bipolar disorder. <i>Bipolar Disorders</i> , 2015, 17, 1-16.	1.1	265
81	The Effect of Personality on Daily Life Emotional Processes. <i>PLoS ONE</i> , 2014, 9, e110907.	1.1	71
82	Post-discharge suicides of inpatients with bipolar disorder in Finland. <i>Bipolar Disorders</i> , 2014, 16, 867-874.	1.1	34
83	A prospective latent analysis study of Axis I psychiatric co-morbidity of DSM-IV major depressive disorder. <i>Psychological Medicine</i> , 2014, 44, 949-959.	2.7	3
84	Differences in incidence of suicide attempts between bipolar I and II disorders and major depressive disorder. <i>Bipolar Disorders</i> , 2014, 16, 652-661.	1.1	141
85	Predictors of adherence to psychopharmacological and psychosocial treatment in bipolar I or II disorders – an 18-month prospective study. <i>Journal of Affective Disorders</i> , 2014, 155, 110-117.	2.0	42
86	Author's response. <i>Brain Stimulation</i> , 2014, 7, 338.	0.7	0
87	Borderline personality disorder among primary care depressive patients: A five-year study. <i>Journal of Affective Disorders</i> , 2014, 155, 303-306.	2.0	23
88	White matter hyperintensities and cognitive performance in adult patients with bipolar I, bipolar II, and major depressive disorders. <i>European Psychiatry</i> , 2014, 29, 226-232.	0.1	37
89	Temperament and character traits predict future burden of depression. <i>Journal of Affective Disorders</i> , 2014, 158, 139-147.	2.0	46
90	Relationship between sunlight and the age of onset of bipolar disorder: An international multisite study. <i>Journal of Affective Disorders</i> , 2014, 167, 104-111.	2.0	43

#	ARTICLE	IF	CITATIONS
91	Incidence and predictors of suicide attempts among primary-care patients with depressive disorders: a 5-year prospective study. <i>Psychological Medicine</i> , 2014, 44, 291-302.	2.7	44
92	Suicidal Behaviour in Mood Disordersâ€”Who, When, and Why?. <i>Canadian Journal of Psychiatry</i> , 2014, 59, 120-130.	0.9	231
93	Temperament, character and personality disorders. <i>European Psychiatry</i> , 2013, 28, 483-491.	0.1	21
94	Typology of clinical course in bipolar disorder based on 18-month naturalistic follow-up. <i>Psychological Medicine</i> , 2013, 43, 789-799.	2.7	15
95	DEPRESSION AND SMOKING: A 5-YEAR PROSPECTIVE STUDY OF PATIENTS WITH MAJOR DEPRESSIVE DISORDER. <i>Depression and Anxiety</i> , 2013, 30, 580-588.	2.0	28
96	Concomitants of Family Histories of Mood Disorders and Alcoholism in a Clinical Cohort of Patients With Bipolar I and II Disorder. <i>Journal of Nervous and Mental Disease</i> , 2012, 200, 388-394.	0.5	7
97	Do antidepressants change personality?â€”A five-year observational study. <i>Journal of Affective Disorders</i> , 2012, 142, 200-207.	2.0	13
98	NEUROTICISM MEDIATES THE EFFECT OF P2RX7 ON OUTCOMES OF MOOD DISORDERS. <i>Depression and Anxiety</i> , 2012, 29, 816-823.	2.0	14
99	Family history of psychiatric disorders and the outcome of psychiatric patients with DSM-IV major depressive disorder. <i>Journal of Affective Disorders</i> , 2011, 131, 251-259.	2.0	11
100	<i>P2RX7</i> gene is associated consistently with mood disorders and predicts clinical outcome in three clinical cohorts. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2011, 156, 435-447.	1.1	79
101	Differences in temperament and character dimensions in patients with bipolar I or II or major depressive disorder and general population subjects. <i>Psychological Medicine</i> , 2011, 41, 1579-1591.	2.7	32
102	Major depressive disorder and white matter abnormalities: A diffusion tensor imaging study with tract-based spatial statistics. <i>Journal of Affective Disorders</i> , 2010, 120, 240-244.	2.0	160
103	Maintenance treatment received by patients with bipolar I and II disorders â€” A naturalistic prospective study. <i>Journal of Affective Disorders</i> , 2010, 121, 116-126.	2.0	12
104	Differences in neuroticism and extraversion between patients with bipolar I or II and general population subjects or major depressive disorder patients. <i>Journal of Affective Disorders</i> , 2010, 125, 42-52.	2.0	45
105	Treatment attitudes and adherence of psychiatric patients with major depressive disorder: A five-year prospective study. <i>Journal of Affective Disorders</i> , 2010, 127, 102-112.	2.0	37
106	A prospective latent analyses study of psychiatric comorbidity of DSM-IV bipolar I and II disorders. <i>Bipolar Disorders</i> , 2010, 12, 271-284.	1.1	48
107	Incidence and Predictors of Suicide Attempts in DSM-IV Major Depressive Disorder: A Five-Year Prospective Study. <i>American Journal of Psychiatry</i> , 2010, 167, 801-808.	4.0	195
108	Categorical and Dimensional Stability of Comorbid Personality Disorder Symptoms in DSM-IV Major Depressive Disorder. <i>Journal of Clinical Psychiatry</i> , 2010, 71, 287-295.	1.1	26

#	ARTICLE	IF	CITATIONS
109	Screening of psychiatric outpatients for borderline personality disorder with the McLean Screening Instrument for Borderline Personality Disorder (MSI-BPD). <i>Nordic Journal of Psychiatry</i> , 2009, 63, 475-479.	0.7	36
110	Relationships of neuroticism and extraversion with axis I and II comorbidity among patients with DSM-IV major depressive disorder. <i>Journal of Affective Disorders</i> , 2009, 114, 110-121.	2.0	44
111	Hopelessness across phases of bipolar I or II disorder: A prospective study. <i>Journal of Affective Disorders</i> , 2009, 115, 11-17.	2.0	15
112	How suicidal bipolar patients are depends on how suicidal ideation is defined. <i>Journal of Affective Disorders</i> , 2009, 118, 48-54.	2.0	22
113	Neuroticism, introversion, and major depressive disorder-traits, states, or scars?. <i>Depression and Anxiety</i> , 2009, 26, 325-334.	2.0	60
114	Treatment of major depressive disorder in the Finnish general population. <i>Depression and Anxiety</i> , 2009, 26, 1049-1059.	2.0	59
115	Gender differences in bipolar disorder type I and II. <i>Acta Psychiatrica Scandinavica</i> , 2009, 120, 464-473.	2.2	51
116	Clinical predictors of unrecognized bipolar I and II disorders. <i>Bipolar Disorders</i> , 2008, 10, 238-244.	1.1	20
117	Differences in outcome of DSM-IV bipolar I and II disorders. <i>Bipolar Disorders</i> , 2008, 10, 413-425.	1.1	75
118	Differences in incidence of suicide attempts during phases of bipolar I and II disorders. <i>Bipolar Disorders</i> , 2008, 10, 588-596.	1.1	107
119	Antidepressant utilisation patterns and determinants of short-term and non-psychiatric use in the Finnish general adult population. <i>Journal of Affective Disorders</i> , 2008, 110, 94-105.	2.0	70
120	Only half of bipolar I and II patients report prodromal symptoms. <i>Journal of Affective Disorders</i> , 2008, 111, 366-371.	2.0	31
121	Maintenance pharmacotherapy for recurrent major depressive disorder: 5-year follow-up study. <i>British Journal of Psychiatry</i> , 2008, 193, 163-164.	1.7	13
122	Long-Term Outcome of Major Depressive Disorder in Psychiatric Patients Is Variable. <i>Journal of Clinical Psychiatry</i> , 2008, 69, 196-205.	1.1	95
123	Predictors for Switch From Unipolar Major Depressive Disorder to Bipolar Disorder Type I or II. <i>Journal of Clinical Psychiatry</i> , 2008, 69, 1267-1275.	1.1	49
124	Do characteristics of patients with major depressive disorder differ between primary and psychiatric care?. <i>Psychological Medicine</i> , 2007, 37, 893-904.	2.7	31
125	Lifetime Prevalence of Psychotic and Bipolar I Disorders in a General Population. <i>Archives of General Psychiatry</i> , 2007, 64, 19.	13.8	1,112
126	Early age at onset of bipolar disorder is associated with more severe clinical features but delayed treatment seeking. <i>Bipolar Disorders</i> , 2007, 9, 698-705.	1.1	103



#	ARTICLE	IF	CITATIONS
127	Suicidal behaviour during different phases of bipolar disorder. <i>Journal of Affective Disorders</i> , 2007, 97, 101-107.	2.0	93
128	Adequacy of Treatment Received by Diagnosed and Undiagnosed Patients With Bipolar I and II Disorders. <i>Journal of Clinical Psychiatry</i> , 2007, 68, 102-110.	1.1	35
129	Determinants of Functional Disability and Social Adjustment in Major Depressive Disorder. <i>Journal of Nervous and Mental Disease</i> , 2006, 194, 570-576.	0.5	44
130	Prospective study of risk factors for attempted suicide among patients with bipolar disorder. <i>Bipolar Disorders</i> , 2006, 8, 576-585.	1.1	112
131	Decline in suicidal ideation among patients with MDD is preceded by decline in depression and hopelessness. <i>Journal of Affective Disorders</i> , 2006, 95, 95-102.	2.0	58
132	The relationship of neuroticism and extraversion to symptoms of anxiety and depression in the general population. <i>Depression and Anxiety</i> , 2006, 23, 281-289.	2.0	260
133	Differences in Axis I and II Comorbidity Between Bipolar I and II Disorders and Major Depressive Disorder. <i>Journal of Clinical Psychiatry</i> , 2006, 67, 584-593.	1.1	85
134	Functional and Work Disability in Major Depressive Disorder. <i>Journal of Nervous and Mental Disease</i> , 2005, 193, 189-195.	0.5	62
135	Depressive disorders in primary care: recurrent, chronic, and co-morbid. <i>Psychological Medicine</i> , 2005, 35, 673-682.	2.7	84
136	Prospective study of risk factors for attempted suicide among patients with DSM-IV major depressive disorder. <i>British Journal of Psychiatry</i> , 2005, 186, 314-318.	1.7	161
137	DSM-IV mood-, anxiety- and alcohol use disorders and their comorbidity in the Finnish general population. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2005, 40, 1-10.	1.6	318
138	Childhood adversities as risk factors for adult mental disorders. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2005, 40, 769-777.	1.6	130
139	Continuity Is the Main Challenge in Treating Major Depressive Disorder in Psychiatric Care. <i>Journal of Clinical Psychiatry</i> , 2005, 66, 220-227.	1.1	90
140	Suicidal Ideation and Attempts in Bipolar I and II Disorders. <i>Journal of Clinical Psychiatry</i> , 2005, 66, 1456-1462.	1.1	204
141	The clinical characteristics of DSM-IV bipolar I and II disorders: baseline findings from the Jorvi Bipolar Study (JoBS). <i>Bipolar Disorders</i> , 2004, 6, 395-405.	1.1	130
142	Use of health services for major depressive episode in Finland. <i>Journal of Affective Disorders</i> , 2004, 79, 105-112.	2.0	60
143	Life Events, Social Support, and Onset of Major Depressive Episode in Finnish Patients. <i>Journal of Nervous and Mental Disease</i> , 2004, 192, 373-381.	0.5	36
144	Co-morbidity and stability of melancholic features in DSM-IV major depressive disorder. <i>Psychological Medicine</i> , 2004, 34, 1443-1452.	2.7	57

#	ARTICLE	IF	CITATIONS
145	Severity and Comorbidity Predict Episode Duration and Recurrence of DSM-IV Major Depressive Disorder. <i>Journal of Clinical Psychiatry</i> , 2004, 65, 810-819.	1.1	135
146	The Mood Disorder Questionnaire improves recognition of bipolar disorder in psychiatric care. <i>BMC Psychiatry</i> , 2003, 3, 8.	1.1	84
147	Do Means Matter?. <i>Journal of Nervous and Mental Disease</i> , 2003, 191, 745-750.	0.5	43
148	Suicidal Ideation and Attempts Among Psychiatric Patients With Major Depressive Disorder. <i>Journal of Clinical Psychiatry</i> , 2003, 64, 1094-1100.	1.1	282
149	Current Comorbidity of Psychiatric Disorders Among DSM-IV Major Depressive Disorder Patients in Psychiatric Care in the Vantaa Depression Study. <i>Journal of Clinical Psychiatry</i> , 2002, 63, 126-134.	1.1	202
150	Cigarette smoking, alcohol intoxication and major depressive episode in a representative population sample. <i>Journal of Epidemiology and Community Health</i> , 2001, 55, 573-576.	2.0	57
151	Disability Pension for Major Depression in Finland. <i>American Journal of Psychiatry</i> , 2000, 157, 1869-1872.	4.0	30
152	The 12-month prevalence and risk factors for major depressive episode in Finland: representative sample of 5993 adults. <i>Acta Psychiatrica Scandinavica</i> , 2000, 102, 178-184.	2.2	134
153	Suicide attempts preceding completed suicide. <i>British Journal of Psychiatry</i> , 1998, 173, 531-535.	1.7	243
154	Employment Status Influences the Weekly Patterns of Suicide among Alcohol Misusers. <i>Alcoholism: Clinical and Experimental Research</i> , 1997, 21, 1704-1706.	1.4	18
155	Differences in Recent Life Events between Alcoholic and Depressive Nonalcoholic Suicides. <i>Alcoholism: Clinical and Experimental Research</i> , 1994, 18, 1143-1149.	1.4	62
156	Completed suicide and recent lithium treatment. <i>Journal of Affective Disorders</i> , 1992, 26, 101-103.	2.0	35