Erkki Isomets

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/457575/erkki-isometsa-publications-by-year.pdf

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

6,859 80 151 44 h-index g-index citations papers 164 7,712 5.2 5.71 L-index avg, IF ext. citations ext. papers

#	Paper	IF	Citations
151	A Randomized Clinical Trial of Attempted Suicide Short Intervention Program versus Crisis Counseling in Preventing Repeat Suicide Attempts: A Two-Year Follow-Up Study <i>Psychotherapy and Psychosomatics</i> , 2022 , 1-10	9.4	O
150	Implementation of CYP2D6 copy-number imputation panel and frequency of key pharmacogenetic variants in Finnish individuals with a psychotic disorder <i>Pharmacogenomics Journal</i> , 2022 ,	3.5	1
149	Suicide Risk Among Psychiatric Patients 2022 , 1-19		
148	A randomized clinical trial of the ASSIP vs. crisis counselling in preventing suicide attempt repetition: a two-year follow-up study <i>Nordic Journal of Psychiatry</i> , 2022 , 1	2.3	
147	SYMPOSIUM 1: suicide prevention: Nordic Perspectives <i>Nordic Journal of Psychiatry</i> , 2022 , 1	2.3	
146	Differences in psychosocial functioning between psychotic disorders in the Finnish SUPER study <i>Schizophrenia Research</i> , 2022 , 244, 10-17	3.6	
145	Reduced visual contrast suppression during major depressive episodes. <i>Journal of Psychiatry and Neuroscience</i> , 2021 , 46, E222-E231	4.5	6
144	White Matter Hyperintensities after Five-Year Follow-Up and a Cross-Sectional FA Decrease in Bipolar I and Major Depressive Patients. <i>Neuropsychobiology</i> , 2021 , 1-12	4	
143	Variants in regulatory elements of PDE4D associate with major mental illness in the Finnish population. <i>Molecular Psychiatry</i> , 2021 , 26, 816-824	15.1	4
142	Presence and Overlap of Bipolar Symptoms and Borderline Features during Major Depressive Episodes. <i>Journal of Affective Disorders</i> , 2021 , 280, 467-477	6.6	3
141	Effectiveness of brief schema group therapy for borderline personality disorder symptoms: a randomized pilot study. <i>Nordic Journal of Psychiatry</i> , 2021 , 75, 176-185	2.3	Ο
140	Escitalopram enhances synchrony of brain responses during emotional narratives in patients with major depressive disorder. <i>NeuroImage</i> , 2021 , 237, 118110	7.9	1
139	The evidence does not support the premises of the environmental mismatch hypothesis. Response to "Bipolar disorder: An evolutionary psychoneuroimmunological approach". <i>Neuroscience and Biobehavioral Reviews</i> , 2021 , 131, 539-540	9	
138	Variations in seasonal solar insolation are associated with a history of suicide attempts in bipolar I disorder. <i>International Journal of Bipolar Disorders</i> , 2021 , 9, 26	5.4	0
137	Diagnostic conversion from unipolar depression to bipolar disorder, schizophrenia, or schizoaffective disorder: A nationwide prospective 15-year register study on 43\(^1495\) inpatients. <i>Bipolar Disorders</i> , 2020 , 22, 582-592	3.8	5
136	Comorbid alcohol use disorder in psychiatric MDD patients: A five-year prospective study. <i>Journal of Affective Disorders</i> , 2020 , 267, 283-288	6.6	5
135	Borderline Personality Disorder With Depression Confers Significant Risk of Suicidal Behavior in Mood Disorder Patients-A Comparative Study. <i>Frontiers in Psychiatry</i> , 2020 , 11, 290	5	7

(2018-2020)

134	Role of Hopelessness in Suicidal Ideation Among Patients With Depressive Disorders. <i>Journal of Clinical Psychiatry</i> , 2020 , 81,	4.6	4
133	Pregnancy, delivery and postpartum in women with schizophrenia or schizoaffective disorder in Finland: a national register-based comparative study. <i>Psychiatry Research</i> , 2020 , 294, 113504	9.9	1
132	T49. SOCIODEMOGRAPHIC AND LIFESTYLE FACTORS ASSOCIATED WITH GOOD PERFORMANCE IN PAIRED ASSOCIATES LEARNING TEST IN PATIENTS WITH SCHIZOPHRENIA. <i>Schizophrenia Bulletin</i> , 2020 , 46, S250-S250	1.3	78
131	Suicides in Mood Disorders in Psychiatric Settings in Nordic National Register-Based Studies. <i>Frontiers in Psychiatry</i> , 2020 , 11, 721	5	6
130	Do Suicide Attempts of Mood Disorder Patients Directly Increase the Risk for a Reattempt?. <i>Frontiers in Psychiatry</i> , 2020 , 11, 547791	5	2
129	Schizophrenia and pregnancy: a national register-based follow-up study among Finnish women born between 1965 and 1980. <i>Archives of Womenrs Mental Health</i> , 2020 , 23, 91-100	5	6
128	Association between solar insolation and a history of suicide attempts in bipolar I disorder. <i>Journal of Psychiatric Research</i> , 2019 , 113, 1-9	5.2	15
127	The Relationship Between Smoking and Suicidal Behavior in Psychiatric Patients with Major Depressive Disorder. <i>Archives of Suicide Research</i> , 2019 , 23, 590-604	2.3	3
126	Clinical course predicts long-term outcomes in bipolar disorder. <i>Psychological Medicine</i> , 2019 , 49, 1109-	16.57	2
125	Machine learning and big data analytics in bipolar disorder: A position paper from the International Society for Bipolar Disorders Big Data Task Force. <i>Bipolar Disorders</i> , 2019 , 21, 582-594	3.8	52
124	Risk factors for suicide in depression in Finland: first-hospitalized patients followed up to 24 lyears. <i>Acta Psychiatrica Scandinavica</i> , 2019 , 139, 154-163	6.5	18
123	Predominant polarity in bipolar I and II disorders: A five-year follow-up study. <i>Journal of Affective Disorders</i> , 2019 , 246, 806-813	6.6	16
122	Maternal schizophrenia and out-of-home placements of offspring: A national follow-up study among Finnish women born 1965-1980 and their children. <i>Psychiatry Research</i> , 2019 , 273, 9-14	9.9	8
121	Short-term escitalopram treatment normalizes aberrant self-referential processing in major depressive disorder. <i>Journal of Affective Disorders</i> , 2018 , 236, 222-229	6.6	35
120	O5.5. SLEEP IN MAJOR PSYCHIATRIC DISORDERS: RESULTS FROM NATIONWIDE SUPER FINLAND STUDY. <i>Schizophrenia Bulletin</i> , 2018 , 44, S88-S88	1.3	78
119	Features of borderline personality disorder as a mediator of the relation between childhood traumatic experiences and psychosis-like experiences in patients with mood disorder. <i>European Psychiatry</i> , 2018 , 49, 9-15	6	3
118	Decline in suicide mortality after psychiatric hospitalization for depression in Finland between 1991 and 2014. <i>World Psychiatry</i> , 2018 , 17, 110-112	14.4	11
117	Self-reported psychosis-like experiences in patients with mood disorders. <i>European Psychiatry</i> , 2018 , 51, 90-97	6	5

116	Schizophrenia and induced abortions: A national register-based follow-up study among Finnish women born between 1965-1980 with schizophrenia or schizoaffective disorder. <i>Schizophrenia Research</i> , 2018 , 192, 142-147	3.6	13
115	Alcohol use and smoking in burn patients at the Helsinki Burn Center. <i>Burns</i> , 2018 , 44, 158-167	2.3	13
114	State and trait hopelessness in a prospective five-year study of patients with depressive disorders. Journal of Affective Disorders, 2018, 239, 107-114	6.6	9
113	Obstetric and perinatal health outcomes related to schizophrenia: A national register-based follow-up study among Finnish women born between 1965 and 1980 and their offspring. <i>European Psychiatry</i> , 2018 , 52, 68-75	6	12
112	Self-reported treatment adherence among psychiatric in- and outpatients. <i>Nordic Journal of Psychiatry</i> , 2018 , 72, 526-533	2.3	4
111	Maintenance pharmacotherapy for recurrent major depressive disorder in primary care: A 5-year follow-up study. <i>European Psychiatry</i> , 2017 , 41, 111-114	6	3
110	Incidence and predictors of suicide attempts in bipolar I and II disorders: A 5-year follow-up study. <i>Bipolar Disorders</i> , 2017 , 19, 13-22	3.8	31
109	Level of functioning, perceived work ability, and work status among psychiatric patients with major mental disorders. <i>European Psychiatry</i> , 2017 , 44, 83-89	6	16
108	Mood Zoom could be a promising tool for daily mood variability monitoring, potentially differentiating bipolar from borderline patients. <i>Evidence-Based Mental Health</i> , 2017 , 20, e9	11.1	1
107	Dangerously mixed. <i>Bipolar Disorders</i> , 2017 , 19, 314-315	3.8	
106	Mediating role of borderline personality disorder traits in the effects of childhood maltreatment on suicidal behaviour among mood disorder patients. <i>European Psychiatry</i> , 2017 , 44, 53-60	3.8 6	9
, ,	Mediating role of borderline personality disorder traits in the effects of childhood maltreatment on		9
106	Mediating role of borderline personality disorder traits in the effects of childhood maltreatment on suicidal behaviour among mood disorder patients. <i>European Psychiatry</i> , 2017 , 44, 53-60 Single dose of mirtazapine modulates whole-brain functional connectivity during emotional	6	
106	Mediating role of borderline personality disorder traits in the effects of childhood maltreatment on suicidal behaviour among mood disorder patients. <i>European Psychiatry</i> , 2017 , 44, 53-60 Single dose of mirtazapine modulates whole-brain functional connectivity during emotional narrative processing. <i>Psychiatry Research - Neuroimaging</i> , 2017 , 263, 61-69 Prevalence of current anxiety disorders in people with bipolar disorder during euthymia: a	6	6
106 105 104	Mediating role of borderline personality disorder traits in the effects of childhood maltreatment on suicidal behaviour among mood disorder patients. <i>European Psychiatry</i> , 2017 , 44, 53-60 Single dose of mirtazapine modulates whole-brain functional connectivity during emotional narrative processing. <i>Psychiatry Research - Neuroimaging</i> , 2017 , 263, 61-69 Prevalence of current anxiety disorders in people with bipolar disorder during euthymia: a meta-analysis. <i>Psychological Medicine</i> , 2017 , 47, 1107-1115 Relationships between self-reported childhood traumatic experiences, attachment style, neuroticism and features of borderline personality disorders in patients with mood disorders.	6.9	25
106 105 104	Mediating role of borderline personality disorder traits in the effects of childhood maltreatment on suicidal behaviour among mood disorder patients. <i>European Psychiatry</i> , 2017 , 44, 53-60 Single dose of mirtazapine modulates whole-brain functional connectivity during emotional narrative processing. <i>Psychiatry Research - Neuroimaging</i> , 2017 , 263, 61-69 Prevalence of current anxiety disorders in people with bipolar disorder during euthymia: a meta-analysis. <i>Psychological Medicine</i> , 2017 , 47, 1107-1115 Relationships between self-reported childhood traumatic experiences, attachment style, neuroticism and features of borderline personality disorders in patients with mood disorders. <i>Journal of Affective Disorders</i> , 2017 , 210, 82-89 Incidence and Predictors of Suicide Attempts in Bipolar I and II Disorders: A Five-year Follow-up.	6 2.9 6.9	25
106 105 104 103	Mediating role of borderline personality disorder traits in the effects of childhood maltreatment on suicidal behaviour among mood disorder patients. <i>European Psychiatry</i> , 2017 , 44, 53-60 Single dose of mirtazapine modulates whole-brain functional connectivity during emotional narrative processing. <i>Psychiatry Research - Neuroimaging</i> , 2017 , 263, 61-69 Prevalence of current anxiety disorders in people with bipolar disorder during euthymia: a meta-analysis. <i>Psychological Medicine</i> , 2017 , 47, 1107-1115 Relationships between self-reported childhood traumatic experiences, attachment style, neuroticism and features of borderline personality disorders in patients with mood disorders. <i>Journal of Affective Disorders</i> , 2017 , 210, 82-89 Incidence and Predictors of Suicide Attempts in Bipolar I and II Disorders: A Five-year Follow-up. <i>European Psychiatry</i> , 2017 , 41, S87-S87	6 2.9 6.9 6.6	6 25 25

(2015-2016)

98	Health-related quality of life of primary care patients with depressive disorders. <i>European Psychiatry</i> , 2016 , 37, 28-34	6	10
97	Pharmacological and psychosocial treatment of depression in primary care: Low intensity and poor adherence and continuity. <i>Journal of Affective Disorders</i> , 2016 , 202, 145-52	6.6	14
96	Large-scale suicide prevention by pharmacological treatment of mood disorders. <i>European Psychiatry</i> , 2016 , 33, S53-S53	6	
95	Differences and similarities of risk factors for suicidal ideation and attempts among patients with depressive or bipolar disorders. <i>Journal of Affective Disorders</i> , 2016 , 193, 318-30	6.6	50
94	A single dose of mirtazapine attenuates neural responses to self-referential processing. <i>Journal of Psychopharmacology</i> , 2016 , 30, 23-32	4.6	13
93	Self-reported symptoms of schizotypal and borderline personality disorder in patients with mood disorders. <i>European Psychiatry</i> , 2016 , 33, 37-44	6	11
92	Response style and severity and chronicity of depressive disorders in primary health care. <i>European Psychiatry</i> , 2016 , 33, 1-8	6	
91	Personality disorders and suicide attempts in unipolar and bipolar mood disorders. <i>Journal of Affective Disorders</i> , 2016 , 190, 632-639	6.6	18
90	Temperament, character, and suicide attempts in unipolar and bipolar mood disorders. <i>Journal of Clinical Psychiatry</i> , 2016 , 77, 252-60	4.6	23
89	Rumination And Primary Care Depression. European Psychiatry, 2016, 33, S422-S422	6	1
88	A review of factors associated with greater likelihood of suicide attempts and suicide deaths in bipolar disorder: Part II of a report of the International Society for Bipolar Disorders Task Force on Suicide in Bipolar Disorder. <i>Australian and New Zealand Journal of Psychiatry</i> , 2015 , 49, 1006-20	2.6	68
87	Epidemiology, neurobiology and pharmacological interventions related to suicide deaths and suicide attempts in bipolar disorder: Part I of a report of the International Society for Bipolar Disorders Task Force on Suicide in Bipolar Disorder. <i>Australian and New Zealand Journal of</i>	2.6	91
86	Influence of light exposure during early life on the age of onset of bipolar disorder. <i>Journal of Psychiatric Research</i> , 2015 , 64, 1-8	5.2	34
85	Differences and overlap in self-reported symptoms of bipolar disorder and borderline personality disorder. <i>European Psychiatry</i> , 2015 , 30, 914-9	6	13
84	Return to work six months after burn: a prospective study at the Helsinki Burn Center. <i>Burns</i> , 2015 , 41, 1152-60	2.3	13
83	A 5-year prospective study of predictors for functional and work disability among primary care patients with depressive disorders. <i>European Psychiatry</i> , 2015 , 30, 51-7	6	16
	taran da antara da a		
82	International Society for Bipolar Disorders Task Force on Suicide: meta-analyses and meta-regression of correlates of suicide attempts and suicide deaths in bipolar disorder. <i>Bipolar Disorders</i> , 2015 , 17, 1-16	3.8	202

80	Five-year outcome of bipolar I and II disorders: findings of the Jorvi Bipolar Study. <i>Bipolar Disorders</i> , 2015 , 17, 363-74	3.8	32
79	Health-related quality of life 6 months after burns among hospitalized patients: Predictive importance of mental disorders and burn severity. <i>Burns</i> , 2015 , 41, 742-8	2.3	20
78	Validity and reliability of the Finnish version of the Functioning Assessment Short Test (FAST) in bipolar disorder. <i>International Journal of Bipolar Disorders</i> , 2015 , 3, 10	5.4	5
77	Long-term personality changes and predictive adaptive responses after depressive episodes. <i>Evolution and Human Behavior</i> , 2015 , 36, 337-344	4	45
76	Predictors of adherence to psychopharmacological and psychosocial treatment in bipolar I or II disorders - an 18-month prospective study. <i>Journal of Affective Disorders</i> , 2014 , 155, 110-7	6.6	39
75	Author⊌ response. Brain Stimulation, 2014, 7, 338	5.1	
74	Borderline personality disorder among primary care depressive patients: a five-year study. <i>Journal of Affective Disorders</i> , 2014 , 155, 303-6	6.6	18
73	White matter hyperintensities and cognitive performance in adult patients with bipolar I, bipolar II, and major depressive disorders. <i>European Psychiatry</i> , 2014 , 29, 226-32	6	31
72	Temperament and character traits predict future burden of depression. <i>Journal of Affective Disorders</i> , 2014 , 158, 139-47	6.6	36
71	Relationship between sunlight and the age of onset of bipolar disorder: an international multisite study. <i>Journal of Affective Disorders</i> , 2014 , 167, 104-11	6.6	37
70	Incidence and predictors of suicide attempts among primary-care patients with depressive disorders: a 5-year prospective study. <i>Psychological Medicine</i> , 2014 , 44, 291-302	6.9	29
69	Suicidal behaviour in mood disorderswho, when, and why?. <i>Canadian Journal of Psychiatry</i> , 2014 , 59, 120-30	4.8	173
68	The effect of personality on daily life emotional processes. PLoS ONE, 2014, 9, e110907	3.7	48
67	Post-discharge suicides of inpatients with bipolar disorder in Finland. <i>Bipolar Disorders</i> , 2014 , 16, 867-7	43.8	29
66	A prospective latent analysis study of Axis I psychiatric co-morbidity of DSM-IV major depressive disorder. <i>Psychological Medicine</i> , 2014 , 44, 949-59	6.9	3
65	Differences in incidence of suicide attempts between bipolar I and II disorders and major depressive disorder. <i>Bipolar Disorders</i> , 2014 , 16, 652-61	3.8	108
64	Temperament, character and personality disorders. European Psychiatry, 2013, 28, 483-91	6	15
63	Typology of clinical course in bipolar disorder based on 18-month naturalistic follow-up. <i>Psychological Medicine</i> , 2013 , 43, 789-99	6.9	13

(2009-2013)

62	Depression and smoking: a 5-year prospective study of patients with major depressive disorder. <i>Depression and Anxiety</i> , 2013 , 30, 580-8	8.4	25
61	Do antidepressants change personality?a five-year observational study. <i>Journal of Affective Disorders</i> , 2012 , 142, 200-7	6.6	10
60	Neuroticism mediates the effect of P2RX7 on outcomes of mood disorders. <i>Depression and Anxiety</i> , 2012 , 29, 816-23	8.4	11
59	Concomitants of family histories of mood disorders and alcoholism in a clinical cohort of patients with bipolar I and II disorder. <i>Journal of Nervous and Mental Disease</i> , 2012 , 200, 388-94	1.8	6
58	Family history of psychiatric disorders and the outcome of psychiatric patients with DSM-IV major depressive disorder. <i>Journal of Affective Disorders</i> , 2011 , 131, 251-9	6.6	7
57	P2RX7 gene is associated consistently with mood disorders and predicts clinical outcome in three clinical cohorts. <i>American Journal of Medical Genetics Part B: Neuropsychiatric Genetics</i> , 2011 , 156B, 435-	47 ⁵	67
56	Differences in temperament and character dimensions in patients with bipolar I or II or major depressive disorder and general population subjects. <i>Psychological Medicine</i> , 2011 , 41, 1579-91	6.9	29
55	A prospective latent analyses study of psychiatric comorbidity of DSM-IV bipolar I and II disorders. <i>Bipolar Disorders</i> , 2010 , 12, 271-84	3.8	39
54	Incidence and predictors of suicide attempts in DSM-IV major depressive disorder: a five-year prospective study. <i>American Journal of Psychiatry</i> , 2010 , 167, 801-8	11.9	147
53	Major depressive disorder and white matter abnormalities: a diffusion tensor imaging study with tract-based spatial statistics. <i>Journal of Affective Disorders</i> , 2010 , 120, 240-4	6.6	143
52	Maintenance treatment received by patients with bipolar I and II disordersa naturalistic prospective study. <i>Journal of Affective Disorders</i> , 2010 , 121, 116-26	6.6	9
51	Differences in neuroticism and extraversion between patients with bipolar I or II and general population subjects or major depressive disorder patients. <i>Journal of Affective Disorders</i> , 2010 , 125, 42-	52.6 52.6	40
50	Treatment attitudes and adherence of psychiatric patients with major depressive disorder: a five-year prospective study. <i>Journal of Affective Disorders</i> , 2010 , 127, 102-12	6.6	30
49	Categorical and dimensional stability of comorbid personality disorder symptoms in DSM-IV major depressive disorder: a prospective study. <i>Journal of Clinical Psychiatry</i> , 2010 , 71, 287-95	4.6	20
48	Screening of psychiatric outpatients for borderline personality disorder with the McLean Screening Instrument for Borderline Personality Disorder (MSI-BPD). <i>Nordic Journal of Psychiatry</i> , 2009 , 63, 475-9	2.3	31
47	Relationships of neuroticism and extraversion with axis I and II comorbidity among patients with DSM-IV major depressive disorder. <i>Journal of Affective Disorders</i> , 2009 , 114, 110-21	6.6	36
46	Hopelessness across phases of bipolar I or II disorder: a prospective study. <i>Journal of Affective Disorders</i> , 2009 , 115, 11-7	6.6	13
45	How suicidal bipolar patients are depends on how suicidal ideation is defined. <i>Journal of Affective Disorders</i> , 2009 , 118, 48-54	6.6	16

44	Neuroticism, introversion, and major depressive disordertraits, states, or scars?. <i>Depression and Anxiety</i> , 2009 , 26, 325-34	8.4	44
43	Treatment of major depressive disorder in the Finnish general population. <i>Depression and Anxiety</i> , 2009 , 26, 1049-59	8.4	56
42	Gender differences in bipolar disorder type I and II. Acta Psychiatrica Scandinavica, 2009, 120, 464-73	6.5	42
41	Clinical predictors of unrecognized bipolar I and II disorders. <i>Bipolar Disorders</i> , 2008 , 10, 238-44	3.8	18
40	Differences in outcome of DSM-IV bipolar I and II disorders. <i>Bipolar Disorders</i> , 2008 , 10, 413-25	3.8	70
39	Differences in incidence of suicide attempts during phases of bipolar I and II disorders. <i>Bipolar Disorders</i> , 2008 , 10, 588-96	3.8	91
38	Antidepressant utilisation patterns and determinants of short-term and non-psychiatric use in the Finnish general adult population. <i>Journal of Affective Disorders</i> , 2008 , 110, 94-105	6.6	60
37	Only half of bipolar I and II patients report prodromal symptoms. <i>Journal of Affective Disorders</i> , 2008 , 111, 366-71	6.6	27
36	Maintenance pharmacotherapy for recurrent major depressive disorder: 5-year follow-up study. <i>British Journal of Psychiatry</i> , 2008 , 193, 163-4	5.4	12
35	Long-term outcome of major depressive disorder in psychiatric patients is variable. <i>Journal of Clinical Psychiatry</i> , 2008 , 69, 196-205	4.6	81
34	Predictors for switch from unipolar major depressive disorder to bipolar disorder type I or II: a 5-year prospective study. <i>Journal of Clinical Psychiatry</i> , 2008 , 69, 1267-75	4.6	41
33	Early age at onset of bipolar disorder is associated with more severe clinical features but delayed treatment seeking. <i>Bipolar Disorders</i> , 2007 , 9, 698-705	3.8	84
32	Suicidal behaviour during different phases of bipolar disorder. <i>Journal of Affective Disorders</i> , 2007 , 97, 101-7	6.6	80
31	Do characteristics of patients with major depressive disorder differ between primary and psychiatric care?. <i>Psychological Medicine</i> , 2007 , 37, 893-904	6.9	28
30	Lifetime prevalence of psychotic and bipolar I disorders in a general population. <i>Archives of General Psychiatry</i> , 2007 , 64, 19-28		888
29	Adequacy of treatment received by diagnosed and undiagnosed patients with bipolar I and II disorders. <i>Journal of Clinical Psychiatry</i> , 2007 , 68, 102-10	4.6	32
28	Decline in suicidal ideation among patients with MDD is preceded by decline in depression and hopelessness. <i>Journal of Affective Disorders</i> , 2006 , 95, 95-102	6.6	44
27	The relationship of neuroticism and extraversion to symptoms of anxiety and depression in the general population. <i>Depression and Anxiety</i> , 2006 , 23, 281-9	8.4	205

(2003-2006)

26	Determinants of functional disability and social adjustment in major depressive disorder: a prospective study. <i>Journal of Nervous and Mental Disease</i> , 2006 , 194, 570-6	1.8	37
25	Prospective study of risk factors for attempted suicide among patients with bipolar disorder. <i>Bipolar Disorders</i> , 2006 , 8, 576-85	3.8	96
24	Differences in Axis I and II comorbidity between bipolar I and II disorders and major depressive disorder. <i>Journal of Clinical Psychiatry</i> , 2006 , 67, 584-93	4.6	71
23	Functional and work disability in major depressive disorder. <i>Journal of Nervous and Mental Disease</i> , 2005 , 193, 189-95	1.8	54
22	Depressive disorders in primary care: recurrent, chronic, and co-morbid. <i>Psychological Medicine</i> , 2005 , 35, 673-82	6.9	73
21	Prospective study of risk factors for attempted suicide among patients with DSM-IV major depressive disorder. <i>British Journal of Psychiatry</i> , 2005 , 186, 314-8	5.4	127
20	DSM-IV mood-, anxiety- and alcohol use disorders and their comorbidity in the Finnish general populationresults from the Health 2000 Study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2005 , 40, 1-10	4.5	281
19	Childhood adversities as risk factors for adult mental disorders: results from the Health 2000 study. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2005 , 40, 769-77	4.5	115
18	Continuity is the main challenge in treating major depressive disorder in psychiatric care. <i>Journal of Clinical Psychiatry</i> , 2005 , 66, 220-7	4.6	71
17	Suicidal ideation and attempts in bipolar I and II disorders. Journal of Clinical Psychiatry, 2005, 66, 1456	-62 .6	174
16	The clinical characteristics of DSM-IV bipolar I and II disorders: baseline findings from the Jorvi Bipolar Study (JoBS). <i>Bipolar Disorders</i> , 2004 , 6, 395-405	3.8	114
15	Use of health services for major depressive episode in Finland. <i>Journal of Affective Disorders</i> , 2004 , 79, 105-12	6.6	55
14	Life events, social support, and onset of major depressive episode in Finnish patients. <i>Journal of Nervous and Mental Disease</i> , 2004 , 192, 373-81	1.8	31
13	Co-morbidity and stability of melancholic features in DSM-IV major depressive disorder. <i>Psychological Medicine</i> , 2004 , 34, 1443-52	6.9	47
12	Severity and comorbidity predict episode duration and recurrence of DSM-IV major depressive disorder. <i>Journal of Clinical Psychiatry</i> , 2004 , 65, 810-9	4.6	115
11	Do means matter?: Differences in characteristics of Finnish suicide completers using different methods. <i>Journal of Nervous and Mental Disease</i> , 2003 , 191, 745-50	1.8	36
10	The mood disorder questionnaire improves recognition of bipolar disorder in psychiatric care. <i>BMC Psychiatry</i> , 2003 , 3, 8	4.2	72
9	Suicidal ideation and attempts among psychiatric patients with major depressive disorder. <i>Journal of Clinical Psychiatry</i> , 2003 , 64, 1094-100	4.6	210

8	Current Comorbidity of Psychiatric Disorders Among DSM-IV Major Depressive Disorder Patients in Psychiatric Care in the Vantaa Depression Study. <i>Journal of Clinical Psychiatry</i> , 2002 , 63, 126-134	4.6	163
7	Cigarette smoking, alcohol intoxication and major depressive episode in a representative population sample. <i>Journal of Epidemiology and Community Health</i> , 2001 , 55, 573-6	5.1	49
6	Disability pension for major depression in Finland. <i>American Journal of Psychiatry</i> , 2000 , 157, 1869-72	11.9	22
5	The 12-month prevalence and risk factors for major depressive episode in Finland: representative sample of 5993 adults. <i>Acta Psychiatrica Scandinavica</i> , 2000 , 102, 178-84	6.5	108
4	Suicide attempts preceding completed suicide. British Journal of Psychiatry, 1998, 173, 531-5	5.4	200
3	Employment Status Influences the Weekly Patterns of Suicide among Alcohol Misusers. <i>Alcoholism: Clinical and Experimental Research</i> , 1997 , 21, 1704-1706	3.7	15
2	Differences in recent life events between alcoholic and depressive nonalcoholic suicides. <i>Alcoholism: Clinical and Experimental Research</i> , 1994 , 18, 1143-9	3.7	50
1	Completed suicide and recent lithium treatment. <i>Journal of Affective Disorders</i> , 1992 , 26, 101-3	6.6	29