

Rein Willemze

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4579468/publications.pdf>

Version: 2024-02-01

573
papers

39,056
citations

2203

99
h-index

4323

173
g-index

583
all docs

583
docs citations

583
times ranked

18372
citing authors

#	ARTICLE	IF	CITATIONS
1	WHO-EORTC classification for cutaneous lymphomas. <i>Blood</i> , 2005, 105, 3768-3785.	0.6	3,529
2	EORTC classification for primary cutaneous lymphomas: a proposal from the Cutaneous Lymphoma Study Group of the European Organization for Research and Treatment of Cancer. <i>Blood</i> , 1997, 90, 354-71.	0.6	908
3	The 2018 update of the WHO-EORTC classification for primary cutaneous lymphomas. <i>Blood</i> , 2019, 133, 1703-1714.	0.6	846
4	Autologous or Allogeneic Bone Marrow Transplantation Compared with Intensive Chemotherapy in Acute Myelogenous Leukemia. <i>New England Journal of Medicine</i> , 1995, 332, 217-223.	13.9	804
5	Primary and secondary cutaneous CD30+lymphoproliferative disorders: a report from the Dutch Cutaneous Lymphoma Group on the long-term follow-up data of 219 patients and guidelines for diagnosis and treatment. <i>Blood</i> , 2000, 95, 3653-3661.	0.6	741
6	Subcutaneous panniculitis-like T-cell lymphoma: definition, classification, and prognostic factors: an EORTC Cutaneous Lymphoma Group Study of 83 cases. <i>Blood</i> , 2008, 111, 838-845.	0.6	617
7	Intravascular lymphoma: clinical presentation, natural history, management and prognostic factors in a series of 38 cases, with special emphasis on the "cutaneous variant"™1. <i>British Journal of Haematology</i> , 2004, 127, 173-183.	1.2	535
8	Primary Cutaneous CD8-Positive Epidermotropic Cytotoxic T Cell Lymphomas. <i>American Journal of Pathology</i> , 1999, 155, 483-492.	1.9	476
9	Update on erythrodermic cutaneous T-cell lymphoma: Report of the international society for cutaneous lymphomas. <i>Journal of the American Academy of Dermatology</i> , 2002, 46, 95-106.	0.6	448
10	Hematopoiesis-restricted minor histocompatibility antigens HA-1- or HA-2-specific T cells can induce complete remissions of relapsed leukemia. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003, 100, 2742-2747.	3.3	400
11	Molecular remission in PML/RAR alpha-positive acute promyelocytic leukemia by combined all-trans retinoic acid and idarubicin (AIDA) therapy. Gruppo Italiano-Malattie Ematologiche Maligne dell'Adulto and Associazione Italiana di Ematologia ed Oncologia Pediatrica Cooperative Groups. <i>Blood</i> , 1997, 90, 1014-21.	0.6	375
12	European Organisation for Research and Treatment of Cancer consensus recommendations for the treatment of mycosis fungoides/S�azary syndrome " Update 2017. <i>European Journal of Cancer</i> , 2017, 77, 57-74.	1.3	363
13	Melanocortin 1 Receptor (MC1R) Gene Variants are Associated with an Increased Risk for Cutaneous Melanoma Which is Largely Independent of Skin Type and Hair Color. <i>Journal of Investigative Dermatology</i> , 2001, 117, 294-300.	0.3	351
14	Spectrum of primary cutaneous CD30 (Ki-1)-positive lymphoproliferative disorders. <i>Journal of the American Academy of Dermatology</i> , 1993, 28, 973-980.	0.6	339
15	Cutaneous Lymphoma International Consortium Study of Outcome in Advanced Stages of Mycosis Fungoides and S�azary Syndrome: Effect of Specific Prognostic Markers on Survival and Development of a Prognostic Model. <i>Journal of Clinical Oncology</i> , 2015, 33, 3766-3773.	0.8	328
16	Primary cutaneous CD30-positive large cell lymphoma: Definition of a new type of cutaneous lymphoma with a favorable prognosis.A European multicenter study of 47 patients. <i>Cancer</i> , 1993, 71, 2097-2104.	2.0	326
17	Mycosis Fungoides. <i>Archives of Dermatology</i> , 2000, 136, 504-10.	1.7	313
18	WHO/EORTC classification of cutaneous lymphomas 2005: histological and molecular aspects. <i>Journal of Cutaneous Pathology</i> , 2005, 32, 647-674.	0.7	313

#	ARTICLE	IF	CITATIONS
19	Progress in allogeneic bone marrow and peripheral blood stem cell transplantation for multiple myeloma: a comparison between transplants performed 1983-93 and 1994-98 at European Group for Blood and Marrow Transplantation centres. <i>British Journal of Haematology</i> , 2001, 113, 209-216.	1.2	307
20	The Influence of Painful Sunburns and Lifetime Sun Exposure on the Risk of Actinic Keratoses, Seborrhic Warts, Melanocytic Nevi, Atypical Nevi, and Skin Cancer. <i>Journal of Investigative Dermatology</i> , 2003, 120, 1087-1093.	0.3	281
21	Reclassification of 300 Primary Cutaneous B-Cell Lymphomas According to the New WHOâ€EORTC Classification for Cutaneous Lymphomas: Comparison With Previous Classifications and Identification of Prognostic Markers. <i>Journal of Clinical Oncology</i> , 2007, 25, 1581-1587.	0.8	278
22	Follicular Mycosis Fungoides, a Distinct Disease Entity With or Without Associated Follicular Mucinosis. <i>Archives of Dermatology</i> , 2002, 138, 191-8.	1.7	277
23	CAMPATH-1H monoclonal antibody in therapy for previously treated low-grade non-Hodgkin's lymphomas: a phase II multicenter study. European Study Group of CAMPATH-1H Treatment in Low-Grade Non-Hodgkin's Lymphoma.. <i>Journal of Clinical Oncology</i> , 1998, 16, 3257-3263.	0.8	272
24	Distinct types of primary cutaneous large B-cell lymphoma identified by gene expression profiling. <i>Blood</i> , 2005, 105, 3671-3678.	0.6	266
25	Activation of the <i>c-myc</i> Oncogene in a Precursor-B-Cell Blast Crisis of Follicular Lymphoma, Presenting as Composite Lymphoma. <i>New England Journal of Medicine</i> , 1988, 318, 1373-1378.	13.9	254
26	Prognostic Factors in Primary Cutaneous Large B-Cell Lymphomas: A European Multicenter Study. <i>Journal of Clinical Oncology</i> , 2001, 19, 3602-3610.	0.8	251
27	KIR-ligand incompatibility in the graft-versus-host direction improves outcomes after umbilical cord blood transplantation for acute leukemia. <i>Leukemia</i> , 2009, 23, 492-500.	3.3	236
28	Prevention of interleukin-8-induced mobilization of hematopoietic progenitor cells in rhesus monkeys by inhibitory antibodies against the Metalloproteinase gelatinase B (MMP-9). <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999, 96, 10863-10868.	3.3	225
29	Epigenetic Profiling of Cutaneous T-Cell Lymphoma: Promoter Hypermethylation of Multiple Tumor Suppressor Genes Including BCL7a, PTPRG, and p73. <i>Journal of Clinical Oncology</i> , 2005, 23, 3886-3896.	0.8	224
30	Peripheral T-cell lymphomas unspecified presenting in the skin: analysis of prognostic factors in a group of 82 patients. <i>Blood</i> , 2003, 102, 2213-2219.	0.6	221
31	An Immunodeficiency Disease with <i>RAG</i> Mutations and Granulomas. <i>New England Journal of Medicine</i> , 2008, 358, 2030-2038.	13.9	219
32	Response to Interferon Alfa-2b in a Patient with Systemic Mastocytosis. <i>New England Journal of Medicine</i> , 1992, 326, 619-623.	13.9	212
33	The clinical and histological spectrum of lymphomatoid papulosis. <i>British Journal of Dermatology</i> , 1982, 107, 131-144.	1.4	210
34	Treatment of T-cell prolymphocytic leukemia with human CD52 antibody.. <i>Journal of Clinical Oncology</i> , 1997, 15, 2667-2672.	0.8	202
35	The Polycomb group protein EZH2 is upregulated in proliferating, cultured human mantle cell lymphoma. <i>British Journal of Haematology</i> , 2001, 112, 950-958.	1.2	200
36	Mixed T cell receptor dimers harbor potentially harmful neoreactivity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010, 107, 10972-10977.	3.3	196

#	ARTICLE	IF	CITATIONS
37	Total skin electron radiation in the management of mycosis fungoides: Consensus of the European Organization for Research and Treatment of Cancer (EORTC) Cutaneous Lymphoma Project Group. <i>Journal of the American Academy of Dermatology</i> , 2002, 47, 364-370.	0.6	194
38	CD56+ hematological neoplasms presenting in the skin: a retrospective analysis of 23 new cases and 130 cases from the literature. <i>Annals of Oncology</i> , 2004, 15, 1097-1108.	0.6	191
39	Oncogenomic analysis of mycosis fungoides reveals major differences with Sézary syndrome. <i>Blood</i> , 2009, 113, 127-136.	0.6	188
40	Prognostic factors in transformed mycosis fungoides: a retrospective analysis of 100 cases. <i>Blood</i> , 2012, 119, 1643-1649.	0.6	186
41	Differences in clinical behaviour and immunophenotype between primary cutaneous and primary nodal anaplastic large cell lymphoma of T-cell or null cell phenotype. <i>Histopathology</i> , 1993, 23, 127-135.	1.6	179
42	Primary cutaneous large cell lymphomas of follicular center cell origin. <i>Journal of the American Academy of Dermatology</i> , 1987, 16, 518-526.	0.6	177
43	Long-term safety aspects of systemic therapy with fumaric acid esters in severe psoriasis. <i>British Journal of Dermatology</i> , 2003, 149, 363-369.	1.4	177
44	Novel and Highly Recurrent Chromosomal Alterations in Sézary Syndrome. <i>Cancer Research</i> , 2008, 68, 2689-2698.	0.4	176
45	Indolent CD8-positive Lymphoid Proliferation of the Ear. <i>American Journal of Surgical Pathology</i> , 2007, 31, 1887-1892.	2.1	175
46	Primary cutaneous lymphomas: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. <i>Annals of Oncology</i> , 2013, 24, vi149-vi154.	0.6	175
47	Primary Cutaneous Marginal Zone B-Cell Lymphoma. <i>Archives of Dermatology</i> , 2005, 141, 1139.	1.7	173
48	Primary cutaneous lymphomas: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. <i>Annals of Oncology</i> , 2018, 29, iv30-iv40.	0.6	171
49	Variations in clinical presentation, frequency of hemophagocytosis and clinical behavior of intravascular lymphoma diagnosed in different geographical regions. <i>Haematologica</i> , 2007, 92, 486-492.	1.7	164
50	Increased Risk of Cancer Other Than Melanoma in CDKN2A Founder Mutation (p16-Leiden)-Positive Melanoma Families. <i>Clinical Cancer Research</i> , 2008, 14, 7151-7157.	3.2	161
51	Neutrophils are indispensable for hematopoietic stem cell mobilization induced by interleukin-8 in mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 6228-6233.	3.3	160
52	Cutaneous immunocytomas: a clinicopathologic study of 26 cases. <i>Histopathology</i> , 1993, 23, 117-125.	1.6	158
53	Granulomatous Mycosis Fungoides and Granulomatous Slack Skin. <i>Archives of Dermatology</i> , 2008, 144, 1609-17.	1.7	158
54	Intensive chemotherapy followed by allogeneic or autologous stem cell transplantation for patients with myelodysplastic syndromes (MDSs) and acute myeloid leukemia following MDS. <i>Blood</i> , 2001, 98, 2326-2331.	0.6	155

#	ARTICLE	IF	CITATIONS
55	Bcl-2, Bcl-6 and CD10 expression in cutaneous B-cell lymphoma: further support for a follicle centre cell origin and differential diagnostic significance. <i>British Journal of Dermatology</i> , 2003, 149, 1183-1191.	1.4	155
56	Aberrant Expression of the Tyrosine Kinase Receptor EphA4 and the Transcription Factor Twist in SÅ@zary Syndrome Identified by Gene Expression Analysis. <i>Cancer Research</i> , 2004, 64, 5578-5586.	0.4	155
57	Inhibition of human macrophage colony formation by interleukin 4.. <i>Journal of Experimental Medicine</i> , 1989, 170, 577-582.	4.2	154
58	Primary extranodal and nodal non-Hodgkin's lymphoma. <i>European Journal of Cancer & Clinical Oncology</i> , 1989, 25, 1203-1210.	0.9	153
59	The prognosis of patients with lymphomatoid papulosis associated with malignant lymphomas. <i>British Journal of Dermatology</i> , 1992, 126, 596-602.	1.4	149
60	Expression of Programmed Death-1 in Primary Cutaneous CD4-Positive Small/Medium-Sized Pleomorphic T-Cell Lymphoma, Cutaneous Pseudo-T-Cell Lymphoma, and Other Types of Cutaneous T-Cell Lymphoma. <i>American Journal of Surgical Pathology</i> , 2012, 36, 109-116.	2.1	148
61	Generation of leukemia-reactive cytotoxic T lymphocyte clones from the HLA-identical bone marrow donor of a patient with leukemia.. <i>Journal of Experimental Medicine</i> , 1992, 176, 1283-1289.	4.2	147
62	Low-Dose Palliative Radiotherapy for Cutaneous B- and T-Cell Lymphomas. <i>International Journal of Radiation Oncology Biology Physics</i> , 2009, 74, 154-158.	0.4	142
63	Human IP-9: A Keratinocyte-Derived High Affinity CXC-Chemokine Ligand for the IP-10/Mig Receptor (CXCR3)1. <i>Journal of Investigative Dermatology</i> , 1999, 112, 716-722.	0.3	140
64	Treatment of primary cutaneous B-cell lymphomas of follicle center cell origin: a clinical follow-up study of 55 patients treated with radiotherapy or polychemotherapy.. <i>Journal of Clinical Oncology</i> , 1996, 14, 549-555.	0.8	137
65	Gene-expression profiling and array-based CGH classify CD4+CD56+ hematodermic neoplasm and cutaneous myelomonocytic leukemia as distinct disease entities. <i>Blood</i> , 2007, 109, 1720-1727.	0.6	137
66	bcl-2 protein expression in primary cutaneous large B-cell lymphoma is site-related.. <i>Journal of Clinical Oncology</i> , 1998, 16, 2080-2085.	0.8	136
67	Transplantation of Peripheral Blood Stem Cells as Compared With Bone Marrow From HLA-Identical Siblings in Adult Patients With Acute Myeloid Leukemia and Acute Lymphoblastic Leukemia. <i>Journal of Clinical Oncology</i> , 2002, 20, 4655-4664.	0.8	136
68	Effect of Smoking and Sun on the Aging Skin. <i>Journal of Investigative Dermatology</i> , 2003, 120, 548-554.	0.3	135
69	Evolution of acquired severe aplastic anaemia to myelodysplasia and subsequent leukaemia in adults. <i>British Journal of Haematology</i> , 1988, 70, 55-62.	1.2	134
70	Megakaryoblastic Leukaemia (Acute Myelofibrosis): a Report of Three Cases. <i>British Journal of Haematology</i> , 1979, 42, 9-20.	1.2	132
71	Most Primary Cutaneous CD30-Positive Lymphoproliferative Disorders Have a CD4-Positive Cytotoxic T-Cell Phenotype. <i>Journal of Investigative Dermatology</i> , 1997, 109, 636-640.	0.3	131
72	Internalization and cell cycle-dependent killing of leukemic cells by Gemtuzumab Ozogamicin: rationale for efficacy in CD33-negative malignancies with endocytic capacity. <i>Leukemia</i> , 2004, 18, 316-325.	3.3	131

#	ARTICLE	IF	CITATIONS
73	The PROCLIP international registry of early-stage mycosis fungoides identifies substantial diagnostic delay in most patients. <i>British Journal of Dermatology</i> , 2019, 181, 350-357.	1.4	127
74	Primary and secondary cutaneous CD30(+) lymphoproliferative disorders: a report from the Dutch Cutaneous Lymphoma Group on the long-term follow-up data of 219 patients and guidelines for diagnosis and treatment. <i>Blood</i> , 2000, 95, 3653-61.	0.6	127
75	Growth inhibition of clonogenic leukemic precursor cells by minor histocompatibility antigen-specific cytotoxic T lymphocytes.. <i>Journal of Experimental Medicine</i> , 1991, 174, 27-33.	4.2	126
76	Array-Based Comparative Genomic Hybridization Analysis Reveals Recurrent Chromosomal Alterations and Prognostic Parameters in Primary Cutaneous Large B-Cell Lymphoma. <i>Journal of Clinical Oncology</i> , 2006, 24, 296-305.	0.8	125
77	Prognostic significance of CD30 (Ki-1/Ber-H2) expression in primary cutaneous large-cell lymphomas of T-cell origin. A clinicopathologic and immunohistochemical study in 20 patients. <i>American Journal of Pathology</i> , 1989, 135, 1169-78.	1.9	124
78	Autologous bone marrow transplantation in acute myeloid leukemia in first remission: results of a Dutch prospective study.. <i>Journal of Clinical Oncology</i> , 1990, 8, 287-294.	0.8	122
79	Retroviral transfer of human CD20 as a suicide gene for adoptive T-cell therapy. <i>Haematologica</i> , 2009, 94, 1316-1320.	1.7	121
80	Results of Radiotherapy in 153 Primary Cutaneous B-Cell Lymphomas Classified According to the WHO-EORTC Classification. <i>Archives of Dermatology</i> , 2007, 143, 1520.	1.7	120
81	Clinical Staging and Prognostic Factors in Folliculotropic Mycosis Fungoides. <i>JAMA Dermatology</i> , 2016, 152, 992.	2.0	119
82	DFFRY codes for a new human male-specific minor transplantation antigen involved in bone marrow graft rejection. <i>Blood</i> , 2000, 95, 1100-1105.	0.6	117
83	Classification of cutaneous T-cell lymphoma: from Alibert to WHO-EORTC. <i>Journal of Cutaneous Pathology</i> , 2006, 33, 18-26.	0.7	117
84	The CXCR3 Activating Chemokines IP-10, Mig, and IP-9 are Expressed in Allergic but not in Irritant Patch Test Reactions. <i>Journal of Investigative Dermatology</i> , 1999, 113, 574-578.	0.3	116
85	MicroRNA-21 Expression in CD4+ T Cells Is Regulated by STAT3 and Is Pathologically Involved in SÅ©zary Syndrome. <i>Journal of Investigative Dermatology</i> , 2011, 131, 762-768.	0.3	116
86	Characterization of T-Cell Subpopulations in Skin and Peripheral Blood of Patients with Cutaneous T-Cell Lymphomas and Benign Inflammatory Dermatoses. <i>Journal of Investigative Dermatology</i> , 1983, 80, 60-66.	0.3	115
87	Repeat polymorphisms in the interleukin-4 gene (IL4). <i>Nucleic Acids Research</i> , 1991, 19, 3763-3763.	6.5	115
88	Classification of primary cutaneous T-cell lymphomas. <i>Histopathology</i> , 1994, 24, 405-415.	1.6	113
89	Differential Expression of Programmed Death-1 (PD-1) in SÅ©zary Syndrome and Mycosis Fungoides. <i>Archives of Dermatology</i> , 2012, 148, 1379.	1.7	113
90	Anthracycline-based chemotherapy as primary treatment for intravascular lymphoma. <i>Annals of Oncology</i> , 2004, 15, 1215-1221.	0.6	111

#	ARTICLE	IF	CITATIONS
91	Human Mesenchymal Stem Cells Derived from Bone Marrow Display a Better Chondrogenic Differentiation Compared with Other Sources. <i>Connective Tissue Research</i> , 2007, 48, 132-140.	1.1	110
92	Allogeneic stem cell transplantation in paroxysmal nocturnal hemoglobinuria. <i>Haematologica</i> , 2012, 97, 1666-1673.	1.7	110
93	Quality of life in patients with acute myelogenous leukemia in prolonged first complete remission after bone marrow transplantation (allogeneic or autologous) or chemotherapy: a cross-sectional study of the EORTC-GIMEMA AML 8A trial. <i>Bone Marrow Transplantation</i> , 1997, 20, 307-315.	1.3	108
94	The early phase of engraftment after murine blood cell transplantation is mediated by hematopoietic stem cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998, 95, 725-729.	3.3	108
95	Primary gastrointestinal non-Hodgkin's lymphoma in a population-based registry. <i>British Journal of Cancer</i> , 1989, 60, 745-750.	2.9	106
96	Rejection of bone-marrow graft by recipient-derived cytotoxic T lymphocytes against minor histocompatibility antigens. <i>Lancet</i> , The, 1990, 335, 131-134.	6.3	105
97	Treatment of Multifocal Primary Cutaneous B-Cell Lymphoma: A Clinical Follow-Up Study of 29 Patients. <i>Journal of Clinical Oncology</i> , 1999, 17, 2471-2471.	0.8	105
98	CD8+ T Cells in Cutaneous T-Cell Lymphoma: Expression of Cytotoxic Proteins, Fas Ligand, and Killing Inhibitory Receptors and Their Relationship With Clinical Behavior. <i>Journal of Clinical Oncology</i> , 2001, 19, 4322-4329.	0.8	105
99	Molecular Cytogenetic Analysis of Chromosomal Breakpoints in the IGH, MYC, BCL6, and MALT1 Gene Loci in Primary Cutaneous B-cell Lymphomas. <i>Journal of Investigative Dermatology</i> , 2004, 123, 213-219.	0.3	105
100	Photodynamic Therapy does not Prevent Cutaneous Squamous-Cell Carcinoma in Organ-Transplant Recipients: Results of a Randomized-Controlled Trial. <i>Journal of Investigative Dermatology</i> , 2006, 126, 569-574.	0.3	105
101	Expression of Fas and Fas-ligand in primary cutaneous T-cell lymphoma (CTCL): association between lack of Fas expression and aggressive types of CTCL. <i>British Journal of Dermatology</i> , 2000, 143, 313-319.	1.4	103
102	Subsequent Squamous- and Basal-Cell Carcinomas in Kidney-Transplant Recipients After the First Skin Cancer: Cumulative Incidence and Risk Factors. <i>Transplantation</i> , 2010, 89, 1231-1238.	0.5	102
103	Effect of centre on outcome of bone-marrow transplantation for acute myeloid leukaemia. <i>Lancet</i> , The, 2000, 355, 1393-1398.	6.3	99
104	Replacement of animal-derived collagen matrix by human fibroblast-derived dermal matrix for human skin equivalent products. <i>Biomaterials</i> , 2009, 30, 71-78.	5.7	99
105	Modification of rhodamine staining allows identification of hematopoietic stem cells with preferential short-term or long-term bone marrow-repopulating ability.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995, 92, 8901-8905.	3.3	98
106	Global patterns of care in advanced stage mycosis fungoides/Sezary syndrome: a multicenter retrospective follow-up study from the Cutaneous Lymphoma International Consortium. <i>Annals of Oncology</i> , 2017, 28, 2517-2525.	0.6	98
107	Histopathologic studies in S�zary syndrome and erythrodermic mycosis fungoides: A comparison with benign forms of erythroderma. <i>Journal of the American Academy of Dermatology</i> , 1986, 15, 1217-1226.	0.6	96
108	The addition of rituximab to anthracycline-based chemotherapy significantly improves outcome in Western patients with intravascular large B-cell lymphoma. <i>British Journal of Haematology</i> , 2008, 143, 253-257.	1.2	96

#	ARTICLE	IF	CITATIONS
109	The majority of cutaneous marginal zone B-cell lymphomas expresses class-switched immunoglobulins and develops in a T-helper type 2 inflammatory environment. <i>Blood</i> , 2008, 112, 3355-3361.	0.6	92
110	Applicability and Prognostic Value of the New TNM Classification System in 135 Patients With Primary Cutaneous Anaplastic Large Cell Lymphoma. <i>Archives of Dermatology</i> , 2009, 145, 1399-404.	1.7	92
111	miRNA expression profiling of mycosis fungoides. <i>Molecular Oncology</i> , 2011, 5, 273-280.	2.1	91
112	Cutaneous pseudo-T-cell lymphomas. A clinicopathologic study of 20 patients. <i>Cancer</i> , 1992, 69, 717-724.	2.0	89
113	Identification of prognostic factors predicting outcome in Hodgkin's lymphoma patients relapsing after autologous stem cell transplantation. <i>Annals of Oncology</i> , 2013, 24, 2430-2434.	0.6	89
114	Generation of dendritic cells expressing bcr-abl from CD34-positive chronic myeloid leukemia precursor cells. <i>Human Immunology</i> , 1997, 53, 216-223.	1.2	88
115	Gemtuzumab ozogamicin (Mylotarg®) as single-agent treatment for frail patients 61 years of age and older with acute myeloid leukemia: final results of AML-15B, a phase 2 study of the European Organisation for Research and Treatment of Cancer and Gruppo Italiano Malattie Ematologiche dell'Adulto Leukemia Groups. <i>Leukemia</i> , 2005, 19, 1768-1773.	3.3	88
116	Expression profiling reveals that methylation of TIMP3 is involved in uveal melanoma development. <i>International Journal of Cancer</i> , 2003, 106, 472-479.	2.3	86
117	IgM Expression on Paraffin Sections Distinguishes Primary Cutaneous Large B-cell Lymphoma, Leg Type From Primary Cutaneous Follicle Center Lymphoma. <i>American Journal of Surgical Pathology</i> , 2010, 34, 1043-1048.	2.1	86
118	Aggressive epidermotropic cutaneous CD8 ⁺ lymphoma: a cutaneous lymphoma with distinct clinical and pathological features. Report of an EORTC Cutaneous Lymphoma Task Force Workshop. <i>Histopathology</i> , 2015, 67, 425-441.	1.6	86
119	Differentiation between lymphadenosis benigna cutis and primary cutaneous follicular center cell lymphomas a comparative clinicopathologic study of 57 patients. <i>Cancer</i> , 1990, 65, 2301-2306.	2.0	85
120	T cells recognizing leukemic CD34 ⁺ progenitor cells mediate the antileukemic effect of donor lymphocyte infusions for relapsed chronic myeloid leukemia after allogeneic stem cell transplantation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998, 95, 10152-10157.	3.3	85
121	Cyclosporin A combined with vincristine, doxorubicin and dexamethasone (VAD) compared with VAD alone in patients with advanced refractory multiple myeloma: an EORTC-HOVON randomized phase III study (06914). <i>British Journal of Haematology</i> , 2001, 115, 895-902.	1.2	84
122	Fine-Mapping Chromosomal Loss at 9p21: Correlation with Prognosis in Primary Cutaneous Diffuse Large B-Cell Lymphoma, Leg Type. <i>Journal of Investigative Dermatology</i> , 2009, 129, 1149-1155.	0.3	84
123	Mycosis fungoides-like lesions associated with phenytoin and carbamazepine therapy. <i>Journal of the American Academy of Dermatology</i> , 1991, 24, 216-220.	0.6	83
124	Expression of Cytotoxic Proteins by Neoplastic T Cells in Mycosis Fungoides Increases with Progression from Plaque Stage to Tumor Stage Disease. <i>American Journal of Pathology</i> , 1999, 154, 1203-1210.	1.9	82
125	Diagnostic Criteria in Szary's Syndrome: A Multiparameter Study of Peripheral Blood Lymphocytes in 32 Patients with Erythroderma. <i>Journal of Investigative Dermatology</i> , 1983, 81, 392-397.	0.3	81
126	The influence of HLA-matched sibling donor availability on treatment outcome for patients with AML: an analysis of the AML 8A study of the EORTC Leukaemia Cooperative Group and GIMEMA. <i>British Journal of Haematology</i> , 1998, 102, 1344-1353.	1.2	80

#	ARTICLE	IF	CITATIONS
127	Dual HLA class I and class II restricted recognition of alloreactive T lymphocytes mediated by a single T cell receptor complex. Proceedings of the National Academy of Sciences of the United States of America, 2001, 98, 6806-6811.	3.3	79
128	Interleukin 6 is a permissive factor for monocytic colony formation by human hematopoietic progenitor cells.. Journal of Experimental Medicine, 1992, 175, 1151-1154.	4.2	78
129	Evaluation of Immunophenotypic and Molecular Biomarkers for SÅ©zary Syndrome Using Standard Operating Procedures: A Multicenter Study of 59 Patients. Journal of Investigative Dermatology, 2016, 136, 1364-1372.	0.3	78
130	Response to pentostatin in hairy-cell leukemia refractory to interferon-alpha. The European Organization for Research and Treatment of Cancer Leukemia Cooperative Group.. Journal of Clinical Oncology, 1989, 7, 1533-1538.	0.8	77
131	Recognition of clonogenic leukemic cells, remission bone marrow and HLA-identical donor bone marrow by CD8+ or CD4+ minor histocompatibility antigen-specific cytotoxic T lymphocytes.. Journal of Clinical Investigation, 1995, 96, 877-883.	3.9	77
132	Efficacy of a hypofractionated schedule in electron beam radiotherapy for epithelial skin cancer: Analysis of 434 cases. Radiotherapy and Oncology, 2010, 95, 245-249.	0.3	75
133	A Meta-Analysis of Gene Expression Data Identifies a Molecular Signature Characteristic for Tumor-Stage Mycosis Fungoides. Journal of Investigative Dermatology, 2012, 132, 2050-2059.	0.3	75
134	EORTC Classification for Primary Cutaneous Lymphomas: The Best Guide to Good Clinical Management. American Journal of Dermatopathology, 1999, 21, 265-273.	0.3	75
135	Complete remission of accelerated phase chronic myeloid leukemia by treatment with leukemia-reactive cytotoxic T lymphocytes. Blood, 1999, 94, 1201-8.	0.6	75
136	Blastic NK-cell lymphomas (agranular CD4+CD56+ hematodermic neoplasms): a review. American Journal of Clinical Pathology, 2005, 123, 662-75.	0.4	75
137	Cucurbitacin I Inhibits Stat3 and Induces Apoptosis in SÅ©zary Cells. Journal of Investigative Dermatology, 2008, 128, 1691-1695.	0.3	74
138	The antileukaemic activity of 5-Aza-2 deoxycytidine (Aza-dC) in patients with relapsed and resistant leukaemia. British Journal of Cancer, 1991, 64, 144-148.	2.9	73
139	Value of allogeneic versus autologous stem cell transplantation and chemotherapy in patients with myelodysplastic syndromes and secondary acute myeloid leukemia. Final results of a prospective randomized European Intergroup Trial. Haematologica, 2010, 95, 1754-1761.	1.7	73
140	Intensified chemotherapy inspired by a pediatric regimen combined with allogeneic transplantation in adult patients with acute lymphoblastic leukemia up to the age of 40. Leukemia, 2011, 25, 1697-1703.	3.3	73
141	High Dose Cytosine Arabinoside in the Management of Refractory Acute Leukaemia. Scandinavian Journal of Haematology, 1982, 29, 141-146.	0.0	71
142	Diffuse large cell lymphomas of follicular center cell origin presenting in the skin. A clinicopathologic and immunologic study of 16 patients. American Journal of Pathology, 1987, 126, 325-333.	1.9	71
143	Demonstration of Clonal Immunoglobulin Gene Rearrangements in Cutaneous B-Cell Lymphomas and Pseudo-B-Cell Lymphomas: Differential Diagnostic and Pathogenetic Aspects. Journal of Investigative Dermatology, 1992, 99, 749-754.	0.3	69
144	Biology of IL-8-Induced Stem Cell Mobilization. Annals of the New York Academy of Sciences, 1999, 872, 71-82.	1.8	69

#	ARTICLE	IF	CITATIONS
145	Aberrant DNA Methylation in Cutaneous Malignancies. <i>Seminars in Oncology</i> , 2005, 32, 479-487.	0.8	69
146	High prevalence of MYD88 and CD79B mutations in intravascular large B-cell lymphoma. <i>Blood</i> , 2018, 131, 2086-2089.	0.6	69
147	Results of bone marrow examination in 275 patients with histological features that suggest an indolent type of cutaneous B-cell lymphoma. <i>British Journal of Haematology</i> , 2008, 142, 52-56.	1.2	68
148	Identification of phosphatidylinositol 4-kinase type II β as HLA class II-restricted target in graft versus leukemia reactivity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 3837-3842.	3.3	68
149	Cytotoxic T-lymphocyte (CTL) responses against acute or chronic myeloid leukemia. <i>Immunological Reviews</i> , 1997, 157, 223-230.	2.8	67
150	Epidermal Interferon- β Inducible Protein-10 (IP-10) and Monokine Induced by β -Interferon (Mig) but not IL-8 mRNA Expression is Associated with Epidermotropism in Cutaneous T Cell Lymphomas. <i>Journal of Investigative Dermatology</i> , 1998, 111, 222-226.	0.3	67
151	VH Gene Analysis of Primary Cutaneous B-Cell Lymphomas: Evidence for Ongoing Somatic Hypermutation and Isotype Switching. <i>Blood</i> , 1998, 92, 3857-3864.	0.6	67
152	Chromosomal Aberration Patterns Differ in Subtypes of Primary Cutaneous B Cell Lymphomas. <i>Journal of Investigative Dermatology</i> , 2004, 122, 1495-1502.	0.3	67
153	Histopathological and immunophenotypical criteria for the diagnosis of SÄzary syndrome in differentiation from other erythrodermic skin diseases: a European Organisation for Research and Treatment of Cancer (EORTC) Cutaneous Lymphoma Task Force Study of 9. <i>British Journal of Dermatology</i> , 2015, 173, 93-105.	1.4	67
154	Leiden reconstructed human epidermal model as a tool for the evaluation of the skin corrosion and irritation potential according to the ECVAM guidelines. <i>Toxicology in Vitro</i> , 2008, 22, 1311-1320.	1.1	66
155	Combined CD8+ and CD4+ adenovirus hexon-specific T cells associated with viral clearance after stem cell transplantation as treatment for adenovirus infection. <i>Haematologica</i> , 2010, 95, 1943-1951.	1.7	66
156	Allogeneic bone marrow transplantation for acute leukemia in patients over the age of 40 years. <i>Leukemia</i> , 1997, 11, 416-419.	3.3	65
157	The stem cell mobilizing capacity of patients with acute myeloid leukemia in complete remission correlates with relapse risk: results of the EORTC-GIMEMA AML-10 trial. <i>Leukemia</i> , 2003, 17, 60-67.	3.3	65
158	Non-Mycosis Fungoides Cutaneous T-Cell Lymphomas. <i>American Journal of Clinical Pathology</i> , 2013, 139, 491-514.	0.4	64
159	Immunohistochemical Studies of Lymphadenosis Benigna Cutis Occurring in a Tattoo. <i>American Journal of Dermatopathology</i> , 1988, 10, 518-523.	0.3	63
160	The generation of dendritic-like cells with increased allostimulatory function from acute myeloid leukemia cells of various FAB subclasses. <i>Human Immunology</i> , 2000, 61, 565-574.	1.2	63
161	Incidence of cancer in kidney-transplant recipients: A long-term cohort study in a single center. <i>Cancer Epidemiology</i> , 2011, 35, 105-111.	0.8	63
162	Intensive postremission chemotherapy without maintenance therapy in adults with acute lymphoblastic leukemia. Dutch Hemato-Oncology Research Group.. <i>Journal of Clinical Oncology</i> , 1997, 15, 476-482.	0.8	62

#	ARTICLE	IF	CITATIONS
163	Cutaneous Anaplastic Large Cell Lymphoma and Peripheral T-Cell Lymphoma NOS Show Distinct Chromosomal Alterations and Differential Expression of Chemokine Receptors and Apoptosis Regulators. <i>Journal of Investigative Dermatology</i> , 2010, 130, 563-575.	0.3	62
164	HLA-DP as specific target for cellular immunotherapy in HLA class II-expressing B-cell leukemia. <i>Leukemia</i> , 2008, 22, 1387-1394.	3.3	60
165	Generation and administration of HA-1-specific T-cell lines for the treatment of patients with relapsed leukemia after allogeneic stem cell transplantation: a pilot study. <i>Haematologica</i> , 2012, 97, 1205-1208.	1.7	60
166	Mycosis Fungoides. <i>American Journal of Clinical Pathology</i> , 2013, 139, 466-490.	0.4	60
167	Immunological, cytochemical and ultrastructural studies in lymphomatoid papulosis. <i>British Journal of Dermatology</i> , 1983, 108, 381-394.	1.4	59
168	Angiotropic (intravascular) large cell lymphoma of the skin previously classified as malignant angioendotheliomatosis. <i>British Journal of Dermatology</i> , 1987, 116, 393-399.	1.4	59
169	Diagnostic and prognostic evaluation of phenotypic markers TRAF1, MUM1, BCL2 and CD15 in cutaneous CD30-positive lymphoproliferative disorders. <i>British Journal of Dermatology</i> , 2009, 161, 121-127.	1.4	59
170	Comparative value of post-remission treatment in cytogenetically normal AML subclassified by NPM1 and FLT3-ITD allelic ratio. <i>Leukemia</i> , 2017, 31, 26-33.	3.3	59
171	Direct cloning of leukemia-reactive T cells from patients treated with donor lymphocyte infusion shows a relative dominance of hematopoiesis-restricted minor histocompatibility antigen HA-1 and HA-2 specific T cells. <i>Leukemia</i> , 2004, 18, 798-808.	3.3	58
172	Oral valganciclovir as pre-emptive therapy has similar efficacy on cytomegalovirus DNA load reduction as intravenous ganciclovir in allogeneic stem cell transplantation recipients. <i>Bone Marrow Transplantation</i> , 2006, 37, 693-698.	1.3	58
173	Primary cutaneous lymphoma: recommendations for clinical trial design and staging update from the ISCL, USCLC, and EORTC. <i>Blood</i> , 2022, 140, 419-437.	0.6	58
174	Chemotherapy only compared to chemotherapy followed by transplantation in high risk myelodysplastic syndrome and secondary acute myeloid leukemia; two parallel studies adjusted for various prognostic factors. <i>Leukemia</i> , 2002, 16, 1615-1621.	3.3	57
175	Long-term follow-up of myeloablative allogeneic stem cell transplantation using Campath α in the bag α as T-cell depletion: the Leiden experience. <i>Bone Marrow Transplantation</i> , 2006, 37, 1129-1134.	1.3	57
176	A genomic and expression study of AP α in primary cutaneous T α cell lymphoma: evidence for dysregulated expression of JUNB and JUND in MF and SS. <i>Journal of Cutaneous Pathology</i> , 2008, 35, 899-910.	0.7	57
177	Langerhans cell histiocytosis first presenting in the skin in adults: frequent association with a second haematological malignancy. <i>British Journal of Dermatology</i> , 2012, 167, 1287-1294.	1.4	57
178	Effect of subcutaneously administered human recombinant erythropoietin on erythropoiesis in patients with myelodysplasia. <i>British Journal of Haematology</i> , 1991, 78, 488-493.	1.2	56
179	Chemokine IP-10 expression in cultured human keratinocytes. <i>Archives of Dermatological Research</i> , 1998, 290, 335-341.	1.1	56
180	Lymphomatoid papulosis in children: a study of 10 children registered by the Dutch Cutaneous Lymphoma Working Group. <i>British Journal of Dermatology</i> , 2001, 144, 351-354.	1.4	56

#	ARTICLE	IF	CITATIONS
181	Lymphocytic infiltration of the skin (Jessner): a T-cell lymphoproliferative disease. <i>British Journal of Dermatology</i> , 1984, 110, 523-529.	1.4	56
182	Serpina1 is a potent inhibitor of IL-8-induced hematopoietic stem cell mobilization. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 1469-1474.	3.3	56
183	Genomic analysis reveals recurrent deletion of JAK-STAT signaling inhibitors <i>HNRNPK</i> and <i>SOCS1</i> in mycosis fungoides. <i>Genes Chromosomes and Cancer</i> , 2018, 57, 653-664.	1.5	56
184	Immunophenotyping and gene rearrangement analysis provide additional criteria to differentiate between cutaneous T-cell lymphomas and pseudo-T-cell lymphomas. <i>American Journal of Pathology</i> , 1997, 150, 1941-9.	1.9	56
185	Multicenter randomized phase II trial of idarubicin vs mitoxantrone, combined with VP-16 and cytarabine for induction/consolidation therapy, followed by a feasibility study of autologous peripheral blood stem cell transplantation in elderly patients with acute myeloid leukemia. <i>Leukemia</i> , 1999, 13, 843-849.	3.3	55
186	Primary cutaneous B-cell lymphoma: classification and treatment. <i>Current Opinion in Oncology</i> , 2006, 18, 425-431.	1.1	55
187	Cutaneous B-Cell Lymphoproliferative Disorders. <i>American Journal of Clinical Pathology</i> , 2013, 139, 515-535.	0.4	55
188	Clinical and histologic differentiation between lymphomatoid papulosis and pityriasis lichenoides. <i>Journal of the American Academy of Dermatology</i> , 1985, 13, 418-428.	0.6	54
189	Nuclear Factor- κ B Pathway-Activating Gene Aberrancies in Primary Cutaneous Large B-Cell Lymphoma, Leg Type. <i>Journal of Investigative Dermatology</i> , 2014, 134, 290-292.	0.3	54
190	Mimics of Cutaneous Lymphoma. <i>American Journal of Clinical Pathology</i> , 2013, 139, 536-551.	0.4	53
191	HLA-B8 and HLA-A3 coexpressed with HLA-B8 are associated with a reduced risk of the development of chronic myeloid leukemia. The Chronic Leukemia Working Party of the EBMT. <i>Blood</i> , 1999, 93, 3863-5.	0.6	53
192	Quantitative skin cultures of <i>Pityrosporum</i> yeasts in patients seropositive for the human immunodeficiency virus with and without seborrheic dermatitis. <i>Journal of the American Academy of Dermatology</i> , 1992, 27, 37-39.	0.6	52
193	Primary cutaneous follicle center lymphoma and primary cutaneous large B-cell lymphoma, leg type, are both targeted by aberrant somatic hypermutation but demonstrate differential expression of AID. <i>Blood</i> , 2006, 107, 4926-4929.	0.6	51
194	Studies on clonality by PCR analysis of the PGK-1 gene. <i>Nucleic Acids Research</i> , 1991, 19, 2794-2794.	6.5	50
195	Monitoring of engraftment and progression of acute lymphoblastic leukemia in individual NOD/SCID mice. <i>Experimental Hematology</i> , 2001, 29, 322-329.	0.2	49
196	The applicability and prognostic value of the new TNM classification system for primary cutaneous lymphomas other than mycosis fungoides and SÅ©zary syndrome: results on a large cohort of primary cutaneous B-cell lymphomas and comparison with the system used by the Dutch Cutaneous Lymphoma Group. <i>British Journal of Dermatology</i> , 2007, 157, 1205-1211.	1.4	49
197	A randomized phase-I/II multicenter study of recombinant human granulocyte-macrophage colony-stimulating factor (GM-CSF) therapy for patients with myelodysplastic syndromes and a relatively low risk of acute leukemia. <i>Annals of Hematology</i> , 1992, 64, 173-180.	0.8	48
198	Problems and possible solutions in finding an unrelated bone marrow donor. Results of consecutive searches for 240 Dutch patients. <i>Bone Marrow Transplantation</i> , 1997, 20, 1011-1017.	1.3	48

#	ARTICLE	IF	CITATIONS
199	Cutaneous T-Cell Lymphoma: Epidemiology, Etiology, and Classification. <i>Leukemia and Lymphoma</i> , 2003, 44, S49-S54.	0.6	48
200	Normalized ultraviolet (UV) induction of Langerhans cell depletion and neutrophil infiltrates after artificial UVB hardening of patients with polymorphic light eruption.. <i>British Journal of Dermatology</i> , 2005, 152, 1268-1274.	1.4	48
201	Expression of B-cell transcription factors in primary cutaneous B-cell lymphoma. <i>Modern Pathology</i> , 2006, 19, 1270-1276.	2.9	48
202	Phase I/II feasibility study evaluating the generation of leukemia-reactive cytotoxic T lymphocyte lines for treatment of patients with relapsed leukemia after allogeneic stem cell transplantation. <i>Haematologica</i> , 2007, 92, 72-80.	1.7	48
203	Skin diseases among schoolchildren in Ghana, Gabon, and Rwanda. <i>International Journal of Dermatology</i> , 2013, 52, 589-600.	0.5	48
204	A novel splice variant of the Fas gene in patients with cutaneous T-cell lymphoma. <i>Cancer Research</i> , 2002, 62, 5389-92.	0.4	48
205	Primary cutaneous anaplastic large cell lymphoma shows a distinct miR expression profile and reveals differences from tumor-stage mycosis fungoides. <i>Experimental Dermatology</i> , 2012, 21, 632-634.	1.4	47
206	Stage Is a Better Prognostic Indicator than Morphologic Subtype in Primary Noncutaneous T-Cell Lymphoma. <i>American Journal of Clinical Pathology</i> , 1990, 93, 49-57.	0.4	46
207	Diagnostic Value of T-Cell Receptor Beta Gene Rearrangement Analysis on Peripheral Blood Lymphocytes of Patients with Erythroderma. <i>Journal of Investigative Dermatology</i> , 1991, 97, 782-786.	0.3	46
208	Effects of thrombopoietin on the proliferation and differentiation of primitive and mature haemopoietic progenitor cells in cord blood. <i>British Journal of Haematology</i> , 1998, 101, 425-435.	1.2	46
209	Mantle cell lymphoma proliferates upon IL-10 in the CD40 system. <i>Leukemia</i> , 2000, 14, 1483-1489.	3.3	46
210	Expression and induction of costimulatory and adhesion molecules on acute myeloid leukemic cells Implications for adoptive immunotherapy. <i>Experimental Hematology</i> , 2000, 28, 161-168.	0.2	46
211	Minor histocompatibility antigens as targets of graft-versus-leukemia reactions. <i>Current Opinion in Hematology</i> , 2002, 9, 497-502.	1.2	46
212	Cost analysis of HLA-identical sibling and voluntary unrelated allogeneic bone marrow and peripheral blood stem cell transplantation in adults with acute myelocytic leukaemia or acute lymphoblastic leukaemia. <i>Bone Marrow Transplantation</i> , 2002, 30, 243-251.	1.3	46
213	An <i>in vitro</i> three-dimensional model of primary human cutaneous squamous cell carcinoma. <i>Experimental Dermatology</i> , 2009, 18, 849-856.	1.4	46
214	Epigenomic Analysis of SÃ©zary Syndrome Defines Patterns of Aberrant DNA Methylation and Identifies Diagnostic Markers. <i>Journal of Investigative Dermatology</i> , 2016, 136, 1876-1884.	0.3	46
215	Sustained engraftment of mice transplanted with IL-1-primed blood-derived stem cells. <i>Journal of Immunology</i> , 1992, 148, 417-21.	0.4	46
216	Lymphomatoid papulosis and Hodgkin's disease: Are they related?. <i>Archives of Dermatological Research</i> , 1983, 275, 159-167.	1.1	45

#	ARTICLE	IF	CITATIONS
217	Respiratory failure elicited by streptococcal septicaemia in patients treated with cytosine arabinoside, and its prevention by penicillin. <i>Infection</i> , 1990, 18, 131-137.	2.3	45
218	Cutaneous pseudolymphomas: classification and differential diagnosis. <i>Seminars in Dermatology</i> , 1994, 13, 187-96.	0.6	45
219	UTY gene codes for an HLA-B60-restricted human male-specific minor histocompatibility antigen involved in stem cell graft rejection: characterization of the critical polymorphic amino acid residues for T-cell recognition. <i>Blood</i> , 2000, 96, 3126-32.	0.6	45
220	A randomized comparison of maintenance treatment with androgens, immunotherapy, and chemotherapy in adult acute myelogenous leukemia: A leukemia-lymphoma group trial of the EORTC. <i>Cancer</i> , 1986, 58, 617-623.	2.0	44
221	Immunohistochemical studies in lymphocytic infiltration of the skin (Jessner) and discoid lupus erythematosus. <i>Journal of the American Academy of Dermatology</i> , 1984, 11, 832-840.	0.6	43
222	Pulmonary complications of cytosine-arabioside therapy: radiographic findings. <i>American Journal of Roentgenology</i> , 1987, 149, 23-27.	1.0	43
223	The biological activities of interleukin-1. <i>Blut</i> , 1989, 59, 147-156.	1.2	43
224	Characterization of the dermal infiltrates in Jessner's lymphocytic infiltrate of the skin, polymorphous light eruption and cutaneous lupus erythematosus: differential diagnostic and pathogenetic aspects. <i>Journal of Cutaneous Pathology</i> , 1990, 17, 2-8.	0.7	43
225	Recommendations for treatment in folliculotropic mycosis fungoides: report of the Dutch Cutaneous Lymphoma Group. <i>British Journal of Dermatology</i> , 2017, 177, 223-228.	1.4	43
226	Cytokine production by the bladder carcinoma cell line 5637: rapid analysis of mRNA expression levels using a cDNA-PCR procedure. <i>Lymphokine and Cytokine Research</i> , 1991, 10, 231-5.	0.7	43
227	Interleukin 1 and poly(rI).poly(rC) induce production of granulocyte CSF, macrophage CSF, and granulocyte-macrophage CSF by human endothelial cells. <i>Experimental Hematology</i> , 1989, 17, 229-34.	0.2	43
228	CD30 expression in normal and neoplastic lymphoid tissue: biological aspects and clinical implications. <i>Leukemia</i> , 1995, 9, 1620-7.	3.3	43
229	A histologic study of lymph nodes from patients with the S�zary syndrome. <i>Cancer</i> , 1986, 57, 2375-2380.	2.0	42
230	Lymphomatoid papulosis with a natural killer-cell phenotype.. <i>British Journal of Dermatology</i> , 2001, 145, 318-322.	1.4	42
231	Deep-Sequencing Analysis Reveals that the miR-199a2/214 Cluster within DN3os Represents the Vast Majority of Aberrantly Expressed MicroRNAs in S�zary Syndrome. <i>Journal of Investigative Dermatology</i> , 2012, 132, 1520-1522.	0.3	42
232	Methotrexate-associated B-cell Lymphoproliferative Disorders Presenting in the Skin. <i>American Journal of Surgical Pathology</i> , 2014, 38, 999-1006.	2.1	42
233	High Incidence and Clinical Significance of MYC Rearrangements in Primary Cutaneous Diffuse Large B-Cell Lymphoma, Leg Type. <i>American Journal of Surgical Pathology</i> , 2018, 42, 1488-1494.	2.1	42
234	Analysis of Major Histocompatibility Antigens and the Mononuclear Cell Infiltrate in Halo Nevi. <i>Journal of Investigative Dermatology</i> , 1985, 85, 25-29.	0.3	41

#	ARTICLE	IF	CITATIONS
235	Myeloablative T cell-depleted alloSCT with early sequential prophylactic donor lymphocyte infusion is an efficient and safe post-remission treatment for adult ALL. Bone Marrow Transplantation, 2014, 49, 287-291.	1.3	41
236	TCL1 and CLA expression in agranular CD4/CD56 hematodermic neoplasms (blastic NK-cell lymphomas) and leukemia cutis. American Journal of Clinical Pathology, 2004, 122, 307-13.	0.4	41
237	Alopecia Mucinososa Progressing into Mycosis Fungoides. American Journal of Dermatopathology, 1988, 10, 478-486.	0.3	40
238	The presence of an HLA-identical sibling donor has no impact on outcome of patients with high-risk MDS or secondary AML (sAML) treated with intensive chemotherapy followed by transplantation: results of a prospective study of the EORTC, EBMT, SAKK and GIMEMA Leukemia Groups (EORTC study) Tj ETQq0 00rgBT /Overlock 10	3.3	40
239	The Role of Interleukin-1 in Hematopoiesis. Acta Haematologica, 1991, 86, 148-154.	0.7	39
240	SUCCESSFUL OUTCOME WITH A QUINTUPLE APPROACH OF POSTTRANSPLANT LYMPHOPROLIFERATIVE DISORDER. Transplantation, 2001, 71, 47-52.	0.5	39
241	More epidermal p53 patches adjacent to skin carcinomas in renal transplant recipients than in immunocompetent patients: the role of azathioprine. Experimental Dermatology, 2008, 17, 349-355.	1.4	39
242	Treatment of early-stage mycosis fungoides: results from the PROspective Cutaneous Lymphoma International Prognostic Index (PROCLIPI) study*. British Journal of Dermatology, 2021, 184, 722-730.	1.4	39
243	Diffuse cutaneous mastocytosis with multiple cutaneous mastocytomas.. British Journal of Dermatology, 1980, 102, 601-607.	1.4	38
244	The cell population of cutaneous B-cell lymphomas. British Journal of Dermatology, 1983, 108, 395-409.	1.4	38
245	Peripheral T-cell lymphomas: a clinicopathological and immunological study of 10 cases. Histopathology, 1986, 10, 235-249.	1.6	38
246	Interleukin-4. Blut, 1990, 60, 269-274.	1.2	38
247	Minor histocompatibility antigen-specific cytotoxic T cell lines, capable of lysing human hematopoietic progenitor cells, can be generated in vitro by stimulation with HLA-identical bone marrow cells.. Journal of Experimental Medicine, 1991, 173, 101-109.	4.2	38
248	Assessment of telomere length in hematopoietic interphase cells using in situ hybridization and digital fluorescence microscopy. Cytometry, 1998, 32, 163-169.	1.8	38
249	Frequency of central nervous system involvement in primary cutaneous B-cell lymphoma. Cancer, 2000, 89, 913-919.	2.0	38
250	Differential Expression of Thymus and Activation Regulated Chemokine and Its Receptor CCR4 in Nodal and Cutaneous Anaplastic Large-Cell Lymphomas and Hodgkin's Disease. Modern Pathology, 2002, 15, 838-844.	2.9	38
251	Low-dose cytosine arabinoside (LD-AraC) vs LD-AraC plus granulocyte/macrophage colony stimulating factor vs LD-AraC plus Interleukin-3 for myelodysplastic syndrome patients with a high risk of developing acute leukemia: final results of a randomized phase III study (06903) of the EORTC Leukemia Cooperative Group. Leukemia, 2005, 19, 1929-1933.	3.3	38
252	Differential expression of the HECA-452 antigen (cutaneous lymphocyte associated antigen, CLA) in cutaneous and non-cutaneous T-cell lymphomas. Histopathology, 2007, 21, 59-64.	1.6	38

#	ARTICLE	IF	CITATIONS
253	Genetic engineering of virus-specific T cells with T-cell receptors recognizing minor histocompatibility antigens for clinical application. <i>Haematologica</i> , 2008, 93, 1535-1543.	1.7	38
254	High-dose cytosine arabinoside: pharmacological and clinical aspects. <i>Blut</i> , 1988, 56, 1-11.	1.2	37
255	Pentostatin in T-cell malignancies – a phase II trial of the EORTC. <i>Annals of Oncology</i> , 1999, 10, 1493-1498.	0.6	37
256	Primary cutaneous lymphomas: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. <i>Annals of Oncology</i> , 2010, 21, v177-v180.	0.6	37
257	Autocrine IL-21 Stimulation Is Involved in the Maintenance of Constitutive STAT3 Activation in SÅ©zary Syndrome. <i>Journal of Investigative Dermatology</i> , 2012, 132, 440-447.	0.3	37
258	Multiple minor histocompatibility antigen-specific cytotoxic T lymphocyte clones can be generated during graft rejection after HLA-identical bone marrow transplantation. <i>Bone Marrow Transplantation</i> , 1995, 16, 125-32.	1.3	37
259	Expression of co-stimulatory and adhesion molecules and chemokine or apoptosis receptors on acute myeloid leukaemia: high CD40 and CD11a expression correlates with poor prognosis. <i>British Journal of Haematology</i> , 2001, 115, 298-308.	1.2	36
260	DFFRY codes for a new human male-specific minor transplantation antigen involved in bone marrow graft rejection. <i>Blood</i> , 2000, 95, 1100-5.	0.6	36
261	Leucostasis, an underestimated cause of death in leukaemia. <i>Blut</i> , 1988, 56, 39-44.	1.2	35
262	Generation of leukemia-reactive cytotoxic T lymphocytes from HLA-identical donors of patients with chronic myeloid leukemia using modifications of a limiting dilution assay. <i>Bone Marrow Transplantation</i> , 1998, 21, 553-560.	1.3	35
263	Site-Specific Expression of Polycomb-Group Genes Encoding the HPC-HPH/PRC1 Complex in Clinically Defined Primary Nodal and Cutaneous Large B-Cell Lymphomas. <i>American Journal of Pathology</i> , 2004, 164, 533-542.	1.9	35
264	Reticulum cell sarcomas (large cell lymphomas) presenting in the skin. High frequency of true histiocytic lymphoma. <i>Cancer</i> , 1982, 50, 1367-1379.	2.0	34
265	In vitro-induced resistance to the deoxycytidine analogues cytarabine (AraC) and 5-aza-2â€²-deoxycytidine (DAC) in a rat model for acute myeloid leukemia is mediated by mutations in the deoxycytidine kinase (dck) gene. <i>Annals of Hematology</i> , 1995, 71, 41-47.	0.8	34
266	EBV-positive cutaneous B-cell lymphoproliferative disease after imatinib mesylate. <i>Blood</i> , 2003, 102, 4243-4243.	0.6	34
267	Histiocytic sarcoma (true histiocytic lymphoma): a clinicopathological study of 20 cases. <i>Histopathology</i> , 1984, 8, 105-123.	1.6	33
268	Are most primary cutaneous B-cell lymphomas 'marginal cell lymphomas'? <i>British Journal of Dermatology</i> , 1995, 133, 950-952.	1.4	33
269	Differential involvement of caspases in apoptosis of myeloid leukemic cells induced by chemotherapy versus growth factor withdrawal. <i>FEBS Letters</i> , 1997, 409, 207-210.	1.3	33
270	The number of nucleated cells reflects the hematopoietic content of umbilical cord blood for transplantation. <i>Bone Marrow Transplantation</i> , 1999, 24, 965-970.	1.3	33

#	ARTICLE	IF	CITATIONS
271	Primary cutaneous CD30-positive lymphoproliferative disorders. <i>Hematology/Oncology Clinics of North America</i> , 2003, 17, 1319-1332.	0.9	33
272	Increased Risk of Squamous-Cell Carcinoma in Simultaneous Pancreas Kidney Transplant Recipients Compared with Kidney Transplant Recipients. <i>Journal of Investigative Dermatology</i> , 2009, 129, 2886-2894.	0.3	33
273	Human growth factor (huGRO), interleukin-8 (IL-8) and interferon- γ -inducible protein (γ -IP-10) gene expression in cultured normal human keratinocytes. <i>Archives of Dermatological Research</i> , 1994, 286, 471-475.	1.1	32
274	Intermediate and high-dose ARA-C and m-AMSA (or daunorubicin) as remission and consolidation treatment for patients with relapsed acute leukaemia and lymphoblastic non-Hodgkin lymphoma. <i>Scandinavian Journal of Haematology</i> , 1985, 34, 83-87.	0.0	32
275	Proliferation of B cell malignancies in all stages of differentiation upon stimulation in the 'CD40 system'. <i>Leukemia</i> , 1996, 10, 488-93.	3.3	32
276	Primary Plasma Cell Leukaemia. <i>Acta Haematologica</i> , 1983, 70, 122-129.	0.7	31
277	The many faces of lymphocytoma cutis. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2003, 17, 3-6.	1.3	31
278	Frequency and prognosis of associated malignancies in 504 patients with lymphomatoid papulosis. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2020, 34, 260-266.	1.3	31
279	T cells selectively infiltrate bone marrow areas with residual haemopoiesis of patients with acquired aplastic anaemia. <i>British Journal of Haematology</i> , 1997, 99, 517-519.	1.2	30
280	High susceptibility of human leukemic cells to Fas-induced apoptosis is restricted to G1 phase of the cell cycle and can be increased by interferon treatment. <i>Leukemia</i> , 2003, 17, 576-584.	3.3	30
281	<i>Ex Vivo</i> Expansion and Engraftment Potential of Cord Blood-Derived CD34 ⁺ Cells in NOD/SCID Mice. <i>Annals of the New York Academy of Sciences</i> , 2001, 938, 9-17.	1.8	30
282	Cutaneous Gamma/Delta T-cell Lymphoma During Treatment with Etanercept for Rheumatoid Arthritis. <i>Acta Dermato-Venereologica</i> , 2009, 89, 653-654.	0.6	30
283	Recommendations for treatment of lymphomatoid papulosis with methotrexate: a report from the Dutch Cutaneous Lymphoma Group. <i>British Journal of Dermatology</i> , 2015, 173, 1319-1322.	1.4	30
284	Cutaneous lymphomas with a panniculitic presentation. <i>Seminars in Diagnostic Pathology</i> , 2017, 34, 36-43.	1.0	30
285	EORTC Classification for primary cutaneous lymphomas: A comparison with the R.E.A.L. Classification and the proposed WHO Classification. <i>Annals of Oncology</i> , 2000, 11, S11-S15.	0.6	29
286	Comparison of allogeneic T cell-depleted peripheral blood stem cell and bone marrow transplantation: effect of stem cell source on short- and long-term outcome. <i>Bone Marrow Transplantation</i> , 2001, 27, 1053-1058.	1.3	29
287	Functional characterization of cancer-associated fibroblasts of human cutaneous squamous cell carcinoma. <i>Experimental Dermatology</i> , 2011, 20, 737-742.	1.4	29
288	Production of tumor necrosis factor-alpha by normal and malignant B lymphocytes in response to interferon-alpha, interferon-gamma and interleukin-4. <i>Leukemia</i> , 1992, 6, 116-9.	3.3	29

#	ARTICLE	IF	CITATIONS
289	Prevention of viridans-group streptococcal septicemia in oncohematologic patients: A controlled comparative study on the effect of penicillin G and cotrimoxazole. <i>Annals of Hematology</i> , 1992, 64, 260-265.	0.8	28
290	Generation of B-cell chronic lymphocytic leukemia (B-CLL)-reactive T-cell lines and clones from HLA class I-matched donors using modified B-CLL cells as stimulators: implications for adoptive immunotherapy. <i>Leukemia</i> , 2004, 18, 1278-1287.	3.3	28
291	From sporadic atypical nevi to familial melanoma: Risk analysis for melanoma in sporadic atypical nevus patients. <i>Journal of the American Academy of Dermatology</i> , 2007, 56, 748-752.	0.6	28
292	Prevalence and causative fungal species of tinea capitis among schoolchildren in Gabon. <i>Mycoses</i> , 2011, 54, e354-9.	1.8	28
293	Cytomegalovirus antigenemia assay or PCR can be used to monitor ganciclovir treatment in bone marrow transplant recipients. <i>Bone Marrow Transplantation</i> , 1992, 9, 247-53.	1.3	28
294	Improved survival of lethally irradiated recipient mice transplanted with circulating progenitor cells mobilized by IL-8 after pretreatment with stem cell factor. <i>Experimental Hematology</i> , 1996, 24, 1387-93.	0.2	28
295	<i>Mycosis fungoides</i> simulating acanthosis nigricans. <i>American Journal of Dermatopathology</i> , 1985, 7, 367-372.	0.3	27
296	Neutropenic enterocolitis following treatment with cytosine arabinoside-containing regimens for hematological malignancies: A potentiating role for amsacrine. <i>Annals of Hematology</i> , 1991, 62, 129-134.	0.8	27
297	Increased risk of second cancers in managing Hodgkin's disease: the 20-year Leiden experience. <i>Annals of Hematology</i> , 1992, 65, 213-218.	0.8	27
298	The occurrence of residual or recurrent squamous cell carcinomas in organ transplant recipients after curettage and electrodesiccation. <i>British Journal of Dermatology</i> , 2006, 154, 493-497.	1.4	27
299	Identification of selectively expressed genes and antigens in CTCL. <i>Experimental Dermatology</i> , 2008, 17, 324-334.	1.4	27
300	Trends of skin diseases in organ-transplant recipients transplanted between 1966 and 2006: a cohort study with follow-up between 1994 and 2006. <i>British Journal of Dermatology</i> , 2010, 162, 390-396.	1.4	27
301	TOX expression in cutaneous T-cell lymphomas: an adjunctive diagnostic marker that is not tumour specific and not restricted to the CD4 ⁺ CD8 ⁺ phenotype. <i>British Journal of Dermatology</i> , 2016, 175, 382-386.	1.4	27
302	Side effects of intermediate- and high-dose cytosine arabinoside in the treatment of refractory or relapsed acute leukaemia and non-Hodgkin's lymphoma. <i>Netherlands Journal of Medicine</i> , 1987, 30, 64-74.	0.6	27
303	HLA-DR4 is associated with a diminished risk of the development of chronic myeloid leukemia (CML). <i>Leukemia</i> , 2000, 14, 859-862.	3.3	26
304	Minor histocompatibility antigen-specific T cells with multiple distinct specificities can be isolated by direct cloning of IFN γ -secreting T cells from patients with relapsed leukemia responding to donor lymphocyte infusion. <i>Leukemia</i> , 2005, 19, 83-90.	3.3	26
305	Human allo-reactive CD4 ⁺ T cells as strong mediators of anti-tumor immunity in NOD/scid mice engrafted with human acute lymphoblastic leukemia. <i>Leukemia</i> , 2012, 26, 312-322.	3.3	26
306	Effects of serially passaged fibroblasts on dermal and epidermal morphogenesis in human skin equivalents. <i>Biogerontology</i> , 2013, 14, 131-140.	2.0	26

#	ARTICLE	IF	CITATIONS
307	In vivo use of Campath-1G to prevent graft-versus-host disease and graft rejection after bone marrow transplantation. <i>Bone Marrow Transplantation</i> , 1992, 9, 255-61.	1.3	26
308	Primary cutaneous large B-cell lymphomas of the legs. A distinct type of cutaneous B-cell lymphoma with an intermediate prognosis. Dutch Cutaneous Lymphoma Working Group. <i>Archives of Dermatology</i> , 1996, 132, 1304-8.	1.7	26
309	Pentoxifylline and skin inflammation. <i>Clinical and Experimental Dermatology</i> , 1998, 23, 168-172.	0.6	25
310	Similar myeloid recovery despite superior overall engraftment in NOD/SCID mice after transplantation of human CD34+ cells from umbilical cord blood as compared to adult sources. <i>Bone Marrow Transplantation</i> , 2001, 28, 163-171.	1.3	25
311	Efficacy and toxicity of gemtuzumab ozogamicin in patients with acute myeloid leukemia. <i>European Journal of Haematology</i> , 2006, 76, 409-413.	1.1	25
312	Clinical, Histologic, and Molecular Characteristics of Anaplastic Lymphoma Kinase-positive Primary Cutaneous Anaplastic Large Cell Lymphoma. <i>American Journal of Surgical Pathology</i> , 2020, 44, 776-781.	2.1	25
313	Granzyme B-expressing peripheral T-cell lymphomas: neoplastic equivalents of activated cytotoxic T cells with preference for mucosa-associated lymphoid tissue localization. <i>Blood</i> , 1994, 84, 3785-91.	0.6	25
314	Oral etoposide in the treatment of cutaneous large-cell lymphomas. A preliminary report of four cases. <i>British Journal of Dermatology</i> , 1992, 127, 524-528.	1.4	24
315	Mosaicism of the 5q deletion as assessed by interphase FISH is a common phenomenon in MDS and restricted to myeloid cells. <i>Leukemia</i> , 1997, 11, 519-523.	3.3	24
316	Analysis of T-cell clonality in bone marrow of patients with acquired aplastic anaemia. <i>British Journal of Haematology</i> , 1997, 96, 85-91.	1.2	24
317	HLA class II restricted T-cell receptor gene transfer generates CD4+ T cells with helper activity as well as cytotoxic capacity. <i>Gene Therapy</i> , 2005, 12, 1686-1695.	2.3	24
318	Prevalence and risk factors of inflammatory acne vulgaris in rural and urban Ghanaian schoolchildren. <i>British Journal of Dermatology</i> , 2009, 161, 475-477.	1.4	24
319	Kidney Transplant Recipients with Cutaneous Squamous Cell Carcinoma Have an Increased Risk of Internal Malignancy. <i>Journal of Investigative Dermatology</i> , 2012, 132, 2176-2183.	0.3	24
320	Mycosis fungoides variantsâ€”clinicopathologic features, differential diagnosis, and treatment. <i>Seminars in Cutaneous Medicine and Surgery</i> , 2018, 37, 11-17.	1.6	24
321	Role of deoxycytidine kinase in an in vitro model for AraC- and DAC-resistance: substrate-enzyme interactions with deoxycytidine, 1-beta-D-arabinofuranosylcytosine and 5-aza-2'-deoxycytidine. <i>Leukemia</i> , 1993, 7, 1005-11.	3.3	24
322	Intermediate and high dose Ara-C and m-AMSA for remission induction and consolidation treatment of patients with acute myeloid leukemia: an EORTC leukemia cooperative group phase II study. <i>European Journal of Cancer & Clinical Oncology</i> , 1988, 24, 1721-1725.	0.9	23
323	Gastrointestinal complications of cytosine-arabioside chemotherapy: findings on plain abdominal radiographs. <i>American Journal of Roentgenology</i> , 1990, 154, 95-98.	1.0	23
324	CD44 Antibody Stimulates Adhesion of Peripheral Blood T Cells to Keratinocytes Through the Leukocyte Function-Associated Antigen-1/Intercellular Adhesion Molecule-1 Pathway. <i>Journal of Investigative Dermatology</i> , 1993, 100, 424-428.	0.3	23

#	ARTICLE	IF	CITATIONS
325	Biology and treatment of myelodysplastic syndromes – developments in the past decade. <i>Annals of Hematology</i> , 1993, 66, 107-115.	0.8	23
326	High-dose Ara-C for remission induction and consolidation of previously untreated adults with ALL or lymphoblastic lymphoma. <i>Annals of Hematology</i> , 1995, 70, 71-74.	0.8	23
327	Proliferation of precursor B-lineage acute lymphoblastic leukaemia by activating the CD40 antigen. <i>British Journal of Haematology</i> , 1996, 95, 319-326.	1.2	23
328	Classification of cutaneous lymphomas: crosstalk between pathologist and clinician. <i>Current Diagnostic Pathology</i> , 1998, 5, 23-33.	0.4	23
329	Proteolytic Enzyme Levels Are Increased During Granulocyte Colony-Stimulating Factor-Induced Hematopoietic Stem Cell Mobilization in Human Donors but Do Not Predict the Number of Mobilized Stem Cells. <i>Journal of Hematotherapy and Stem Cell Research</i> , 2002, 11, 513-521.	1.8	23
330	Expression of programmed death-1 in skin biopsies of benign inflammatory vs. lymphomatous erythroderma. <i>British Journal of Dermatology</i> , 2014, 171, 499-504.	1.4	23
331	Programmed death-1 expression in cutaneous B-cell lymphoma. <i>Journal of Cutaneous Pathology</i> , 2014, 41, 14-21.	0.7	23
332	Amsacrine with high-dose cytarabine in acute leukemia. <i>Cancer Treatment Reports</i> , 1985, 69, 1447-8.	0.5	23
333	The human interleukin-6 receptor γ chain gene is localized on chromosome 1 band q21. <i>Human Genetics</i> , 1993, 90, 542-4.	1.8	22
334	Costs of Diagnosis, Treatment, and Follow Up of Patients with Acute Myeloid Leukemia in the Netherlands. <i>Journal of Hematotherapy and Stem Cell Research</i> , 2001, 10, 187-192.	1.8	22
335	Varicella zoster virus (VZV)-related progressive outer retinal necrosis (PORN) after allogeneic stem cell transplantation. <i>Bone Marrow Transplantation</i> , 2005, 36, 467-469.	1.3	22
336	Efficacy of escalated imatinib combined with cytarabine in newly diagnosed patients with chronic myeloid leukemia. <i>Haematologica</i> , 2010, 95, 914-921.	1.7	22
337	Mechanisms underlying hematopoietic stem cell mobilization induced by the CXC chemokine interleukin-8. <i>Current Opinion in Hematology</i> , 1999, 6, 152.	1.2	22
338	Correction of neutropenia associated with chronic lymphocytic leukaemia following treatment with granulocyte-macrophage colony-stimulating factor. <i>Annals of Hematology</i> , 1991, 62, 32-34.	0.8	21
339	Umbilical cord blood contains normal frequencies of cytotoxic T-lymphocyte precursors (ctlp) and helper T-lymphocyte precursors against noninherited maternal antigens and noninherited paternal antigens. <i>Annals of Hematology</i> , 1996, 72, 260-264.	0.8	21
340	Processing of natural and recombinant CXCR3-targeting chemokines and implications for biological activity. <i>FEBS Journal</i> , 2001, 268, 4992-4999.	0.2	21
341	Non-infusional vs intravenous consolidation chemotherapy in elderly patients with acute myeloid leukemia: final results of the EORTC-GIMEMA AML-13 randomized phase III trial. <i>Leukemia</i> , 2006, 20, 1723-1730.	3.3	21
342	Reduced IL-1 α /IL-1 β ratio in ultraviolet B-exposed skin of patients with polymorphic light eruption. <i>Experimental Dermatology</i> , 2009, 18, 212-217.	1.4	21

#	ARTICLE	IF	CITATIONS
343	De novo induced mutations in the deoxycytidine kinase (dck) gene in rat leukemic clonal cell lines confer resistance to cytarabine (AraC) and 5-aza-2'-deoxycytidine (DAC). <i>Leukemia</i> , 1995, 9, 1032-8.	3.3	21
344	p53 and bcl-2 expression do not correlate with prognosis in primary cutaneous large T-cell lymphomas. <i>Journal of Cutaneous Pathology</i> , 1997, 24, 462-467.	0.7	20
345	Differences in Epstein-Barr Virus Expression Between Primary and Secondary Cutaneous Angiocentric Lymphomas. <i>Archives of Dermatology</i> , 1998, 134, 479.	1.7	20
346	Impaired Expression of CD28 on T Cells in Hairy Cell Leukemia. <i>Clinical Immunology</i> , 1999, 93, 256-262.	1.4	20
347	Intensive chemotherapy to improve outcome in patients with acute lymphoblastic leukemia over the age of 40: a phase II study for efficacy and feasibility by HOVON. <i>Leukemia</i> , 2012, 26, 1726-1729.	3.3	20
348	MicroRNA Profiling of Primary Cutaneous Large B-Cell Lymphomas. <i>PLoS ONE</i> , 2013, 8, e82471.	1.1	20
349	Plaque stage folliculotropic mycosis fungoides: histopathologic features and prognostic factors in a series of 40 patients. <i>Journal of Cutaneous Pathology</i> , 2020, 47, 241-250.	0.7	20
350	Deregulation of JAK2 signaling underlies primary cutaneous CD8 ⁺ aggressive epidermotropic cytotoxic T-cell lymphoma. <i>Haematologica</i> , 2022, 107, 702-714.	1.7	20
351	Quality of life in patients with Mycosis Fungoides and SÅžary Syndrome: a systematic review of the literature. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2021, 35, 2377-2387.	1.3	20
352	Palmar-Plantar Skin Changes and Cytarabine. <i>Annals of Internal Medicine</i> , 1985, 103, 805.	2.0	19
353	Comparison of the antileukaemic activity of 5 aza-2-deoxycytidine and arabinofuranosyl-cytosine in rats with myelocytic leukaemia. <i>British Journal of Cancer</i> , 1988, 58, 730-733.	2.9	19
354	Primary Cutaneous B-Cell Lymphomas. <i>Leukemia and Lymphoma</i> , 1994, 14, 213-218.	0.6	19
355	Relation between expression of adhesion molecules and clinical behavior in cutaneous follicle center cell lymphomas. <i>Journal of the American Academy of Dermatology</i> , 1997, 37, 34-40.	0.6	19
356	The role of cytokines and hematopoietic growth factors in the autocrine/paracrine regulation of inducible hematopoiesis. <i>Annals of Hematology</i> , 1997, 75, 27-31.	0.8	19
357	Primary cutaneous lymphomas. <i>Current Opinion in Oncology</i> , 2000, 12, 419-425.	1.1	19
358	Chemical exposures other than arsenic are probably not important risk factors for squamous cell carcinoma, basal cell carcinoma and malignant melanoma of the skin. <i>British Journal of Dermatology</i> , 2005, 152, 194-197.	1.4	19
359	Characterization of graft-versus-leukemia responses in patients treated for advanced chronic lymphocytic leukemia with donor lymphocyte infusions after in vitro T-cell depleted allogeneic stem cell transplantation following reduced-intensity conditioning. <i>Leukemia</i> , 2007, 21, 2569-2574.	3.3	19
360	Primary cutaneous lymphoma: ESMO Clinical Recommendations for diagnosis, treatment and follow-up. <i>Annals of Oncology</i> , 2009, 20, iv115-iv118.	0.6	19

#	ARTICLE	IF	CITATIONS
361	Results of induction and consolidation treatment with intermediate and high-dose cytosine arabinoside and m-AMSA of patients with poor-risk acute myelogenous leukaemia. <i>European Journal of Haematology</i> , 1988, 40, 198-204.	1.1	19
362	Dexamethasone compared to prednisolone for adults with acute lymphoblastic leukemia or lymphoblastic lymphoma: final results of the ALL-4 randomized, phase III trial of the EORTC Leukemia Group. <i>Haematologica</i> , 2010, 95, 1489-1495.	1.7	19
363	Predictive impact of allele-matching and EBMT risk score for outcome after T-cell depleted unrelated donor transplantation in poor-risk acute leukemia and myelodysplasia. <i>Leukemia</i> , 2011, 25, 1548-1554.	3.3	19
364	Recommendations for the Optimal Radiation Dose in Patients With Primary Cutaneous Anaplastic Large Cell Lymphoma: A Report of the Dutch Cutaneous Lymphoma Group. <i>International Journal of Radiation Oncology Biology Physics</i> , 2017, 99, 1279-1285.	0.4	19
365	Evaluation of treatment results in multifocal primary cutaneous anaplastic large cell lymphoma: report of the Dutch Cutaneous Lymphoma Group. <i>British Journal of Dermatology</i> , 2018, 179, 724-731.	1.4	19
366	Immunohistochemical studies using monoclonal antibodies on lymph nodes from patients with mycosis fungoides and S�azary's syndrome. <i>American Journal of Pathology</i> , 1985, 120, 46-54.	1.9	19
367	Is mycosis fungoides exacerbated by fluoxetine?. <i>Journal of the American Academy of Dermatology</i> , 1996, 35, 635-636.	0.6	18
368	Pentoxifylline inhibits human T-cell adhesion to dermal endothelial cells. <i>Archives of Dermatological Research</i> , 1997, 289, 189-193.	1.1	18
369	Long-term survival in adolescent and adult acute lymphoblastic leukemia. <i>Cancer</i> , 2006, 52, 30-34.	2.0	18
370	Involvement of caspase-8 in chemotherapy-induced apoptosis of patient derived leukemia cell lines independent of the death receptor pathway and downstream from mitochondria. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2007, 12, 181-193.	2.2	18
371	Bone marrow examination has limited value in the staging of patients with an anaplastic large cell lymphoma first presenting in the skin. Retrospective analysis of 107 patients. <i>British Journal of Dermatology</i> , 2008, 159, 1148-51.	1.4	18
372	Diagnostic and prognostic significance of <i>CDKN2A</i> / <i>CDKN2B</i> deletions in patients with transformed mycosis fungoides and primary cutaneous CD30-positive lymphoproliferative disease. <i>British Journal of Dermatology</i> , 2015, 172, 784-788.	1.4	18
373	Should we be imaging lymph nodes at initial diagnosis of early-stage mycosis fungoides? Results from the PROspective Cutaneous Lymphoma International Prognostic Index (PROCLIPI) international study*. <i>British Journal of Dermatology</i> , 2021, 184, 524-531.	1.4	18
374	Anti-LFA-1 blocking antibodies prevent mobilization of hematopoietic progenitor cells induced by interleukin-8. <i>Blood</i> , 1998, 91, 4099-105.	0.6	18
375	High incidence of alternatively spliced forms of deoxycytidine kinase in patients with resistant acute myeloid leukemia. <i>Blood</i> , 2000, 96, 1517-24.	0.6	18
376	The hematopoietic activities of interleukin-1. <i>Biotherapy</i> (Dordrecht, Netherlands), 1989, 1, 263-271.	0.7	17
377	Bone marrow transplantation or chemotherapy as post-remission treatment of adult acute myelogenous leukemia. <i>Annals of Hematology</i> , 1991, 62, 59-63.	0.8	17
378	TCR�+ cells expressing V� 9V� 2, which normally predominate the blood, are found in the spleens of patients with hairy cell leukemia. <i>Leukemia</i> , 1997, 11, 106-109.	3.3	17

#	ARTICLE	IF	CITATIONS
379	UVB-induced leucocyte trafficking in the epidermis of photosensitive lupus erythematosus patients: Normal depletion of Langerhans cells. <i>Experimental Dermatology</i> , 2005, 14, 138-142.	1.4	17
380	Similar Potential to Become Activated and Proliferate but Differential Kinetics and Profiles of Cytokine Production of Umbilical Cord Blood T Cells and Adult Blood Naive and Memory T Cells. <i>Human Immunology</i> , 2006, 67, 874-883.	1.2	17
381	Genome-wide linkage scan for atypical nevi in p16-Leiden melanoma families. <i>European Journal of Human Genetics</i> , 2008, 16, 1135-1141.	1.4	17
382	Differential expression of TOX by skin-infiltrating T cells in SÅ©zary syndrome and erythrodermic dermatitis. <i>Journal of Cutaneous Pathology</i> , 2015, 42, 604-609.	0.7	17
383	The Use of Central Pathology Review With Digital Slide Scanning in Advanced-stage Mycosis Fungoides and SÅ©zary Syndrome. <i>American Journal of Surgical Pathology</i> , 2018, 42, 726-734.	2.1	17
384	Substrate-specific deoxycytidine kinase deficiency in 1-beta-D-arabinofuranosylcytosine-resistant leukemic cells. <i>Cancer Research</i> , 1990, 50, 6515-9.	0.4	17
385	Experience with Intermediate and High Dose Cytosine Arabinoside in Refractory Acute Leukaemia. <i>Oncology Research and Treatment</i> , 1983, 6, 200-204.	0.8	16
386	Deoxycytidine kinase, thymidine kinase and cytidine deaminase and the formation of ARA-CTP in leukemic cells in different phases of the cell cycle. <i>Leukemia Research</i> , 1990, 14, 363-369.	0.4	16
387	Spectrophotometric quantification of cell-cell adherence by an enzyme-linked immuno-cell adhesion assay. <i>Journal of Immunological Methods</i> , 1990, 132, 51-56.	0.6	16
388	Increase in Ara-C sensitivity in Ara-C sensitive and -resistant leukemia by stimulation of the salvage and inhibition of the de novo pathway. <i>Annals of Hematology</i> , 1992, 65, 26-32.	0.8	16
389	Colony growth of human hematopoietic progenitor cells in the absence of serum is supported by a proteinase inhibitor identified as antileukoproteinase.. <i>Journal of Experimental Medicine</i> , 1996, 184, 1305-1312.	4.2	16
390	Molecular persistence of chronic myeloid leukemia caused by donor T cells specific for lineage-restricted maturation antigens not recognizing immature progenitor-cells. <i>Leukemia</i> , 2006, 20, 1040-1046.	3.3	16
391	Separation, enrichment, and characterization of human hematopoietic progenitor cells from umbilical cord blood. <i>Annals of Hematology</i> , 1993, 67, 231-236.	0.8	15
392	Pentoxifylline Inhibits T-Cell Adherence to Keratinocytes. <i>Journal of Investigative Dermatology</i> , 1995, 104, 1004-1007.	0.3	15
393	Incidence of mycosis fungoides and SÅ©zary syndrome in the Netherlands between 2000 and 2020. <i>British Journal of Dermatology</i> , 2021, 185, 434-435.	1.4	15
394	TREATMENT OF REFRACTORY ACUTE LEUKAEMIA WITH HIGH DOSE CYTOSINE ARABINOSIDE. <i>British Journal of Haematology</i> , 1982, 51, 497-504.	1.2	14
395	Interleukin-9 stimulates the proliferation of enriched human erythroid progenitor cells: additive effect with GM-CSF. <i>Annals of Hematology</i> , 1993, 66, 45-49.	0.8	14
396	Interleukin-10, interleukin-12, and tumor necrosis factor-Î± differentially influence the proliferation of human CD8 + and CD4 + T-cell clones. <i>Annals of Hematology</i> , 1996, 72, 245-252.	0.8	14

#	ARTICLE	IF	CITATIONS
397	IP-10 mRNA EXPRESSION IN CULTURED KERATINOCYTES IS SUPPRESSED BY INHIBITION OF PROTEIN KINASE-C AND TYROSINE KINASE AND ELEVATION OF cAMP. <i>Cytokine</i> , 1999, 11, 469-475.	1.4	14
398	MICA Gene Polymorphism is Not Associated With an Increased Risk for Skin Cancer. <i>Journal of Investigative Dermatology</i> , 2002, 118, 686-691.	0.3	14
399	Epidermal transit of replication-arrested, undifferentiated keratinocytes in UV-exposed XPC mice: An alternative to in situ apoptosis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 18980-18985.	3.3	14
400	Differential activation of the death receptor pathway in human target cells induced by cytotoxic T lymphocytes showing different kinetics of killing. <i>Haematologica</i> , 2007, 92, 1671-1678.	1.7	14
401	Results of syngeneic hematopoietic stem cell transplantation for acute leukemia: risk factors for outcomes of adults transplanted in first complete remission. <i>Haematologica</i> , 2008, 93, 834-841.	1.7	14
402	Short-term intensive treatment (V.A.A.P.) of adult acute lymphoblastic leukemia and lymphoblastic lymphoma. <i>European Journal of Haematology</i> , 2009, 41, 489-495.	1.1	14
403	Rapamycin impairs UV induction of mutant p53 overexpressing cell clusters without affecting tumor onset. <i>International Journal of Cancer</i> , 2012, 131, 1267-1276.	2.3	14
404	Whole-genome analysis uncovers recurrent IKZF1 inactivation and aberrant cell adhesion in blastic plasmacytoid dendritic cell neoplasm. <i>Genes Chromosomes and Cancer</i> , 2020, 59, 295-308.	1.5	14
405	HOXC4, HOXC5, and HOXC6 expression in primary cutaneous lymphoid lesions. High expression of HOXC5 in anaplastic large-cell lymphomas. <i>American Journal of Pathology</i> , 1997, 151, 1067-74.	1.9	14
406	Development of pulmonary leukostasis in experimental myelocytic leukemia in the Brown-Norway rat. <i>Leukemia</i> , 1992, 6, 142-9.	3.3	14
407	Experience with intermediate- and high-dose cytosine arabinoside in relapsed and refractory acute leukaemia. <i>Netherlands Journal of Medicine</i> , 1983, 26, 215-9.	0.6	14
408	Accelerated reconstitution of platelets and erythrocytes after syngeneic transplantation of bone marrow cells derived from thrombopoietin pretreated donor mice. <i>Blood</i> , 1995, 86, 3308-13.	0.6	14
409	Preliminary results with 5-aza-2'-deoxycytidine (DAC)-containing chemotherapy in patients with relapsed or refractory acute leukemia. The EORTC Leukemia Cooperative Group. <i>Leukemia</i> , 1993, 7 Suppl 1, 49-50.	3.3	14
410	Regulatory effects of T cell lymphokines on cytokine gene expression in monocytes. <i>Lymphokine and Cytokine Research</i> , 1993, 12, 457-64.	0.7	14
411	Presence of Epstein-Barr virus in extranodal T-cell lymphomas: differences in relation to site. <i>Blood</i> , 1994, 83, 1612-8.	0.6	14
412	Neurological complications in patients with cutaneous T-cell lymphoma. <i>British Journal of Dermatology</i> , 1986, 114, 359-366.	1.4	13
413	Pentostatin (2-deoxycoformycin) for the treatment of lymphoid neoplasms. <i>Cancer Treatment Reviews</i> , 1990, 17, 213-215.	3.4	13
414	Absence of mutations in the RET gene in acute myeloid leukemia. <i>Annals of Hematology</i> , 1997, 75, 87-90.	0.8	13

#	ARTICLE	IF	CITATIONS
415	Secondary T-acute lymphoblastic leukaemia mimicking blast crisis in chronic myeloid leukaemia. <i>British Journal of Haematology</i> , 1999, 106, 104-106.	1.2	13
416	Recruitment of leukemic cells from G0 phase of the cell cycle by interferons results in conversion of resistance to daunorubicin. <i>Leukemia</i> , 2003, 17, 2049-2051.	3.3	13
417	Recent advances in primary cutaneous B-cell lymphomas. <i>Current Opinion in Oncology</i> , 2014, 26, 230-236.	1.1	13
418	High numbers of mobilized CD34+ cells collected in AML in first remission are associated with high relapse risk irrespective of treatment with autologous peripheral blood SCT or autologous BMT. <i>Bone Marrow Transplantation</i> , 2015, 50, 341-347.	1.3	13
419	Quantitative electron microscopy in the early diagnosis of cutaneous T-cell lymphomas. A long-term follow-up study of 77 patients. <i>American Journal of Pathology</i> , 1986, 123, 166-73.	1.9	13
420	Abnormally activated T lymphocytes in the spleen of patients with hairy-cell leukemia. <i>Leukemia</i> , 1994, 8, 2095-101.	3.3	13
421	Percentage of S-phase cells in bone marrow aspirates, biopsy specimens and bone marrow aspirates corrected for blood dilution from patients with acute leukemia. <i>Leukemia Research</i> , 1987, 11, 209-213.	0.4	12
422	Differential Diagnosis of Cutaneous Large Cell Lymphomas Using Monoclonal Antibodies Reactive in Paraffin-Embedded Skin Biopsy Specimens. <i>American Journal of Dermatopathology</i> , 1991, 13, 342-349.	0.3	12
423	Clofarabine in combination with a standard remission induction regimen (cytosine arabinoside and) 10.784314 rgBT /Over leukemia (AML) or high-risk myelodysplastic syndrome (HR-MDS): phase I results of an ongoing phase I/II study of the leukemia groups of EORTC and GIMEMA (EORTC GIMEMA 06061/AML-14A trial). <i>Annals of Hematology</i> , 2014, 93, 965-75.	0.8	12
424	No TP63 rearrangements in a selected group of primary cutaneous CD30+ lymphoproliferative disorders with aggressive clinical course. <i>Blood</i> , 2016, 128, 141-143.	0.6	12
425	Folliculotropic mycosis fungoides presenting with a solitary lesion: Clinicopathological features and long-term follow-up data in a series of 9 cases. <i>Journal of Cutaneous Pathology</i> , 2018, 45, 122-128.	0.7	12
426	Acquired N-Linked Glycosylation Motifs in B-Cell Receptors of Primary Cutaneous B-Cell Lymphoma and the Normal B-Cell Repertoire. <i>Journal of Investigative Dermatology</i> , 2019, 139, 2195-2203.	0.3	12
427	Primary cutaneous peripheral T-cell lymphoma, not otherwise specified: results of a multicentre European Organization for Research and Treatment of Cancer (EORTC) cutaneous lymphoma taskforce study on the clinico-pathological and prognostic features. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2021, 35, 658-668.	1.3	12
428	Analysis of helper activity on pokeweed mitogen- and interleukin 2-driven immunoglobulin synthesis by neoplastic T4+ cells.. <i>Journal of Clinical Investigation</i> , 1985, 76, 2139-2143.	3.9	12
429	Cell-of-origin classification using the Hans and Lymph2Cx algorithms in primary cutaneous large B-cell lymphomas. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2022, 480, 667-675.	1.4	12
430	Clinical, histopathological and prognostic features of primary cutaneous acral CD8+ T-cell lymphoma and other dermal CD8+ cutaneous lymphoproliferations: results of an EORTC Cutaneous Lymphoma Group workshop*. <i>British Journal of Dermatology</i> , 2022, 186, 887-897.	1.4	12
431	Continuous infusion of interleukin-6 in sublethally irradiated mice accelerates platelet reconstitution and the recovery of myeloid but not of megakaryocytic progenitor cells in bone marrow. <i>Experimental Hematology</i> , 1993, 21, 1621-7.	0.2	12
432	Persistent clonal excess and skewed T-cell repertoire in T cells from patients with hairy cell leukemia. <i>Blood</i> , 1996, 87, 3795-802.	0.6	12

#	ARTICLE	IF	CITATIONS
433	Regulatory activity of neoplastic T cells in Sezary syndrome on in vitro immunoglobulin production. <i>Leukemia Research</i> , 1984, 8, 873-884.	0.4	11
434	Array-based CGH of primary cutaneous CD8+ aggressive EPIDERMOTROPIC cytotoxic T-cell lymphoma. <i>Genes Chromosomes and Cancer</i> , 2018, 57, 622-629.	1.5	11
435	VH gene analysis of primary cutaneous B-cell lymphomas: evidence for ongoing somatic hypermutation and isotype switching. <i>Blood</i> , 1998, 92, 3857-64.	0.6	11
436	The significance of Leu 8 negative T cells in lymphoid skin infiltrates: malignant transformation, selective homing or T-cell activation?. <i>British Journal of Dermatology</i> , 1990, 123, 587-593.	1.4	10
437	A simplified assay for measurement of cytosine arabinoside incorporation into DNA in Ara-C-sensitive and-resistant leukemic cells. <i>Cancer Chemotherapy and Pharmacology</i> , 1990, 27, 151-156.	1.1	10
438	Induction of ICAM-1 expression by epidermal keratinocytes via a paracrine pathway possibly involving dermal dendritic cells. <i>Archives of Dermatological Research</i> , 1992, 284, 250-252.	1.1	10
439	Multiple minor histocompatibility antigen disparities between a recipient and four HLA-identical potential sibling donors for bone marrow transplantation. <i>Human Immunology</i> , 1993, 37, 221-228.	1.2	10
440	IL-4 down-regulates IL-2-, IL-3-, and GM-CSF-induced cytokine gene expression in peripheral blood monocytes. <i>Annals of Hematology</i> , 1994, 68, 293-298.	0.8	10
441	CD30-Negative Cutaneous T-Cell Lymphomas Other than Mycosis Fungoides. <i>Surgical Pathology Clinics</i> , 2014, 7, 229-252.	0.7	10
442	Human interleukin for DA cells (HILDA) does not affect the proliferation and differentiation of hematopoietic progenitor cells in human long-term bone marrow cultures. <i>Experimental Hematology</i> , 1992, 20, 6-10.	0.2	10
443	Fournier's gangrene or fulminant necrotizing fasciitis of the scrotum and penis as a complication of granulocytopenia in a patient with acute myelogenous leukaemia (AML). <i>Netherlands Journal of Medicine</i> , 1988, 32, 235-9.	0.6	10
444	Involvement of the complement system in the pathogenesis of pulmonary leukostasis in experimental myelocytic leukemia. <i>Leukemia</i> , 1993, 7, 1608-14.	3.3	10
445	Marrow culture in diffusion chambers in rabbits: III. Effect of endotoxin and leukocyte products on cell production. <i>American Journal of Hematology</i> , 1979, 7, 33-43.	2.0	9
446	Intercellular distribution of HLA-DR and T6 antigens in the epidermis of various dermatoses. <i>British Journal of Dermatology</i> , 1985, 113, 802-804.	1.4	9
447	Order of human hematopoietic growth factor and receptor genes on the long arm of chromosome 5, as determined by fluorescence in situ hybridization. <i>Annals of Hematology</i> , 1993, 66, 15-20.	0.8	9
448	The Role of the CD40 Antigen on Malignant B Cells. <i>Leukemia and Lymphoma</i> , 1996, 22, 229-235.	0.6	9
449	Accelerated platelet reconstitution following transplantation of bone marrow cells derived from IL-6-treated donor mice. <i>Annals of Hematology</i> , 1996, 73, 239-245.	0.8	9
450	Factors influencing release of granulocyte-macrophage colony-stimulating activity from human mononuclear phagocytes*. <i>European Journal of Haematology</i> , 1988, 41, 352-358.	1.1	9

#	ARTICLE	IF	CITATIONS
451	Dietary Immunosuppressants Do Not Enhance UV-Induced Skin Carcinogenesis, and Reveal Discordance between p53-Mutant Early Clones and Carcinomas. <i>Cancer Prevention Research</i> , 2013, 6, 129-138.	0.7	9
452	TOX expression in cutaneous B-cell lymphomas. <i>Archives of Dermatological Research</i> , 2016, 308, 423-427.	1.1	9
453	Outcomes of rare patients with a primary cutaneous CD30+ lymphoproliferative disorder developing extracutaneous disease. <i>Blood</i> , 2020, 135, 769-773.	0.6	9
454	Whole-genome profiling of primary cutaneous anaplastic large cell lymphoma. <i>Haematologica</i> , 2022, 107, 1619-1632.	1.7	9
455	Murine hematopoietic progenitor cells with colony-forming or radioprotective capacity lack expression of the beta 2-integrin LFA-1. <i>Blood</i> , 1999, 93, 107-12.	0.6	9
456	Haemopoietic growth factor tyrosine kinase receptor expression profiles in normal haemopoiesis. <i>British Journal of Haematology</i> , 1996, 94, 236-241.	1.2	8
457	Commentary on "Cutaneous Manifestations of Lymphoma: A Clinical Guide Based on the WHO Classification". <i>Clinical Lymphoma and Myeloma</i> , 2001, 2, 101-102.	2.1	8
458	Cost analysis of CHOP (-like) chemotherapy regimens for patients with newly diagnosed aggressive non-Hodgkin's lymphoma. <i>European Journal of Haematology</i> , 2002, 69, 213-220.	1.1	8
459	Aleukaemic leukaemia cutis in a patient with acute T-cell lymphoblastic leukaemia. <i>Clinical and Experimental Dermatology</i> , 2003, 28, 330-332.	0.6	8
460	Umbilical cord blood "naive T cells but not adult blood "naive T cells require HLA class II on antigen-presenting cells for allo-immune activation. <i>Human Immunology</i> , 2004, 65, 328-339.	1.2	8
461	Pitfalls in diagnosing primary cutaneous aggressive epidermotropic CD8+ T-cell lymphoma. <i>British Journal of Dermatology</i> , 2019, 180, 411-412.	1.4	8
462	Rationale of a New Classification for the Group of Primary Cutaneous Lymphomas. <i>Seminars in Cutaneous Medicine and Surgery</i> , 2000, 19, 71-77.	1.6	8
463	Clonal hematopoiesis in patients with acquired aplastic anemia. <i>Blood</i> , 1991, 78, 3209-14.	0.6	8
464	Effect of the interval between high dose 1-beta-D-arabinofuranosylcytosine injections on leukemic cell load, intestinal toxicity, and normal hematopoietic stem cells in a rat model for acute myelogenous leukemia. <i>Cancer Research</i> , 1986, 46, 3825-7.	0.4	8
465	Reduction and repopulation of T-lymphocytes after cytoreductive therapy with or without autologous bone marrow rescue. <i>Experimental Hematology</i> , 1986, 14, 173-7.	0.2	8
466	Inhibitory effect of interleukin-4 on the proliferation of acute myeloid leukemia cells with myelo-monocytic differentiation (AML-M4/M5); the role of interleukin-6. <i>Leukemia</i> , 1993, 7, 643-5.	3.3	8
467	Thirty years of progress in cutaneous lymphoma research. <i>Giornale Italiano Di Dermatologia E Venereologia</i> , 2012, 147, 515-21.	0.8	8
468	Primary cutaneous lymphoma: the 2018 update of the WHO-EORTC classification. <i>Presse Medicale</i> , 2022, 51, 104126.	0.8	8

#	ARTICLE	IF	CITATIONS
469	B-cell colony growth of malignant and normal B-cells. <i>British Journal of Cancer</i> , 1987, 55, 397-405.	2.9	7
470	APLASTIC ANAEMIA PATIENTS WITH CLONAL Xâ€CHROMOSOME INACTIVATION PATTERN IN HAEMOPOIETIC CELLS EXHIBIT POLYCLONAL TCRÎ³ AND IgH GENE REARRANGEMENTS. <i>British Journal of Haematology</i> , 1996, 93, 326-332.	1.2	7
471	Leukemia Inhibitory Factor Induces In Vivo Expansion of Bone Marrow Progenitor Cells that Accelerate Hematopoietic Reconstitution but Do Not Enhance Radioprotection in Lethally Irradiated Mice. <i>Stem Cells</i> , 1997, 15, 50-55.	1.4	7
472	Unconventional rapid ERK1,2 activation is indispensable for proliferation of the growth factor-independent myeloid leukemic cell line KG1. <i>Leukemia</i> , 1998, 12, 699-704.	3.3	7
473	The Morphology of Dyserythropoiesis in a Patient with Acute Erythroleukaemia Associated with Multiple Myeloma. <i>Scandinavian Journal of Haematology</i> , 1976, 17, 353-368.	0.0	7
474	Point and period prevalences of eczema in rural and urban schoolchildren in Ghana, Gabon and Rwanda. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2012, 26, 488-494.	1.3	7
475	Increased Expression of PLS3 Correlates with Better Outcome in SÃ©zary Syndrome. <i>Journal of Investigative Dermatology</i> , 2017, 137, 754-757.	0.3	7
476	The prognostic significance of chromosomal findings in patients with acute myeloid leukemia in a study comparing the efficacy of autologous and allogeneic bone marrow transplantation. <i>Bone Marrow Transplantation</i> , 1991, 8, 377-81.	1.3	7
477	Jessner's lymphocytic infiltration of the skin. A clinical study of 100 patients. <i>Archives of Dermatology</i> , 1989, 125, 1525-30.	1.7	7
478	Homing and antigenic characterization of a human non-Hodgkin's lymphoma B cell line in severe combined immunodeficient (SCID) mice. <i>Leukemia</i> , 1994, 8, 1385-91.	3.3	7
479	Anti-CD45 and anti-CD52 (Campath) monoclonal antibodies effectively eliminate systematically disseminated human non-Hodgkin's lymphoma B cells in Scid mice. <i>Experimental Hematology</i> , 1996, 24, 919-26.	0.2	7
480	Correction of abnormal T-cell receptor repertoire during interferon-alpha therapy in patients with hairy cell leukemia. <i>Blood</i> , 1998, 91, 4224-31.	0.6	7
481	Day-6 bone marrow aspirate for the prediction of response to remission induction therapy for acute myelogenous leukaemia. <i>Blut</i> , 1988, 57, 91-95.	1.2	6
482	Cloning of the Dck gene encoding rat deoxycytidine kinase. <i>Gene</i> , 1994, 150, 351-354.	1.0	6
483	A novel RT-PCR-based protein activity truncation assay for direct assessment of deoxycytidine kinase in small numbers of purified leukemic cells. <i>Leukemia</i> , 2000, 14, 1678-1684.	3.3	6
484	Should primary cutaneous Ki-1(CD30)-positive anaplastic large cell lymphoma in childhood be treated with multiple-agent chemotherapy?. <i>Journal of the American Academy of Dermatology</i> , 2001, 45, 638-639.	0.6	6
485	Azathioprine-Induced Microsatellite Instability Is Not Observed in Skin Carcinomas of Organ Transplant Recipients. <i>Journal of Investigative Dermatology</i> , 2009, 129, 1307-1309.	0.3	6
486	XV. Primary cutaneous lymphomas. <i>Annals of Oncology</i> , 2011, 22, iv72-iv75.	0.6	6

#	ARTICLE	IF	CITATIONS
487	Tumor-stage mycosis fungoides of the vulva successfully treated with local low-dose radiotherapy. <i>Dermatologic Therapy</i> , 2015, 28, 36-39.	0.8	6
488	The clinical spectrum of mycosis fungoides in Tanzania, East Africa. <i>British Journal of Dermatology</i> , 2017, 176, 1653-1656.	1.4	6
489	Generalized Molluscum Contagiosum Successfully Treated with Interferon-Alpha in a Patient with Folliculotropic Mycosis Fungoides. <i>Case Reports in Dermatology</i> , 2019, 11, 52-56.	0.3	6
490	Ethnicity in mycosis fungoides: white patients present at an older age and with more advanced disease. <i>British Journal of Dermatology</i> , 2019, 180, 1264-1265.	1.4	6
491	Rationale of a new classification for the group of primary cutaneous lymphomas. <i>Seminars in Cutaneous Medicine and Surgery</i> , 2000, 19, 71-7.	1.6	6
492	Thrombocytopenic purpura, coeliac disease and IgA deficiency. <i>Netherlands Journal of Medicine</i> , 1986, 29, 165-6.	0.6	6
493	HOXC4, HOXC5, and HOXC6 expression in non-Hodgkin's lymphoma: preferential expression of the HOXC5 gene in primary cutaneous anaplastic T-cell and oro-gastrointestinal tract mucosa-associated B-cell lymphomas. <i>Blood</i> , 1997, 90, 4116-25.	0.6	6
494	Hairy cell leukemia-specific recognition by multiple autologous HLA-DQ or DP-restricted T-cell clones. <i>Blood</i> , 1999, 93, 251-9.	0.6	6
495	Solitary Plasmocytoma of the Vagina. <i>Acta Haematologica</i> , 1981, 66, 140-144.	0.7	5
496	Hairy cell leukemia: In-vitro proliferation and pseudo-colony formation. <i>Leukemia Research</i> , 1987, 11, 911-921.	0.4	5
497	Selective Response of CD5+B Cell Malignancies to Activation of the CD72 Antigen. <i>Clinical Immunology and Immunopathology</i> , 1998, 87, 42-49.	2.1	5
498	Comparable engraftment and chimerism kinetics using oral and intravenous fludarabine as part of a reduced intensity conditioning regimen. <i>Bone Marrow Transplantation</i> , 2008, 42, 137-138.	1.3	5
499	Quantitation of tumour development correlates with prognosis in tumour stage (stage IIB) mycosis fungoides. <i>British Journal of Dermatology</i> , 2014, 170, 1080-1086.	1.4	5
500	Absence of T(H)2 cytokine messenger RNA expression in CD30-negative primary cutaneous large T-cell lymphomas. <i>Archives of Dermatology</i> , 2001, 137, 901-5.	1.7	5
501	Engraftment potential into NOD/SCID mice of CD34+ cells derived from human fetal liver as compared to fetal bone marrow. <i>Haematologica</i> , 2001, 86, 1021-8.	1.7	5
502	Primary lymphoblastic non-Hodgkin's lymphoma of the skin. <i>British Journal of Dermatology</i> , 1981, 104, 333-338.	1.4	4
503	PREDICTION OF RESPONSE OF ACUTE NON-LYMPHOCYTIC LEUKAEMIA TO THERAPY WITH "HIGH DOSE" CYTOSINE ARABINOSIDE. <i>British Journal of Haematology</i> , 1985, 59, 733-734.	1.2	4
504	Human B cell colony assays. <i>Blut</i> , 1987, 55, 189-198.	1.2	4

#	ARTICLE	IF	CITATIONS
505	No Acceleration of UV-Induced Skin Carcinogenesis from Evenly Spread Dietary Intake of Cyclosporine in Contrast to Oral Bolus Dosages. <i>Transplantation</i> , 2013, 96, 871-876.	0.5	4
506	Corresponding anaplastic lymphoma kinase-tropomyosin 3 (ALK-TPM3) fusion in a patient with a primary cutaneous anaplastic large-cell lymphoma and a Spitz nevus. <i>JAAD Case Reports</i> , 2019, 5, 970-972.	0.4	4
507	Interleukin 4 down-regulates the expression of CD14 and the production of interleukin 6 in acute myeloid leukemia cells. <i>Lymphokine and Cytokine Research</i> , 1991, 10, 457-61.	0.7	4
508	Colony growth of normal and neoplastic cells in various concentrations of methylcellulose. <i>Experimental Hematology</i> , 1988, 16, 922-8.	0.2	4
509	The in vivo diffusion chamber technique for bone marrow or blood cell culture. <i>Blut</i> , 1981, 42, 59-67.	1.2	3
510	E-rosette positive acute lymphoblastic leukaemia in adolescents and adults. <i>British Journal of Haematology</i> , 1983, 55, 295-304.	1.2	3
511	A comparison of two culture techniques: An in vitro & an in vivo tumour colony-forming assay. <i>British Journal of Cancer</i> , 1985, 52, 713-717.	2.9	3
512	Detection of partial cDNA sequences differentially expressed in patients with myelodysplasia. <i>Annals of Hematology</i> , 1996, 72, 231-236.	0.8	3
513	Extramedullary hematopoietic growth factor and cytokine gene expression indicate endocrine regulation of hematopoiesis in patients with acute appendicitis. <i>Annals of Hematology</i> , 1997, 75, 201-205.	0.8	3
514	The risk of cancer is not increased in patients with multiple kidney transplantations. <i>Transplant Immunology</i> , 2012, 27, 189-194.	0.6	3
515	A rare case of cutaneous Epstein-Barr virus-negative intravascular cytotoxic T-cell lymphoma. <i>JAAD Case Reports</i> , 2019, 5, 548-551.	0.4	3
516	High Dose Intensive Chemotherapy, as Is Standard in Childhood Leukemia, Is Feasible and Efficacious in Adult Patients with Acute Lymphoblastic Leukemia (ALL) up to the Age of 40: Results From the Dutch-Belgian HOVON-70 Study. <i>Blood</i> , 2009, 114, 323-323.	0.6	3
517	Primary Cutaneous Follicle Center Lymphomas (PCFCL) Express Heavily Mutated B-Cell Receptors with Acquired N-Glycosylation Motifs and Lack Ongoing Somatic Hypermutation. <i>Blood</i> , 2018, 132, 1573-1573.	0.6	3
518	Experimental myelocytic leukemia in the Brown-Norway rat as a model for pulmonary leukostasis. <i>American Journal of Pathology</i> , 1991, 138, 777-80.	1.9	3
519	Total skin electron beam therapy for cutaneous T-cell lymphomas in the Netherlands: A retrospective analysis of treatment outcomes and selection for high or low dose schedule. <i>Clinical and Translational Radiation Oncology</i> , 2022, 33, 77-82.	0.9	3
520	Frequency and prognostic significance of clonal T-cell receptor beta-gene rearrangements in the peripheral blood of patients with mycosis fungoides. <i>Archives of Dermatology</i> , 1992, 128, 1602-7.	1.7	3
521	Serum monocyte colony-stimulating factor, erythropoietin and interleukin-6 in relation to pancytopenia in hairy cell leukemia. <i>Leukemia</i> , 1992, 6, 735-7.	3.3	3
522	Lymphomatoid papulosis. <i>Dermatologic Clinics</i> , 1985, 3, 735-47.	1.0	3

#	ARTICLE	IF	CITATIONS
523	Results of a multicentre study of the treatment of acute lymphoblastic leukaemia in adolescents and adults (1978-1981). <i>Netherlands Journal of Medicine</i> , 1982, 25, 303-7.	0.6	3
524	Constitutive in vivo cytokine and hematopoietic growth factor gene expression in the bone marrow and peripheral blood of healthy individuals. <i>Blood</i> , 1995, 85, 2038-44.	0.6	3
525	Direct display of hematopoietic tyrosine kinase receptor expression profiles in KG1 cells by PCR using degenerate primers. <i>Leukemia</i> , 1996, 10, 1395-9.	3.3	3
526	Unrelated bone marrow transplantation as a rescue procedure following inadvertent loss of an autologous bone marrow graft. <i>Bone Marrow Transplantation</i> , 1996, 18, 461-3.	1.3	3
527	Characterization of a novel malignant B cell line with t(14;18) and t(4;11) established from a patient with acute monoblastic leukemia. <i>Experimental Hematology</i> , 1997, 25, 1062-8.	0.2	3
528	Preliminary results of consolidation therapy with high-dose cytosine arabinoside for patients with bad-risk or relapsed acute leukemia or lymphoblastic non-Hodgkin's lymphoma. <i>European Journal of Cancer & Clinical Oncology</i> , 1987, 23, 401-405.	0.9	2
529	Recent developments in the early diagnosis of mycosis fungoides and SÅ©zary's syndrome. <i>European Journal of Cancer & Clinical Oncology</i> , 1987, 23, 1581-1583.	0.9	2
530	Deoxycoformycin in Therapy of Refractory Lymphoid Neoplasms. <i>Oncology Research and Treatment</i> , 1988, 11, 35-37.	0.8	2
531	Sublocalization of the breakpoints of a t(5;16) in myelodysplasia. <i>Cancer Genetics and Cytogenetics</i> , 1998, 100, 5-9.	1.0	2
532	Tumor necrosis factor alpha (Tnf-Î±) Production by acute myeloid leukemic (Aml) Blasts results in impaired proliferation of normal hematopoietic progenitor cells (Hpc). <i>Experimental Hematology</i> , 2000, 28, 69-70.	0.2	2
533	Clear cell acanthoma on the dorsum of the hand. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2016, 30, 378-380.	1.3	2
534	p16 immunostaining in keratinocytic neoplasia in organ transplant recipients: Bowen's disease shows a characteristic pattern. <i>Journal of Cutaneous Pathology</i> , 2017, 44, 28-33.	0.7	2
535	Serum and cutaneous transcriptional expression levels of IL31 are minimal in cutaneous T cell lymphoma variants. <i>Biochemistry and Biophysics Reports</i> , 2021, 26, 101007.	0.7	2
536	High-dose Ara-C for remission induction and consolidation of previously untreated adults with ALL or lymphoblastic lymphoma. <i>Annals of Hematology</i> , 1995, 70, 71-74.	0.8	2
537	Up-Front Window Trial of Gemtuzumab Ozogamicin (GO) in Previously Untreated Elderly Patients with AML: An EORTC Leukemia Group Study.. <i>Blood</i> , 2004, 104, 877-877.	0.6	2
538	Allogeneic Stem Cell Transplantation (Allo-SCT) in the Management of Advanced Waldenströmâ€™s Macroglobulinemia (WM).. <i>Blood</i> , 2007, 110, 619-619.	0.6	2
539	Gall bladder perforation in response to chemotherapy in a patient with a histiocytic lymphoma. <i>Netherlands Journal of Medicine</i> , 1988, 33, 78-82.	0.6	2
540	Extranodal non-Hodgkin's lymphoma. <i>Netherlands Journal of Medicine</i> , 1988, 33, 49-51.	0.6	2

#	ARTICLE	IF	CITATIONS
541	Monoclonal antibodies in the histopathological diagnosis of non-Hodgkin lymphomas. <i>Netherlands Journal of Medicine</i> , 1985, 28, 138-41.	0.6	2
542	Classification of primary cutaneous large cell lymphomas. <i>Dermatologic Clinics</i> , 1994, 12, 361-73.	1.0	2
543	Mycosis fungoides with extracutaneous localization in the breast. <i>British Journal of Dermatology</i> , 1996, 134, 1125-30.	1.4	2
544	An in vivo clonogenic assay for human tumors using plasma clot diffusion chambers implanted in mice. <i>European Journal of Cancer & Clinical Oncology</i> , 1985, 21, 127-134.	0.9	1
545	Radioresistant pseudo-colony formation in the PHA-leukocyte feeder colony assay. <i>Leukemia Research</i> , 1988, 12, 917-922.	0.4	1
546	(C11) Clonality of hematopoiesis in patients with aplastic anemia and myelodysplastic syndromes. <i>Leukemia Research</i> , 1991, 15, 7.	0.4	1
547	GM-CSF and IL-3 in Myelodysplastic Syndromes at Low Risk of Developing Acute Leukemia (MDS-LR). <i>Leukemia and Lymphoma</i> , 1992, 7, 64-64.	0.6	1
548	71 Clonality of hematopoiesis in myelodysplasia. <i>Leukemia Research</i> , 1997, 21, S19.	0.4	1
549	Classification and recommended treatment options for folliculotropic mycosis fungoides. <i>Expert Opinion on Orphan Drugs</i> , 2018, 6, 35-45.	0.5	1
550	Allogeneic (Allo-) or Autologous (Auto-) Peripheral Blood Stem Cell Transplantation (SCT) Randomized Versus a Second Intensive Consolidation after a Common Induction and Consolidation Course in Patients with Bad Prognosis Myelodysplastic Syndromes and Acute Myelogenous Leukemia Following MDS of More Than 6 Months Duration: The Final Analysis of a Joint Study (CRIANT) of the EORTC, EBMT, SAKK, HOVON and GIMEMA Leukemia Groups. <i>Blood</i> , 2005, 106, 794-794.	0.6	1
551	Incidence and Clinical Impact of TET2 Mutations in Acute Myeloid Leukemia Patients Treated within the EORTC/GIMEMA AML-12/06991 AML Trial. <i>Blood</i> , 2009, 114, 2609-2609.	0.6	1
552	Antigen-Independent, Autonomous B-Cell Receptor Signaling As a Dominant Candidate Oncogenic Mechanism in ABC DLBCL. <i>Blood</i> , 2016, 128, 778-778.	0.6	1
553	Treatment of acute myelogenous leukaemia in adults. A retrospective study of 70 patients (1968-1973). <i>Netherlands Journal of Medicine</i> , 1975, 18, 17-28.	0.6	1
554	Systemic polychemotherapy in patients with mycosis fungoides and lymph node involvement: a follow-up study of 17 patients. <i>Acta Dermato-Venereologica</i> , 1985, 65, 179-83.	0.6	1
555	Haematopoietic growth factors. <i>Netherlands Journal of Medicine</i> , 1989, 35, 151-68.	0.6	1
556	New techniques in the evaluation of cutaneous T-cell lymphoma. <i>Dermatologic Clinics</i> , 1985, 3, 665-72.	1.0	1
557	Bone marrow disorders in Hodgkin's disease: cyclic pancytopenia coinciding with Pel-Ebstein fever. <i>Netherlands Journal of Medicine</i> , 1983, 26, 74-6.	0.6	1
558	The diagnostic value of morphometry on blood lymphocytes in erythrodermic actinic reticuloid. <i>Archives of Dermatology</i> , 1995, 131, 1298-303.	1.7	1

#	ARTICLE	IF	CITATIONS
559	Mass Cytometric Analysis of Early-Stage Mycosis Fungoides. <i>Cells</i> , 2022, 11, 1062.	1.8	1
560	Marrow culture in diffusion chambers in rabbits. II. Effect of competing demands for red cell and white cell production on cell growth. <i>Blood</i> , 1978, 51, 1129-1135.	0.6	1
561	Willem A. van Vloten zum 60. Geburtstag. <i>Hautarzt</i> , 2001, 52, 547-548.	1.2	0
562	Cutaneous lymphomas. <i>Lancet Oncology</i> , The, 2005, 6, 839.	5.1	0
563	Limited effect of chemotherapy in cutaneous lymphoma. <i>Blood</i> , 2015, 125, 4-5.	0.6	0
564	176 The Prospective Cutaneous Lymphoma International Prognostic Index (PROCLIFI) Study identifies clinical prognostic markers and establishes the foundation for large-scale translational research in cutaneous lymphoma. <i>Journal of Investigative Dermatology</i> , 2019, 139, S31.	0.3	0
565	Primary Allogeneic T-Cell Responses Against In Vitro Modified Mantle Cell Lymphoma (MCL) Cells: Implications for Adoptive Immunotherapy after HLA-Matched Stem Cell Transplantation (SCT).. <i>Blood</i> , 2004, 104, 2118-2118.	0.6	0
566	Retroviral Interference with Execution Pathways in Target Cells To Unravel Mechanisms of Cytotoxic T Cell-Mediated Cell Death.. <i>Blood</i> , 2004, 104, 1267-1267.	0.6	0
567	First Line Treatment of Aggressive Cutaneous NK/T Cell Lymphomas Based on High Dose Cytarabine and Stem Cell Transplantation Does Not Lead to Durable Remissions.. <i>Blood</i> , 2006, 108, 4669-4669.	0.6	0
568	The B-Cell Receptor of Primary Cutaneous Follicle Center Lymphoma: Implications for Pathogenesis. <i>Blood</i> , 2016, 128, 4136-4136.	0.6	0
569	Post remission treatment in acute myelogenous leukaemia in first remission. The EORTC experience. EORTC Leukemia Cooperative Group. <i>Bone Marrow Transplantation</i> , 1989, 4 Suppl 3, 72.	1.3	0
570	Comparison of intensive consolidation therapy and autologous bone marrow transplantation in acute myeloid leukaemia. The Leiden experience. <i>Bone Marrow Transplantation</i> , 1989, 4 Suppl 3, 73.	1.3	0
571	The unrelated bone marrow donor. <i>Bone Marrow Transplantation</i> , 1988, 3, 371-7.	1.3	0
572	Effect of calcitriol on growth, differentiation, chemokine mRNA expression of cultured keratinocytes and on keratinocyte-T cell binding. <i>Acta Dermato-venereologica Supplementum</i> , 1994, 186, 52-4.	0.0	0
573	Tumor Clone Frequency Calculation Using High-Throughput Sequencing of the TCR β Gene in Patients with Folliculotropic Mycosis Fungoides. <i>Journal of Investigative Dermatology</i> , 2022, 142, 2544-2546.e2.	0.3	0