

Rein Willemze

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4579468/rein-willemze-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

555
papers

32,134
citations

92
h-index

156
g-index

583
ext. papers

35,408
ext. citations

4.9
avg, IF

6.56
L-index

#	Paper	IF	Citations
555	Cell-of-origin classification using the Hans and Lymph2Cx algorithms in primary cutaneous large B-cell lymphomas.. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2022 , 480, 667	5.1	3
554	Total skin electron beam therapy for cutaneous T-cell lymphomas in the Netherlands: A retrospective analysis of treatment outcomes and selection for high or low dose schedule.. <i>Clinical and Translational Radiation Oncology</i> , 2022 , 33, 77-82	4.6	
553	Primary cutaneous lymphoma: the 2018 update of the WHO-EORTC classification.. <i>Presse Medicale</i> , 2022 , 104126	2.2	0
552	Primary cutaneous peripheral T-cell lymphoma, not otherwise specified: results of a multicentre European Organization for Research and Treatment of Cancer (EORTC) cutaneous lymphoma taskforce study on the clinico-pathological and prognostic features. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2021 , 35, 658-668	4.6	4
551	Deregulation of JAK2 signaling underlies primary cutaneous CD8+ aggressive epidermotropic cytotoxic T-cell lymphoma. <i>Haematologica</i> , 2021 ,	6.6	5
550	Incidence of mycosis fungoides and Sary syndrome in the Netherlands between 2000 and 2020. <i>British Journal of Dermatology</i> , 2021 , 185, 434-435	4	5
549	Serum and cutaneous transcriptional expression levels of IL31 are minimal in cutaneous T cell lymphoma variants. <i>Biochemistry and Biophysics Reports</i> , 2021 , 26, 101007	2.2	1
548	Treatment of early-stage mycosis fungoides: results from the PROspective Cutaneous Lymphoma International Prognostic Index (PROCLIP) study. <i>British Journal of Dermatology</i> , 2021 , 184, 722-730	4	14
547	Should we be imaging lymph nodes at initial diagnosis of early-stage mycosis fungoides? Results from the PROspective Cutaneous Lymphoma International Prognostic Index (PROCLIP) international study. <i>British Journal of Dermatology</i> , 2021 , 184, 524-531	4	4
546	Quality of life in patients with Mycosis Fungoides and Sary Syndrome: a systematic review of the literature. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2021 , 35, 2377-2387	4.6	1
545	Outcomes of rare patients with a primary cutaneous CD30+ lymphoproliferative disorder developing extracutaneous disease. <i>Blood</i> , 2020 , 135, 769-773	2.2	4
544	Clinical, Histologic, and Molecular Characteristics of Anaplastic Lymphoma Kinase-positive Primary Cutaneous Anaplastic Large Cell Lymphoma. <i>American Journal of Surgical Pathology</i> , 2020 , 44, 776-781	6.7	11
543	Frequency and prognosis of associated malignancies in 504 patients with lymphomatoid papulosis. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2020 , 34, 260-266	4.6	14
542	Whole-genome analysis uncovers recurrent IKZF1 inactivation and aberrant cell adhesion in blastic plasmacytoid dendritic cell neoplasm. <i>Genes Chromosomes and Cancer</i> , 2020 , 59, 295-308	5	7
541	Plaque stage folliculotropic mycosis fungoides: histopathologic features and prognostic factors in a series of 40 patients. <i>Journal of Cutaneous Pathology</i> , 2020 , 47, 241-250	1.7	6
540	Generalized Molluscum Contagiosum Successfully Treated with Interferon-Alpha in a Patient with Folliculotropic Mycosis Fungoides. <i>Case Reports in Dermatology</i> , 2019 , 11, 52-56	1.1	2
539	Acquired N-Linked Glycosylation Motifs in B-Cell Receptors of Primary Cutaneous B-Cell Lymphoma and the Normal B-Cell Repertoire. <i>Journal of Investigative Dermatology</i> , 2019 , 139, 2195-2203	4.3	5

538	A rare case of cutaneous Epstein-Barr virus-negative intravascular cytotoxic T-cell lymphoma. <i>JAAD Case Reports</i> , 2019 , 5, 548-551	1.4	2
537	Corresponding anaplastic lymphoma kinase-tropomyosin 3 () fusion in a patient with a primary cutaneous anaplastic large-cell lymphoma and a Spitz nevus. <i>JAAD Case Reports</i> , 2019 , 5, 970-972	1.4	2
536	Ethnicity in mycosis fungoides: white patients present at an older age and with more advanced disease. <i>British Journal of Dermatology</i> , 2019 , 180, 1264-1265	4	4
535	The 2018 update of the WHO-EORTC classification for primary cutaneous lymphomas. <i>Blood</i> , 2019 , 133, 1703-1714	2.2	431
534	Pitfalls in diagnosing primary cutaneous aggressive epidermotropic CD8 T-cell lymphoma. <i>British Journal of Dermatology</i> , 2019 , 180, 411-412	4	5
533	The PROCLIFI international registry of early-stage mycosis fungoides identifies substantial diagnostic delay in most patients. <i>British Journal of Dermatology</i> , 2019 , 181, 350-357	4	66
532	Evaluation of treatment results in multifocal primary cutaneous anaplastic large cell lymphoma: report of the Dutch Cutaneous Lymphoma Group. <i>British Journal of Dermatology</i> , 2018 , 179, 724-731	4	14
531	High prevalence of and mutations in intravascular large B-cell lymphoma. <i>Blood</i> , 2018 , 131, 2086-2089	2.2	36
530	Classification and recommended treatment options for folliculotropic mycosis fungoides. <i>Expert Opinion on Orphan Drugs</i> , 2018 , 6, 35-45	1.1	
529	The Use of Central Pathology Review With Digital Slide Scanning in Advanced-stage Mycosis Fungoides and Szary Syndrome: A Multi-institutional and International Pathology Study. <i>American Journal of Surgical Pathology</i> , 2018 , 42, 726-734	6.7	14
528	Genomic analysis reveals recurrent deletion of JAK-STAT signaling inhibitors HNRNPK and SOCS1 in mycosis fungoides. <i>Genes Chromosomes and Cancer</i> , 2018 , 57, 653-664	5	35
527	Mycosis fungoides variants-clinicopathologic features, differential diagnosis, and treatment. <i>Seminars in Cutaneous Medicine and Surgery</i> , 2018 , 37, 11-17	1.4	18
526	Primary Cutaneous Follicle Center Lymphomas (PCFCL) Express Heavily Mutated B-Cell Receptors with Acquired N-Glycosylation Motifs and Lack Ongoing Somatic Hypermutation. <i>Blood</i> , 2018 , 132, 1573-1573	2.2	
525	Folliculotropic mycosis fungoides presenting with a solitary lesion: Clinicopathological features and long-term follow-up data in a series of 9 cases. <i>Journal of Cutaneous Pathology</i> , 2018 , 45, 122-128	1.7	9
524	High Incidence and Clinical Significance of MYC Rearrangements in Primary Cutaneous Diffuse Large B-Cell Lymphoma, Leg Type. <i>American Journal of Surgical Pathology</i> , 2018 , 42, 1488-1494	6.7	29
523	Primary cutaneous lymphomas: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. <i>Annals of Oncology</i> , 2018 , 29, iv30-iv40	10.3	99
522	Array-based CGH of primary cutaneous CD8+ aggressive EPIDERMOTROPIC cytotoxic T-cell lymphoma. <i>Genes Chromosomes and Cancer</i> , 2018 , 57, 622-629	5	6
521	Comparative value of post-remission treatment in cytogenetically normal AML subclassified by NPM1 and FLT3-ITD allelic ratio. <i>Leukemia</i> , 2017 , 31, 26-33	10.7	48

520	Increased Expression of PLS3 Correlates with Better Outcome in S \ddot{z} ary Syndrome. <i>Journal of Investigative Dermatology</i> , 2017 , 137, 754-757	4.3	5
519	Recommendations for treatment in folliculotropic mycosis fungoides: report of the Dutch Cutaneous Lymphoma Group. <i>British Journal of Dermatology</i> , 2017 , 177, 223-228	4	28
518	European Organisation for Research and Treatment of Cancer consensus recommendations for the treatment of mycosis fungoides/S \ddot{z} ary syndrome - Update 2017. <i>European Journal of Cancer</i> , 2017 , 77, 57-74	7.5	240
517	Cutaneous lymphomas with a panniculitic presentation. <i>Seminars in Diagnostic Pathology</i> , 2017 , 34, 36-43	4.3	21
516	Global patterns of care in advanced stage mycosis fungoides/Sezary syndrome: a multicenter retrospective follow-up study from the Cutaneous Lymphoma International Consortium. <i>Annals of Oncology</i> , 2017 , 28, 2517-2525	10.3	58
515	Recommendations for the Optimal Radiation Dose in Patients With Primary Cutaneous Anaplastic Large Cell Lymphoma: A Report of the Dutch Cutaneous Lymphoma Group. <i>International Journal of Radiation Oncology Biology Physics</i> , 2017 , 99, 1279-1285	4	12
514	p16 immunostaining in keratinocytic neoplasia in organ transplant recipients: Bowen's disease shows a characteristic pattern. <i>Journal of Cutaneous Pathology</i> , 2017 , 44, 28-33	1.7	2
513	The clinical spectrum of mycosis fungoides in Tanzania, East Africa. <i>British Journal of Dermatology</i> , 2017 , 176, 1653-1656	4	3
512	No TP63 rearrangements in a selected group of primary cutaneous CD30+ lymphoproliferative disorders with aggressive clinical course. <i>Blood</i> , 2016 , 128, 141-3	2.2	7
511	TOX expression in cutaneous T-cell lymphomas: an adjunctive diagnostic marker that is not tumour specific and not restricted to the CD4(+) 'CD8(-) phenotype. <i>British Journal of Dermatology</i> , 2016 , 175, 382-6	4	18
510	Clinical Staging and Prognostic Factors in Folliculotropic Mycosis Fungoides. <i>JAMA Dermatology</i> , 2016 , 152, 992-1000	5.1	85
509	TOX expression in cutaneous B-cell lymphomas. <i>Archives of Dermatological Research</i> , 2016 , 308, 423-7	3.3	6
508	Clear cell acanthoma on the dorsum of the hand. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2016 , 30, 378-80	4.6	1
507	The B-Cell Receptor of Primary Cutaneous Follicle Center Lymphoma: Implications for Pathogenesis. <i>Blood</i> , 2016 , 128, 4136-4136	2.2	
506	Antigen-Independent, Autonomous B-Cell Receptor Signaling As a Dominant Candidate Oncogenic Mechanism in ABC DLBCL. <i>Blood</i> , 2016 , 128, 778-778	2.2	
505	Evaluation of Immunophenotypic and Molecular Biomarkers for S \ddot{z} ary Syndrome Using Standard Operating Procedures: A Multicenter Study of 59 Patients. <i>Journal of Investigative Dermatology</i> , 2016 , 136, 1364-1372	4.3	59
504	Epigenomic Analysis of S \ddot{z} ary Syndrome Defines Patterns of Aberrant DNA Methylation and Identifies Diagnostic Markers. <i>Journal of Investigative Dermatology</i> , 2016 , 136, 1876-1884	4.3	31
503	Cutaneous Lymphoma International Consortium Study of Outcome in Advanced Stages of Mycosis Fungoides and S \ddot{z} ary Syndrome: Effect of Specific Prognostic Markers on Survival and Development of a Prognostic Model. <i>Journal of Clinical Oncology</i> , 2015 , 33, 3766-73	2.2	237

502	Limited effect of chemotherapy in cutaneous lymphoma. <i>Blood</i> , 2015 , 125, 4-5	2.2	
501	Recommendations for treatment of lymphomatoid papulosis with methotrexate: a report from the Dutch Cutaneous Lymphoma Group. <i>British Journal of Dermatology</i> , 2015 , 173, 1319-22	4	21
500	Histopathological and immunophenotypical criteria for the diagnosis of Szary syndrome in differentiation from other erythrodermic skin diseases: a European Organisation for Research and Treatment of Cancer (EORTC) Cutaneous Lymphoma Task Force Study of 97 cases. <i>British Journal of Dermatology</i> , 2015 , 173, 93-105	4	44
499	Differential expression of TOX by skin-infiltrating T cells in Szary syndrome and erythrodermic dermatitis. <i>Journal of Cutaneous Pathology</i> , 2015 , 42, 604-9	1.7	14
498	Aggressive epidermotropic cutaneous CD8+ lymphoma: a cutaneous lymphoma with distinct clinical and pathological features. Report of an EORTC Cutaneous Lymphoma Task Force Workshop. <i>Histopathology</i> , 2015 , 67, 425-41	7.3	61
497	Diagnostic and prognostic significance of CDKN2A/CDKN2B deletions in patients with transformed mycosis fungoides and primary cutaneous CD30-positive lymphoproliferative disease. <i>British Journal of Dermatology</i> , 2015 , 172, 784-8	4	14
496	Tumor-stage mycosis fungoides of the vulva successfully treated with local low-dose radiotherapy. <i>Dermatologic Therapy</i> , 2015 , 28, 36-9	2.2	6
495	High numbers of mobilized CD34+ cells collected in AML in first remission are associated with high relapse risk irrespective of treatment with autologous peripheral blood SCT or autologous BMT. <i>Bone Marrow Transplantation</i> , 2015 , 50, 341-7	4.4	11
494	Clofarabine in combination with a standard remission induction regimen (cytosine arabinoside and idarubicin) in patients with previously untreated intermediate and bad-risk acute myelogenous leukemia (AML) or high-risk myelodysplastic syndrome (HR-MDS): phase I results of an ongoing	3	8
493	phase I/II study of the leukemia groups of EORTC and GIMEMA (EORTC-GIMEMA 06051/AML-14A). Myeloablative T cell-depleted alloSCT with early sequential prophylactic donor lymphocyte infusion is an efficient and safe post-remission treatment for adult ALL. <i>Bone Marrow Transplantation</i> , 2014 , 49, 287-91	4.4	33
492	Nuclear factor- κ B pathway-activating gene aberrancies in primary cutaneous large B-cell lymphoma, leg type. <i>Journal of Investigative Dermatology</i> , 2014 , 134, 290-292	4.3	41
491	Expression of programmed death-1 in skin biopsies of benign inflammatory vs. lymphomatous erythroderma. <i>British Journal of Dermatology</i> , 2014 , 171, 499-504	4	20
490	Programmed death-1 expression in cutaneous B-cell lymphoma. <i>Journal of Cutaneous Pathology</i> , 2014 , 41, 14-21	1.7	17
489	Quantitation of tumour development correlates with prognosis in tumour stage (stage IIB) mycosis fungoides. <i>British Journal of Dermatology</i> , 2014 , 170, 1080-6	4	3
488	Recent advances in primary cutaneous B-cell lymphomas. <i>Current Opinion in Oncology</i> , 2014 , 26, 230-6	4.2	7
487	Methotrexate-associated B-cell lymphoproliferative disorders presenting in the skin: A clinicopathologic and immunophenotypical study of 10 cases. <i>American Journal of Surgical Pathology</i> , 2014 , 38, 999-1006	6.7	27
486	CD30-Negative Cutaneous T-Cell Lymphomas Other than Mycosis Fungoides. <i>Surgical Pathology Clinics</i> , 2014 , 7, 229-52	3.9	7
485	Mycosis fungoides: report of the 2011 Society for Hematopathology/European Association for Haematopathology workshop. <i>American Journal of Clinical Pathology</i> , 2013 , 139, 466-90	1.9	46

484	Non-mycosis fungoides cutaneous T-cell lymphomas: report of the 2011 Society for Hematopathology/European Association for Haematopathology workshop. <i>American Journal of Clinical Pathology</i> , 2013 , 139, 491-514	1.9	53
483	Effects of serially passaged fibroblasts on dermal and epidermal morphogenesis in human skin equivalents. <i>Biogerontology</i> , 2013 , 14, 131-40	4.5	21
482	Identification of prognostic factors predicting outcome in Hodgkin's lymphoma patients relapsing after autologous stem cell transplantation. <i>Annals of Oncology</i> , 2013 , 24, 2430-4	10.3	74
481	Dietary immunosuppressants do not enhance UV-induced skin carcinogenesis, and reveal discordance between p53-mutant early clones and carcinomas. <i>Cancer Prevention Research</i> , 2013 , 6, 129-33	3.3	9
480	Mimics of cutaneous lymphoma: report of the 2011 Society for Hematopathology/European Association for Haematopathology workshop. <i>American Journal of Clinical Pathology</i> , 2013 , 139, 536-51	1.9	42
479	Primary cutaneous lymphomas: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. <i>Annals of Oncology</i> , 2013 , 24 Suppl 6, vi149-54	10.3	98
478	Skin diseases among schoolchildren in Ghana, Gabon, and Rwanda. <i>International Journal of Dermatology</i> , 2013 , 52, 589-600	1.7	28
477	Cutaneous B-cell lymphoproliferative disorders: report of the 2011 Society for Hematopathology/European Association for Haematopathology workshop. <i>American Journal of Clinical Pathology</i> , 2013 , 139, 515-35	1.9	43
476	No acceleration of UV-induced skin carcinogenesis from evenly spread dietary intake of cyclosporine in contrast to oral bolus dosages. <i>Transplantation</i> , 2013 , 96, 871-6	1.8	4
475	MicroRNA profiling of primary cutaneous large B-cell lymphomas. <i>PLoS ONE</i> , 2013 , 8, e82471	3.7	15
474	Primary cutaneous anaplastic large cell lymphoma shows a distinct miRNA expression profile and reveals differences from tumor-stage mycosis fungoides. <i>Experimental Dermatology</i> , 2012 , 21, 632-4	4	39
473	A meta-analysis of gene expression data identifies a molecular signature characteristic for tumor-stage mycosis fungoides. <i>Journal of Investigative Dermatology</i> , 2012 , 132, 2050-9	4.3	64
472	Differential expression of programmed death-1 (PD-1) in Szary syndrome and mycosis fungoides. <i>Archives of Dermatology</i> , 2012 , 148, 1379-85		80
471	Langerhans cell histiocytosis first presenting in the skin in adults: frequent association with a second haematological malignancy. <i>British Journal of Dermatology</i> , 2012 , 167, 1287-94	4	44
470	The risk of cancer is not increased in patients with multiple kidney transplantations. <i>Transplant Immunology</i> , 2012 , 27, 189-94	1.7	1
469	Prognostic factors in transformed mycosis fungoides: a retrospective analysis of 100 cases. <i>Blood</i> , 2012 , 119, 1643-9	2.2	136
468	Generation and administration of HA-1-specific T-cell lines for the treatment of patients with relapsed leukemia after allogeneic stem cell transplantation: a pilot study. <i>Haematologica</i> , 2012 , 97, 1205-8	6.6	45
467	Rapamycin impairs UV induction of mutant-p53 overexpressing cell clusters without affecting tumor onset. <i>International Journal of Cancer</i> , 2012 , 131, 1267-76	7.5	11

466	Kidney transplant recipients with cutaneous squamous cell carcinoma have an increased risk of internal malignancy. <i>Journal of Investigative Dermatology</i> , 2012 , 132, 2176-83	4.3	14
465	Point and period prevalences of eczema in rural and urban schoolchildren in Ghana, Gabon and Rwanda. <i>Journal of the European Academy of Dermatology and Venereology</i> , 2012 , 26, 488-94	4.6	7
464	Autocrine IL-21 stimulation is involved in the maintenance of constitutive STAT3 activation in Sjögren syndrome. <i>Journal of Investigative Dermatology</i> , 2012 , 132, 440-7	4.3	32
463	Deep-sequencing analysis reveals that the miR-199a2/214 cluster within DNMT3os represents the vast majority of aberrantly expressed microRNAs in Sjögren syndrome. <i>Journal of Investigative Dermatology</i> , 2012 , 132, 1520-2	4.3	38
462	Human allo-reactive CD4+ T cells as strong mediators of anti-tumor immunity in NOD/scid mice engrafted with human acute lymphoblastic leukemia. <i>Leukemia</i> , 2012 , 26, 312-22	10.7	24
461	Intensive chemotherapy to improve outcome in patients with acute lymphoblastic leukemia over the age of 40: a phase II study for efficacy and feasibility by HOVON. <i>Leukemia</i> , 2012 , 26, 1726-9	10.7	18
460	Expression of programmed death-1 in primary cutaneous CD4-positive small/medium-sized pleomorphic T-cell lymphoma, cutaneous pseudo-T-cell lymphoma, and other types of cutaneous T-cell lymphoma. <i>American Journal of Surgical Pathology</i> , 2012 , 36, 109-16	6.7	123
459	Allogeneic stem cell transplantation in paroxysmal nocturnal hemoglobinuria. <i>Haematologica</i> , 2012 , 97, 1666-73	6.6	81
458	Thirty years of progress in cutaneous lymphoma research. <i>Giornale Italiano Di Dermatologia E Venereologia</i> , 2012 , 147, 515-21	0.8	5
457	miRNA expression profiling of mycosis fungoides. <i>Molecular Oncology</i> , 2011 , 5, 273-80	7.9	79
456	Prevalence and causative fungal species of tinea capitis among schoolchildren in Gabon. <i>Mycoses</i> , 2011 , 54, e354-9	5.2	22
455	Functional characterization of cancer-associated fibroblasts of human cutaneous squamous cell carcinoma. <i>Experimental Dermatology</i> , 2011 , 20, 737-42	4	25
454	Predictive impact of allele-matching and EBMT risk score for outcome after T-cell depleted unrelated donor transplantation in poor-risk acute leukemia and myelodysplasia. <i>Leukemia</i> , 2011 , 25, 1548-54	10.7	16
453	Incidence of cancer in kidney-transplant recipients: a long-term cohort study in a single center. <i>Cancer Epidemiology</i> , 2011 , 35, 105-11	2.8	57
452	MicroRNA-21 expression in CD4+ T cells is regulated by STAT3 and is pathologically involved in Sjögren syndrome. <i>Journal of Investigative Dermatology</i> , 2011 , 131, 762-8	4.3	104
451	Intensified chemotherapy inspired by a pediatric regimen combined with allogeneic transplantation in adult patients with acute lymphoblastic leukemia up to the age of 40. <i>Leukemia</i> , 2011 , 25, 1697-703	10.7	61
450	XV. Primary cutaneous lymphomas. <i>Annals of Oncology</i> , 2011 , 22, iv72-iv75	10.3	6
449	Trends of skin diseases in organ-transplant recipients transplanted between 1966 and 2006: a cohort study with follow-up between 1994 and 2006. <i>British Journal of Dermatology</i> , 2010 , 162, 390-6	4	21

448	Primary cutaneous lymphomas: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. <i>Annals of Oncology</i> , 2010 , 21 Suppl 5, v177-80	10.3	31
447	Mixed T cell receptor dimers harbor potentially harmful neoreactivity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 10972-7	11.5	165
446	Cutaneous anaplastic large cell lymphoma and peripheral T-cell lymphoma NOS show distinct chromosomal alterations and differential expression of chemokine receptors and apoptosis regulators. <i>Journal of Investigative Dermatology</i> , 2010 , 130, 563-75	4.3	54
445	Efficacy of a hypofractionated schedule in electron beam radiotherapy for epithelial skin cancer: Analysis of 434 cases. <i>Radiotherapy and Oncology</i> , 2010 , 95, 245-9	5.3	60
444	IgM expression on paraffin sections distinguishes primary cutaneous large B-cell lymphoma, leg type from primary cutaneous follicle center lymphoma. <i>American Journal of Surgical Pathology</i> , 2010 , 34, 1043-8	6.7	73
443	Subsequent squamous- and basal-cell carcinomas in kidney-transplant recipients after the first skin cancer: cumulative incidence and risk factors. <i>Transplantation</i> , 2010 , 89, 1231-8	1.8	77
442	Efficacy of escalated imatinib combined with cytarabine in newly diagnosed patients with chronic myeloid leukemia. <i>Haematologica</i> , 2010 , 95, 914-21	6.6	18
441	Combined CD8+ and CD4+ adenovirus hexon-specific T cells associated with viral clearance after stem cell transplantation as treatment for adenovirus infection. <i>Haematologica</i> , 2010 , 95, 1943-51	6.6	56
440	Dexamethasone compared to prednisolone for adults with acute lymphoblastic leukemia or lymphoblastic lymphoma: final results of the ALL-4 randomized, phase III trial of the EORTC Leukemia Group. <i>Haematologica</i> , 2010 , 95, 1489-95	6.6	11
439	Value of allogeneic versus autologous stem cell transplantation and chemotherapy in patients with myelodysplastic syndromes and secondary acute myeloid leukemia. Final results of a prospective randomized European Intergroup Trial. <i>Haematologica</i> , 2010 , 95, 1754-61	6.6	62
438	Cutaneous gamma/delta T-cell lymphoma during treatment with etanercept for rheumatoid arthritis. <i>Acta Dermato-Venereologica</i> , 2009 , 89, 653-4	2.2	22
437	Primary cutaneous lymphoma: ESMO clinical recommendations for diagnosis, treatment and follow-up. <i>Annals of Oncology</i> , 2009 , 20 Suppl 4, 115-8	10.3	18
436	Applicability and prognostic value of the new TNM classification system in 135 patients with primary cutaneous anaplastic large cell lymphoma. <i>Archives of Dermatology</i> , 2009 , 145, 1399-404		69
435	Diagnostic and prognostic evaluation of phenotypic markers TRAF1, MUM1, BCL2 and CD15 in cutaneous CD30-positive lymphoproliferative disorders. <i>British Journal of Dermatology</i> , 2009 , 161, 121-7 ⁴		52
434	Prevalence and risk factors of inflammatory acne vulgaris in rural and urban Ghanaian schoolchildren. <i>British Journal of Dermatology</i> , 2009 , 161, 475-7	4	20
433	Fine-mapping chromosomal loss at 9p21: correlation with prognosis in primary cutaneous diffuse large B-cell lymphoma, leg type. <i>Journal of Investigative Dermatology</i> , 2009 , 129, 1149-55	4.3	69
432	Azathioprine-induced microsatellite instability is not observed in skin carcinomas of organ transplant recipients. <i>Journal of Investigative Dermatology</i> , 2009 , 129, 1307-9	4.3	6
431	Increased risk of squamous-cell carcinoma in simultaneous pancreas kidney transplant recipients compared with kidney transplant recipients. <i>Journal of Investigative Dermatology</i> , 2009 , 129, 2886-94	4.3	28

430	KIR-ligand incompatibility in the graft-versus-host direction improves outcomes after umbilical cord blood transplantation for acute leukemia. <i>Leukemia</i> , 2009 , 23, 492-500	10.7	202
429	Reduced IL-1Ra/IL-1 ratio in ultraviolet B-exposed skin of patients with polymorphic light eruption. <i>Experimental Dermatology</i> , 2009 , 18, 212-7	4	18
428	An in vitro three-dimensional model of primary human cutaneous squamous cell carcinoma. <i>Experimental Dermatology</i> , 2009 , 18, 849-56	4	39
427	Replacement of animal-derived collagen matrix by human fibroblast-derived dermal matrix for human skin equivalent products. <i>Biomaterials</i> , 2009 , 30, 71-8	15.6	86
426	Low-dose palliative radiotherapy for cutaneous B- and T-cell lymphomas. <i>International Journal of Radiation Oncology Biology Physics</i> , 2009 , 74, 154-8	4	105
425	Retroviral transfer of human CD20 as a suicide gene for adoptive T-cell therapy. <i>Haematologica</i> , 2009 , 94, 1316-20	6.6	94
424	Oncogenomic analysis of mycosis fungoides reveals major differences with Sezary syndrome. <i>Blood</i> , 2009 , 113, 127-36	2.2	154
423	High Dose Intensive Chemotherapy, as Is Standard in Childhood Leukemia, Is Feasible and Efficacious in Adult Patients with Acute Lymphoblastic Leukemia (ALL) up to the Age of 40: Results From the Dutch-Belgian HOVON-70 Study.. <i>Blood</i> , 2009 , 114, 323-323	2.2	2
422	Incidence and Clinical Impact of TET2 Mutations in Acute Myeloid Leukemia Patients Treated within the EORTC/GIMEMA AML-12/06991 AML Trial.. <i>Blood</i> , 2009 , 114, 2609-2609	2.2	
421	Genome-wide linkage scan for atypical nevi in p16-Leiden melanoma families. <i>European Journal of Human Genetics</i> , 2008 , 16, 1135-41	5.3	17
420	HLA-DP as specific target for cellular immunotherapy in HLA class II-expressing B-cell leukemia. <i>Leukemia</i> , 2008 , 22, 1387-94	10.7	49
419	Comparable engraftment and chimerism kinetics using oral and intravenous fludarabine as part of a reduced intensity conditioning regimen. <i>Bone Marrow Transplantation</i> , 2008 , 42, 137-8	4.4	5
418	Cucurbitacin I inhibits Stat3 and induces apoptosis in Sezary cells. <i>Journal of Investigative Dermatology</i> , 2008 , 128, 1691-5	4.3	60
417	Bone marrow examination has limited value in the staging of patients with an anaplastic large cell lymphoma first presenting in the skin. Retrospective analysis of 107 patients. <i>British Journal of Dermatology</i> , 2008 , 159, 1148-51	4	13
416	Results of bone marrow examination in 275 patients with histological features that suggest an indolent type of cutaneous B-cell lymphoma. <i>British Journal of Haematology</i> , 2008 , 142, 52-6	4.5	53
415	The addition of rituximab to anthracycline-based chemotherapy significantly improves outcome in 'Western' patients with intravascular large B-cell lymphoma. <i>British Journal of Haematology</i> , 2008 , 143, 253-7	4.5	70
414	Leiden reconstructed human epidermal model as a tool for the evaluation of the skin corrosion and irritation potential according to the ECVAM guidelines. <i>Toxicology in Vitro</i> , 2008 , 22, 1311-20	3.6	57
413	Subcutaneous panniculitis-like T-cell lymphoma: definition, classification, and prognostic factors: an EORTC Cutaneous Lymphoma Group Study of 83 cases. <i>Blood</i> , 2008 , 111, 838-45	2.2	492

412	Granulomatous mycosis fungoides and granulomatous slack skin: a multicenter study of the Cutaneous Lymphoma Histopathology Task Force Group of the European Organization For Research and Treatment of Cancer (EORTC). <i>Archives of Dermatology</i> , 2008 , 144, 1609-17		121
411	Novel and highly recurrent chromosomal alterations in Sjögren syndrome. <i>Cancer Research</i> , 2008 , 68, 2689-98		148
410	Results of syngeneic hematopoietic stem cell transplantation for acute leukemia: risk factors for outcomes of adults transplanted in first complete remission. <i>Haematologica</i> , 2008 , 93, 834-41	6.6	9
409	Increased risk of cancer other than melanoma in CDKN2A founder mutation (p16-Leiden)-positive melanoma families. <i>Clinical Cancer Research</i> , 2008 , 14, 7151-7	12.9	135
408	Identification of phosphatidylinositol 4-kinase type II beta as HLA class II-restricted target in graft versus leukemia reactivity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008 , 105, 3837-42	11.5	47
407	An immunodeficiency disease with RAG mutations and granulomas. <i>New England Journal of Medicine</i> , 2008 , 358, 2030-8	59.2	184
406	The majority of cutaneous marginal zone B-cell lymphomas expresses class-switched immunoglobulins and develops in a T-helper type 2 inflammatory environment. <i>Blood</i> , 2008 , 112, 3355-61	2.2	73
405	Genetic engineering of virus-specific T cells with T-cell receptors recognizing minor histocompatibility antigens for clinical application. <i>Haematologica</i> , 2008 , 93, 1535-43	6.6	33
404	A genomic and expression study of AP-1 in primary cutaneous T-cell lymphoma: evidence for dysregulated expression of JUNB and JUND in MF and SS. <i>Journal of Cutaneous Pathology</i> , 2008 , 35, 899-910	1.7	42
403	Identification of selectively expressed genes and antigens in CTCL. <i>Experimental Dermatology</i> , 2008 , 17, 324-34	4	25
402	More epidermal p53 patches adjacent to skin carcinomas in renal transplant recipients than in immunocompetent patients: the role of azathioprine. <i>Experimental Dermatology</i> , 2008 , 17, 349-55	4	34
401	Differential activation of the death receptor pathway in human target cells induced by cytotoxic T lymphocytes showing different kinetics of killing. <i>Haematologica</i> , 2007 , 92, 1671-8	6.6	12
400	Characterization of graft-versus-leukemia responses in patients treated for advanced chronic lymphocytic leukemia with donor lymphocyte infusions after in vitro T-cell depleted allogeneic stem cell transplantation following reduced-intensity conditioning. <i>Leukemia</i> , 2007 , 21, 2569-74	10.7	15
399	The applicability and prognostic value of the new TNM classification system for primary cutaneous lymphomas other than mycosis fungoides and Sjögren syndrome: results on a large cohort of primary cutaneous B-cell lymphomas and comparison with the system used by the Dutch Cutaneous Lymphoma Group. <i>Journal of Cutaneous Medicine and Surgery</i> , 2007 , 13, 1005-11	4	35
398	Involvement of caspase-8 in chemotherapy-induced apoptosis of patient derived leukemia cell lines independent of the death receptor pathway and downstream from mitochondria. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2007 , 12, 181-93	5.4	16
397	Variations in clinical presentation, frequency of hemophagocytosis and clinical behavior of intravascular lymphoma diagnosed in different geographical regions. <i>Haematologica</i> , 2007 , 92, 486-92	6.6	137
396	Reclassification of 300 primary cutaneous B-Cell lymphomas according to the new WHO-EORTC classification for cutaneous lymphomas: comparison with previous classifications and identification of prognostic markers. <i>Journal of Clinical Oncology</i> , 2007 , 25, 1581-7	2.2	215
395	Results of radiotherapy in 153 primary cutaneous B-Cell lymphomas classified according to the WHO-EORTC classification. <i>Archives of Dermatology</i> , 2007 , 143, 1520-6		93

394	Phase I/II feasibility study evaluating the generation of leukemia-reactive cytotoxic T lymphocyte lines for treatment of patients with relapsed leukemia after allogeneic stem cell transplantation. <i>Haematologica</i> , 2007 , 92, 72-80	6.6	44
393	Gene-expression profiling and array-based CGH classify CD4+CD56+ hematodermic neoplasm and cutaneous myelomonocytic leukemia as distinct disease entities. <i>Blood</i> , 2007 , 109, 1720-7	2.2	115
392	Indolent CD8-positive lymphoid proliferation of the ear: a distinct primary cutaneous T-cell lymphoma?. <i>American Journal of Surgical Pathology</i> , 2007 , 31, 1887-92	6.7	144
391	From sporadic atypical nevi to familial melanoma: risk analysis for melanoma in sporadic atypical nevus patients. <i>Journal of the American Academy of Dermatology</i> , 2007 , 56, 748-52	4.5	22
390	Human mesenchymal stem cells derived from bone marrow display a better chondrogenic differentiation compared with other sources. <i>Connective Tissue Research</i> , 2007 , 48, 132-40	3.3	89
389	Allogeneic Stem Cell Transplantation (Allo-SCT) in the Management of Advanced Waldenstrom's Macroglobulinemia (WM).. <i>Blood</i> , 2007 , 110, 619-619	2.2	
388	Efficacy and toxicity of gemtuzumab ozogamicin in patients with acute myeloid leukemia. <i>European Journal of Haematology</i> , 2006 , 76, 409-13	3.8	22
387	Serpina1 is a potent inhibitor of IL-8-induced hematopoietic stem cell mobilization. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006 , 103, 1469-74	11.5	53
386	Array-based comparative genomic hybridization analysis reveals recurrent chromosomal alterations and prognostic parameters in primary cutaneous large B-cell lymphoma. <i>Journal of Clinical Oncology</i> , 2006 , 24, 296-305	2.2	101
385	Similar potential to become activated and proliferate but differential kinetics and profiles of cytokine production of umbilical cord blood T cells and adult blood naive and memory T cells. <i>Human Immunology</i> , 2006 , 67, 874-83	2.3	15
384	Primary cutaneous follicle center lymphoma and primary cutaneous large B-cell lymphoma, leg type, are both targeted by aberrant somatic hypermutation but demonstrate differential expression of AID. <i>Blood</i> , 2006 , 107, 4926-9	2.2	40
383	Primary cutaneous B-cell lymphoma: classification and treatment. <i>Current Opinion in Oncology</i> , 2006 , 18, 425-31	4.2	45
382	Classification of cutaneous T-cell lymphoma: from Alibert to WHO-EORTC. <i>Journal of Cutaneous Pathology</i> , 2006 , 33 Suppl 1, 18-26	1.7	100
381	The occurrence of residual or recurrent squamous cell carcinomas in organ transplant recipients after curettage and electrodesiccation. <i>British Journal of Dermatology</i> , 2006 , 154, 493-7	4	21
380	Expression of B-cell transcription factors in primary cutaneous B-cell lymphoma. <i>Modern Pathology</i> , 2006 , 19, 1270-6	9.8	33
379	Oral valganciclovir as pre-emptive therapy has similar efficacy on cytomegalovirus DNA load reduction as intravenous ganciclovir in allogeneic stem cell transplantation recipients. <i>Bone Marrow Transplantation</i> , 2006 , 37, 693-8	4.4	48
378	Long-term follow-up of myeloablative allogeneic stem cell transplantation using Campath "in the bag" as T-cell depletion: the Leiden experience. <i>Bone Marrow Transplantation</i> , 2006 , 37, 1129-34	4.4	52
377	Photodynamic therapy does not prevent cutaneous squamous-cell carcinoma in organ-transplant recipients: results of a randomized-controlled trial. <i>Journal of Investigative Dermatology</i> , 2006 , 126, 569-74	4.7	83

376	Molecular persistence of chronic myeloid leukemia caused by donor T cells specific for lineage-restricted maturation antigens not recognizing immature progenitor-cells. <i>Leukemia</i> , 2006 , 20, 1040-6	10.7	15
375	Non-infusional vs intravenous consolidation chemotherapy in elderly patients with acute myeloid leukemia: final results of the EORTC-GIMEMA AML-13 randomized phase III trial. <i>Leukemia</i> , 2006 , 20, 1723-30	10.7	19
374	First Line Treatment of Aggressive Cutaneous NK/T Cell Lymphomas Based on High Dose Cytarabine and Stem Cell Transplantation Does Not Lead to Durable Remissions.. <i>Blood</i> , 2006 , 108, 4669-4669	2.2	2980
373	WHO-EORTC classification for cutaneous lymphomas. <i>Blood</i> , 2005 , 105, 3768-85	2.2	2980
372	Primary cutaneous marginal zone B-cell lymphoma: clinical and therapeutic features in 50 cases. <i>Archives of Dermatology</i> , 2005 , 141, 1139-45		130
371	Distinct types of primary cutaneous large B-cell lymphoma identified by gene expression profiling. <i>Blood</i> , 2005 , 105, 3671-8	2.2	214
370	Chemical exposures other than arsenic are probably not important risk factors for squamous cell carcinoma, basal cell carcinoma and malignant melanoma of the skin. <i>British Journal of Dermatology</i> , 2005 , 152, 194-7	4	14
369	Normalized ultraviolet (UV) induction of Langerhans cell depletion and neutrophil infiltrates after artificial UVB hardening of patients with polymorphic light eruption. <i>British Journal of Dermatology</i> , 2005 , 152, 1268-74	4	45
368	Varicella zoster virus (VZV)-related progressive outer retinal necrosis (PORN) after allogeneic stem cell transplantation. <i>Bone Marrow Transplantation</i> , 2005 , 36, 467-9	4.4	19
367	Minor histocompatibility antigen-specific T cells with multiple distinct specificities can be isolated by direct cloning of IFNgamma-secreting T cells from patients with relapsed leukemia responding to donor lymphocyte infusion. <i>Leukemia</i> , 2005 , 19, 83-90	10.7	24
366	Gemtuzumab ozogamicin (Mylotarg) as single-agent treatment for frail patients 61 years of age and older with acute myeloid leukemia: final results of AML-15B, a phase 2 study of the European Organisation for Research and Treatment of Cancer and Gruppo Italiano Malattie Ematologiche dell'Adulto Leukemia Group. <i>Leukemia</i> , 2005 , 19, 1768-73	10.7	78
365	Low-dose cytosine arabinoside (LD-AraC) vs LD-AraC plus granulocyte/macrophage colony stimulating factor vs LD-AraC plus Interleukin-3 for myelodysplastic syndrome patients with a high risk of developing acute leukemia: final results of a randomized phase III study (06903) of the EORTC Leukemia Cooperative Group. <i>Leukemia</i> , 2005 , 19, 1822-33	10.7	29
364	WHO/EORTC classification of cutaneous lymphomas 2005: histological and molecular aspects. <i>Journal of Cutaneous Pathology</i> , 2005 , 32, 647-74	1.7	271
363	UVB-induced leucocyte trafficking in the epidermis of photosensitive lupus erythematosus patients: normal depletion of Langerhans cells. <i>Experimental Dermatology</i> , 2005 , 14, 138-42	4	15
362	Aberrant DNA methylation in cutaneous malignancies. <i>Seminars in Oncology</i> , 2005 , 32, 479-87	5.5	59
361	Epidermal transit of replication-arrested, undifferentiated keratinocytes in UV-exposed XPC mice: an alternative to in situ apoptosis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005 , 102, 18980-5	11.5	14
360	HLA class II restricted T-cell receptor gene transfer generates CD4+ T cells with helper activity as well as cytotoxic capacity. <i>Gene Therapy</i> , 2005 , 12, 1686-95	4	23
359	Epigenetic profiling of cutaneous T-cell lymphoma: promoter hypermethylation of multiple tumor suppressor genes including BCL7a, PTPRG, and p73. <i>Journal of Clinical Oncology</i> , 2005 , 23, 3886-96	2.2	197

358	Allogeneic (Allo-) or Autologous (Auto-) Peripheral Blood Stem Cell Transplantation (SCT) Randomized Versus a Second Intensive Consolidation after a Common Induction and Consolidation Course in Patients with Bad Prognosis Myelodysplastic Syndromes and Acute Myelogenous Leukemia Following MDS of More Than 6 Months Duration: The Final Analysis of a Joint Study	2.2	1
357	Blastic NK-cell lymphomas (agranular CD4+CD56+ hematodermic neoplasms): a review. <i>American Journal of Clinical Pathology</i> , 2005 , 123, 662-75	1.9	54
356	CD56+ hematological neoplasms presenting in the skin: a retrospective analysis of 23 new cases and 130 cases from the literature. <i>Annals of Oncology</i> , 2004 , 15, 1097-108	10.3	165
355	Aberrant expression of the tyrosine kinase receptor EphA4 and the transcription factor twist in Sjögren syndrome identified by gene expression analysis. <i>Cancer Research</i> , 2004 , 64, 5578-86	10.1	140
354	Anthracycline-based chemotherapy as primary treatment for intravascular lymphoma. <i>Annals of Oncology</i> , 2004 , 15, 1215-21	10.3	97
353	Intravascular lymphoma: clinical presentation, natural history, management and prognostic factors in a series of 38 cases, with special emphasis on the 'cutaneous variant'. <i>British Journal of Haematology</i> , 2004 , 127, 173-83	4.5	413
352	Chromosomal aberration patterns differ in subtypes of primary cutaneous B cell lymphomas. <i>Journal of Investigative Dermatology</i> , 2004 , 122, 1495-502	4.3	46
351	Molecular cytogenetic analysis of chromosomal breakpoints in the IGH, MYC, BCL6, and MALT1 gene loci in primary cutaneous B-cell lymphomas. <i>Journal of Investigative Dermatology</i> , 2004 , 123, 213-9	4.3	87
350	Internalization and cell cycle-dependent killing of leukemic cells by Gemtuzumab Ozogamicin: rationale for efficacy in CD33-negative malignancies with endocytic capacity. <i>Leukemia</i> , 2004 , 18, 316-25	10.7	114
349	Direct cloning of leukemia-reactive T cells from patients treated with donor lymphocyte infusion shows a relative dominance of hematopoiesis-restricted minor histocompatibility antigen HA-1 and HA-2 specific T cells. <i>Leukemia</i> , 2004 , 18, 798-808	10.7	56
348	Generation of B-cell chronic lymphocytic leukemia (B-CLL)-reactive T-cell lines and clones from HLA class I-matched donors using modified B-CLL cells as stimulators: implications for adoptive immunotherapy. <i>Leukemia</i> , 2004 , 18, 1278-87	10.7	24
347	Site-specific expression of polycomb-group genes encoding the HPC-HPH/PRC1 complex in clinically defined primary nodal and cutaneous large B-cell lymphomas. <i>American Journal of Pathology</i> , 2004 , 164, 533-42	5.8	30
346	Umbilical cord blood-naïve T cells but not adult blood-naïve T cells require HLA class II on antigen-presenting cells for allo-immune activation. <i>Human Immunology</i> , 2004 , 65, 328-39	2.3	7
345	Up-Front Window Trial of Gemtuzumab Ozogamicin (GO) in Previously Untreated Elderly Patients with AML: An EORTC Leukemia Group Study.. <i>Blood</i> , 2004 , 104, 877-877	2.2	2
344	TCL1 and CLA expression in agranular CD4/CD56 hematodermic neoplasms (blastic NK-cell lymphomas) and leukemia cutis. <i>American Journal of Clinical Pathology</i> , 2004 , 122, 307-13	1.9	20
343	Primary Allogeneic T-Cell Responses Against In Vitro Modified Mantle Cell Lymphoma (MCL) Cells: Implications for Adoptive Immunotherapy after HLA-Matched Stem Cell Transplantation (SCT).. <i>Blood</i> , 2004 , 104, 2118-2118	2.2	
342	Retroviral Interference with Execution Pathways in Target Cells To Unravel Mechanisms of Cytotoxic T Cell-Mediated Cell Death.. <i>Blood</i> , 2004 , 104, 1267-1267	2.2	
341	Hematopoiesis-restricted minor histocompatibility antigens HA-1- or HA-2-specific T cells can induce complete remissions of relapsed leukemia. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003 , 100, 2742-7	11.5	361

340	EBV-positive cutaneous B-cell lymphoproliferative disease after imatinib mesylate. <i>Blood</i> , 2003 , 102, 4243	2.2	29
339	Peripheral T-cell lymphomas unspecified presenting in the skin: analysis of prognostic factors in a group of 82 patients. <i>Blood</i> , 2003 , 102, 2213-9	2.2	181
338	Expression profiling reveals that methylation of TIMP3 is involved in uveal melanoma development. <i>International Journal of Cancer</i> , 2003 , 106, 472-479	7.5	78
337	Effect of smoking and sun on the aging skin. <i>Journal of Investigative Dermatology</i> , 2003 , 120, 548-54	4.3	111
336	The influence of painful sunburns and lifetime sun exposure on the risk of actinic keratoses, seborrheic warts, melanocytic nevi, atypical nevi, and skin cancer. <i>Journal of Investigative Dermatology</i> , 2003 , 120, 1087-93	4.3	234
335	Long-term safety aspects of systemic therapy with fumaric acid esters in severe psoriasis. <i>British Journal of Dermatology</i> , 2003 , 149, 363-9	4	157
334	Aleukaemic leukaemia cutis in a patient with acute T-cell lymphoblastic leukaemia. <i>Clinical and Experimental Dermatology</i> , 2003 , 28, 330-2	1.8	6
333	Bcl-2, Bcl-6 and CD10 expression in cutaneous B-cell lymphoma: further support for a follicle centre cell origin and differential diagnostic significance. <i>British Journal of Dermatology</i> , 2003 , 149, 1183-91	4	128
332	The stem cell mobilizing capacity of patients with acute myeloid leukemia in complete remission correlates with relapse risk: results of the EORTC-GIMEMA AML-10 trial. <i>Leukemia</i> , 2003 , 17, 60-7	10.7	55
331	High susceptibility of human leukemic cells to Fas-induced apoptosis is restricted to G1 phase of the cell cycle and can be increased by interferon treatment. <i>Leukemia</i> , 2003 , 17, 576-84	10.7	27
330	The presence of an HLA-identical sibling donor has no impact on outcome of patients with high-risk MDS or secondary AML (sAML) treated with intensive chemotherapy followed by transplantation: results of a prospective study of the EORTC, EBMT, SAKK and GIMEMA Leukemia Groups (EORTC study 06921). <i>Leukemia</i> , 2003 , 17, 859-68	10.7	34
329	Recruitment of leukemic cells from G0 phase of the cell cycle by interferons results in conversion of resistance to daunorubicin. <i>Leukemia</i> , 2003 , 17, 2049-51	10.7	12
328	Primary cutaneous CD30-positive lymphoproliferative disorders. <i>Hematology/Oncology Clinics of North America</i> , 2003 , 17, 1319-32, vii-viii	3.1	27
327	Cutaneous T-cell lymphoma: epidemiology, etiology, and classification. <i>Leukemia and Lymphoma</i> , 2003 , 44 Suppl 3, S49-54	1.9	35
326	MICA gene polymorphism is not associated with an increased risk for skin cancer. <i>Journal of Investigative Dermatology</i> , 2002 , 118, 686-91	4.3	12
325	Cost analysis of HLA-identical sibling and voluntary unrelated allogeneic bone marrow and peripheral blood stem cell transplantation in adults with acute myelocytic leukaemia or acute lymphoblastic leukaemia. <i>Bone Marrow Transplantation</i> , 2002 , 30, 243-51	4.4	44
324	Chemotherapy only compared to chemotherapy followed by transplantation in high risk myelodysplastic syndrome and secondary acute myeloid leukemia; two parallel studies adjusted for various prognostic factors. <i>Leukemia</i> , 2002 , 16, 1615-21	10.7	52
323	Proteolytic enzyme levels are increased during granulocyte colony-stimulating factor-induced hematopoietic stem cell mobilization in human donors but do not predict the number of mobilized stem cells. <i>Journal of Hematotherapy and Stem Cell Research</i> , 2002 , 11, 513-21		21

322	Neutrophils are indispensable for hematopoietic stem cell mobilization induced by interleukin-8 in mice. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002 , 99, 6228-33	11.5	147
321	Cost analysis of CHOP (-like) chemotherapy regimens for patients with newly diagnosed aggressive non-Hodgkin's lymphoma. <i>European Journal of Haematology</i> , 2002 , 69, 213-20	3.8	6
320	Follicular mycosis fungoides, a distinct disease entity with or without associated follicular mucinosis: a clinicopathologic and follow-up study of 51 patients. <i>Archives of Dermatology</i> , 2002 , 138, 191-8		212
319	Differential expression of thymus and activation regulated chemokine and its receptor CCR4 in nodal and cutaneous anaplastic large-cell lymphomas and Hodgkin's disease. <i>Modern Pathology</i> , 2002 , 15, 838-44	9.8	35
318	Minor histocompatibility antigens as targets of graft-versus-leukemia reactions. <i>Current Opinion in Hematology</i> , 2002 , 9, 497-502	3.3	38
317	Transplantation of peripheral blood stem cells as compared with bone marrow from HLA-identical siblings in adult patients with acute myeloid leukemia and acute lymphoblastic leukemia. <i>Journal of Clinical Oncology</i> , 2002 , 20, 4655-64	2.2	129
316	Update on erythrodermic cutaneous T-cell lymphoma: report of the International Society for Cutaneous Lymphomas. <i>Journal of the American Academy of Dermatology</i> , 2002 , 46, 95-106	4.5	374
315	Total skin electron radiation in the management of mycosis fungoides: Consensus of the European Organization for Research and Treatment of Cancer (EORTC) Cutaneous Lymphoma Project Group. <i>Journal of the American Academy of Dermatology</i> , 2002 , 47, 364-70	4.5	147
314	A novel splice variant of the Fas gene in patients with cutaneous T-cell lymphoma. <i>Cancer Research</i> , 2002 , 62, 5389-92	10.1	44
313	Ex vivo expansion and engraftment potential of cord blood-derived CD34+ cells in NOD/SCID mice. <i>Annals of the New York Academy of Sciences</i> , 2001 , 938, 9-17	6.5	25
312	Intensive chemotherapy followed by allogeneic or autologous stem cell transplantation for patients with myelodysplastic syndromes (MDSs) and acute myeloid leukemia following MDS. <i>Blood</i> , 2001 , 98, 2326-31	2.2	130
311	Lymphomatoid papulosis in children: a study of 10 children registered by the Dutch Cutaneous Lymphoma Working Group. <i>British Journal of Dermatology</i> , 2001 , 144, 351-4	4	44
310	Lymphomatoid papulosis with a natural killer-cell phenotype. <i>British Journal of Dermatology</i> , 2001 , 145, 318-22	4	33
309	The Polycomb group protein EZH2 is upregulated in proliferating, cultured human mantle cell lymphoma. <i>British Journal of Haematology</i> , 2001 , 112, 950-8	4.5	174
308	Progress in allogeneic bone marrow and peripheral blood stem cell transplantation for multiple myeloma: a comparison between transplants performed 1983--93 and 1994--8 at European Group for Blood and Marrow Transplantation centres. <i>British Journal of Haematology</i> , 2001 , 113, 209-16	4.5	259
307	Expression of co-stimulatory and adhesion molecules and chemokine or apoptosis receptors on acute myeloid leukaemia: high CD40 and CD11a expression correlates with poor prognosis. <i>British Journal of Haematology</i> , 2001 , 115, 298-308	4.5	29
306	Cyclosporin A combined with vincristine, doxorubicin and dexamethasone (VAD) compared with VAD alone in patients with advanced refractory multiple myeloma: an EORTC-HOVON randomized phase III study (06914). <i>British Journal of Haematology</i> , 2001 , 115, 895-902	4.5	72
305	Processing of natural and recombinant CXCR3-targeting chemokines and implications for biological activity. <i>FEBS Journal</i> , 2001 , 268, 4992-9		17

304	Melanocortin 1 receptor (MC1R) gene variants are associated with an increased risk for cutaneous melanoma which is largely independent of skin type and hair color. <i>Journal of Investigative Dermatology</i> , 2001 , 117, 294-300	4.3	307
303	Willem A. van Vloten zum 60. Geburtstag. <i>Hautarzt</i> , 2001 , 52, 547-548	1.1	
302	Comparison of allogeneic T cell-depleted peripheral blood stem cell and bone marrow transplantation: effect of stem cell source on short- and long-term outcome. <i>Bone Marrow Transplantation</i> , 2001 , 27, 1053-8	4.4	28
301	Similar myeloid recovery despite superior overall engraftment in NOD/SCID mice after transplantation of human CD34(+) cells from umbilical cord blood as compared to adult sources. <i>Bone Marrow Transplantation</i> , 2001 , 28, 163-71	4.4	23
300	Monitoring of engraftment and progression of acute lymphoblastic leukemia in individual NOD/SCID mice. <i>Experimental Hematology</i> , 2001 , 29, 322-9	3.1	45
299	Dual HLA class I and class II restricted recognition of alloreactive T lymphocytes mediated by a single T cell receptor complex. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2001 , 98, 6806-11	11.5	75
298	Commentary on Cutaneous Manifestations of Lymphoma: A Clinical Guide Based on the WHO Classification <i>Clinical Lymphoma and Myeloma</i> , 2001 , 2, 101-102		6
297	Should primary cutaneous Ki-1(CD30)-positive anaplastic large cell lymphoma in childhood be treated with multiple-agent chemotherapy?. <i>Journal of the American Academy of Dermatology</i> , 2001 , 45, 638-40	4.5	4
296	Costs of diagnosis, treatment, and follow up of patients with acute myeloid leukemia in the Netherlands. <i>Journal of Hematology and Stem Cell Research</i> , 2001 , 10, 187-92		18
295	CD8+ T cells in cutaneous T-cell lymphoma: expression of cytotoxic proteins, Fas Ligand, and killing inhibitory receptors and their relationship with clinical behavior. <i>Journal of Clinical Oncology</i> , 2001 , 19, 4322-9	2.2	80
294	Prognostic factors in primary cutaneous large B-cell lymphomas: a European multicenter study. <i>Journal of Clinical Oncology</i> , 2001 , 19, 3602-10	2.2	198
293	Successful outcome with a "quintuple approach" of posttransplant lymphoproliferative disorder. <i>Transplantation</i> , 2001 , 71, 47-52	1.8	31
292	Absence of T(H)2 cytokine messenger RNA expression in CD30-negative primary cutaneous large T-cell lymphomas. <i>Archives of Dermatology</i> , 2001 , 137, 901-5		5
291	Engraftment potential into NOD/SCID mice of CD34+ cells derived from human fetal liver as compared to fetal bone marrow. <i>Haematologica</i> , 2001 , 86, 1021-8	6.6	5
290	Mycosis fungoides: disease evolution and prognosis of 309 Dutch patients. <i>Archives of Dermatology</i> , 2000 , 136, 504-10		257
289	Primary cutaneous lymphomas. <i>Current Opinion in Oncology</i> , 2000 , 12, 419-25	4.2	11
288	Frequency of central nervous system involvement in primary cutaneous B-cell lymphoma. <i>Cancer</i> , 2000 , 89, 913-9	6.4	32
287	Expression of Fas and Fas-ligand in primary cutaneous T-cell lymphoma (CTCL): association between lack of Fas expression and aggressive types of CTCL. <i>British Journal of Dermatology</i> , 2000 , 143, 313-9	4	92

286	HLA-DR4 is associated with a diminished risk of the development of chronic myeloid leukemia (CML). Chronic Leukemia Working Party of the European Blood and Marrow Transplant Registry. <i>Leukemia</i> , 2000 , 14, 859-62	10.7	25
285	Mantle cell lymphoma proliferates upon IL-10 in the CD40 system. <i>Leukemia</i> , 2000 , 14, 1483-9	10.7	44
284	A novel RT-PCR-based protein activity truncation assay for direct assessment of deoxycytidine kinase in small numbers of purified leukemic cells. <i>Leukemia</i> , 2000 , 14, 1678-84	10.7	6
283	Expression and induction of costimulatory and adhesion molecules on acute myeloid leukemic cells: implications for adoptive immunotherapy. <i>Experimental Hematology</i> , 2000 , 28, 161-8	3.1	41
282	Primary and secondary cutaneous CD30+lymphoproliferative disorders: a report from the Dutch Cutaneous Lymphoma Group on the long-term follow-up data of 219 patients and guidelines for diagnosis and treatment. <i>Blood</i> , 2000 , 95, 3653-3661	2.2	617
281	DFFRY codes for a new human male-specific minor transplantation antigen involved in bone marrow graft rejection. <i>Blood</i> , 2000 , 95, 1100-1105	2.2	106
280	EORTC Classification for primary cutaneous lymphomas: A comparison with the R.E.A.L. Classification and the proposed WHO Classification. <i>Annals of Oncology</i> , 2000 , 11, S11-S15	10.3	17
279	The generation of dendritic-like cells with increased allostimulatory function from acute myeloid leukemia cells of various FAB subclasses. <i>Human Immunology</i> , 2000 , 61, 565-74	2.3	57
278	Effect of centre on outcome of bone-marrow transplantation for acute myeloid leukaemia. Acute Leukaemia Working Party of the European Group for Blood and Marrow Transplantation. <i>Lancet, The</i> , 2000 , 355, 1393-8	4.0	84
277	Rationale of a new classification for the group of primary cutaneous lymphomas. <i>Seminars in Cutaneous Medicine and Surgery</i> , 2000 , 19, 71-7	1.4	6
276	Rationale of a New Classification for the Group of Primary Cutaneous Lymphomas. <i>Seminars in Cutaneous Medicine and Surgery</i> , 2000 , 19, 71-77	1.4	5
275	DFFRY codes for a new human male-specific minor transplantation antigen involved in bone marrow graft rejection. <i>Blood</i> , 2000 , 95, 1100-5	2.2	29
274	Primary and secondary cutaneous CD30(+) lymphoproliferative disorders: a report from the Dutch Cutaneous Lymphoma Group on the long-term follow-up data of 219 patients and guidelines for diagnosis and treatment. <i>Blood</i> , 2000 , 95, 3653-61	2.2	87
273	High incidence of alternatively spliced forms of deoxycytidine kinase in patients with resistant acute myeloid leukemia. <i>Blood</i> , 2000 , 96, 1517-24	2.2	15
272	UTY gene codes for an HLA-B60-restricted human male-specific minor histocompatibility antigen involved in stem cell graft rejection: characterization of the critical polymorphic amino acid residues for T-cell recognition. <i>Blood</i> , 2000 , 96, 3126-32	2.2	38
271	Treatment of multifocal primary cutaneous B-cell lymphoma: a clinical follow-up study of 29 patients. <i>Journal of Clinical Oncology</i> , 1999 , 17, 2471-8	2.2	85
270	Prevention of interleukin-8-induced mobilization of hematopoietic progenitor cells in rhesus monkeys by inhibitory antibodies against the metalloproteinase gelatinase B (MMP-9). <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999 , 96, 10863-8	11.5	208
269	Secondary T-acute lymphoblastic leukaemia mimicking blast crisis in chronic myeloid leukaemia. <i>British Journal of Haematology</i> , 1999 , 106, 104-6	4.5	12

268	Human IP-9: A keratinocyte-derived high affinity CXC-chemokine ligand for the IP-10/Mig receptor (CXCR3). <i>Journal of Investigative Dermatology</i> , 1999 , 112, 716-22	4.3	122
267	The CXCR3 activating chemokines IP-10, Mig, and IP-9 are expressed in allergic but not in irritant patch test reactions. <i>Journal of Investigative Dermatology</i> , 1999 , 113, 574-8	4.3	103
266	Biology of IL-8-induced stem cell mobilization. <i>Annals of the New York Academy of Sciences</i> , 1999 , 872, 71-82	6.5	61
265	Multicenter randomized phase II trial of idarubicin vs mitoxantrone, combined with VP-16 and cytarabine for induction/consolidation therapy, followed by a feasibility study of autologous peripheral blood stem cell transplantation in elderly patients with acute myeloid leukemia. <i>Leukemia</i> , 1999 , 13, 843-9	10.7	49
264	The number of nucleated cells reflects the hematopoietic content of umbilical cord blood for transplantation. <i>Bone Marrow Transplantation</i> , 1999 , 24, 965-70	4.4	28
263	Primary cutaneous CD8-positive epidermotropic cytotoxic T cell lymphomas. A distinct clinicopathological entity with an aggressive clinical behavior. <i>American Journal of Pathology</i> , 1999 , 155, 483-92	5.8	399
262	Expression of cytotoxic proteins by neoplastic T cells in mycosis fungoides increases with progression from plaque stage to tumor stage disease. <i>American Journal of Pathology</i> , 1999 , 154, 1203-10	5.8	62
261	Impaired expression of CD28 on T cells in hairy cell leukemia. <i>Clinical Immunology</i> , 1999 , 93, 256-62	9	18
260	IP-10 mRNA expression in cultured keratinocytes is suppressed by inhibition of protein kinase-C and tyrosine kinase and elevation of cAMP. <i>Cytokine</i> , 1999 , 11, 469-75	4	14
259	Pentostatin in T-cell malignancies--a phase II trial of the EORTC. Leukemia Cooperative Group. <i>Annals of Oncology</i> , 1999 , 10, 1493-8	10.3	33
258	EORTC classification for primary cutaneous lymphomas: the best guide to good clinical management. European Organization for Research and Treatment of Cancer. <i>American Journal of Dermatopathology</i> , 1999 , 21, 265-73	0.9	60
257	Mechanisms underlying hematopoietic stem cell mobilization induced by the CXC chemokine interleukin-8. <i>Current Opinion in Hematology</i> , 1999 , 6, 152-8	3.3	21
256	Murine hematopoietic progenitor cells with colony-forming or radioprotective capacity lack expression of the beta 2-integrin LFA-1. <i>Blood</i> , 1999 , 93, 107-12	2.2	9
255	Hairy cell leukemia-specific recognition by multiple autologous HLA-DQ or DP-restricted T-cell clones. <i>Blood</i> , 1999 , 93, 251-9	2.2	4
254	HLA-B8 and HLA-A3 coexpressed with HLA-B8 are associated with a reduced risk of the development of chronic myeloid leukemia. The Chronic Leukemia Working Party of the EBMT. <i>Blood</i> , 1999 , 93, 3863-5	2.2	52
253	Complete remission of accelerated phase chronic myeloid leukemia by treatment with leukemia-reactive cytotoxic T lymphocytes. <i>Blood</i> , 1999 , 94, 1201-8	2.2	51
252	Selective response of CD5+ B cell malignancies to activation of the CD72 antigen. <i>Clinical Immunology and Immunopathology</i> , 1998 , 87, 42-9		2
251	The influence of HLA-matched sibling donor availability on treatment outcome for patients with AML: an analysis of the AML 8A study of the EORTC Leukaemia Cooperative Group and GIMEMA. European Organization for Research and Treatment of Cancer. Gruppo Italiano Malattie Ematologiche Maligne dell'Adulto. <i>British Journal of Haematology</i> , 1999 , 108, 1344-53	4.5	66

250	Generation of leukemia-reactive cytotoxic T lymphocytes from HLA-identical donors of patients with chronic myeloid leukemia using modifications of a limiting dilution assay. <i>Bone Marrow Transplantation</i> , 1998 , 21, 553-60	4.4	29
249	Unconventional rapid Erk1,2 activation is indispensable for proliferation of the growth factor-independent myeloid leukemic cell line KG1. <i>Leukemia</i> , 1998 , 12, 699-704	10.7	6
248	Sublocalization of the breakpoints of a t(5;16) in myelodysplasia. <i>Cancer Genetics and Cytogenetics</i> , 1998 , 100, 5-9		2
247	Assessment of telomere length in hematopoietic interphase cells using in situ hybridization and digital fluorescence microscopy. <i>Cytometry</i> , 1998 , 32, 163-169		37
246	Chemokine IP-10 expression in cultured human keratinocytes. <i>Archives of Dermatological Research</i> , 1998 , 290, 335-41	3.3	52
245	Epidermal interferon-gamma inducible protein-10 (IP-10) and monokine induced by gamma-interferon (Mig) but not IL-8 mRNA expression is associated with epidermotropism in cutaneous T cell lymphomas. <i>Journal of Investigative Dermatology</i> , 1998 , 111, 222-6	4.3	52
244	Effects of thrombopoietin on the proliferation and differentiation of primitive and mature haemopoietic progenitor cells in cord blood. <i>British Journal of Haematology</i> , 1998 , 101, 425-35	4.5	43
243	Pentoxifylline and skin inflammation. <i>Clinical and Experimental Dermatology</i> , 1998 , 23, 168-72	1.8	22
242	Classification of cutaneous lymphomas: crosstalk between pathologist and clinician. <i>Current Diagnostic Pathology</i> , 1998 , 5, 23-33		16
241	T cells recognizing leukemic CD34(+) progenitor cells mediate the antileukemic effect of donor lymphocyte infusions for relapsed chronic myeloid leukemia after allogeneic stem cell transplantation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998 , 95, 10153-7	11.5	82
240	The early phase of engraftment after murine blood cell transplantation is mediated by hematopoietic stem cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998 , 95, 725-9	11.5	99
239	Differences in Epstein-Barr virus expression between primary and secondary cutaneous angiocentric lymphomas. French Study Group of Cutaneous Lymphomas. <i>Archives of Dermatology</i> , 1998 , 134, 479-84		18
238	CAMPATH-1H monoclonal antibody in therapy for previously treated low-grade non-Hodgkin's lymphomas: a phase II multicenter study. European Study Group of CAMPATH-1H Treatment in Low-Grade Non-Hodgkin's Lymphoma. <i>Journal of Clinical Oncology</i> , 1998 , 16, 3257-63	2.2	245
237	bcl-2 protein expression in primary cutaneous large B-cell lymphoma is site-related. <i>Journal of Clinical Oncology</i> , 1998 , 16, 2080-5	2.2	112
236	VH Gene Analysis of Primary Cutaneous B-Cell Lymphomas: Evidence for Ongoing Somatic Hypermutation and Isotype Switching. <i>Blood</i> , 1998 , 92, 3857-3864	2.2	64
235	Anti-LFA-1 blocking antibodies prevent mobilization of hematopoietic progenitor cells induced by interleukin-8. <i>Blood</i> , 1998 , 91, 4099-105	2.2	17
234	Correction of abnormal T-cell receptor repertoire during interferon-alpha therapy in patients with hairy cell leukemia. <i>Blood</i> , 1998 , 91, 4224-31	2.2	4
233	VH gene analysis of primary cutaneous B-cell lymphomas: evidence for ongoing somatic hypermutation and isotype switching. <i>Blood</i> , 1998 , 92, 3857-64	2.2	7

232	Differential involvement of caspases in apoptosis of myeloid leukemic cells induced by chemotherapy versus growth factor withdrawal. <i>FEBS Letters</i> , 1997 , 409, 207-10	3.8	26
231	Relation between expression of adhesion molecules and clinical behavior in cutaneous follicle center cell lymphomas. <i>Journal of the American Academy of Dermatology</i> , 1997 , 37, 34-40	4.5	16
230	Generation of dendritic cells expressing bcr-abl from CD34-positive chronic myeloid leukemia precursor cells. <i>Human Immunology</i> , 1997 , 53, 216-23	2.3	80
229	Intensive postremission chemotherapy without maintenance therapy in adults with acute lymphoblastic leukemia. Dutch Hemato-Oncology Research Group. <i>Journal of Clinical Oncology</i> , 1997 , 15, 476-82	2.2	55
228	Treatment of T-cell prolymphocytic leukemia with human CD52 antibody. <i>Journal of Clinical Oncology</i> , 1997 , 15, 2667-72	2.2	179
227	p53 and bcl-2 expression do not correlate with prognosis in primary cutaneous large T-cell lymphomas. <i>Journal of Cutaneous Pathology</i> , 1997 , 24, 462-7	1.7	18
226	Cytotoxic T-lymphocyte (CTL) responses against acute or chronic myeloid leukemia. <i>Immunological Reviews</i> , 1997 , 157, 223-30	11.3	61
225	Most primary cutaneous CD30-positive lymphoproliferative disorders have a CD4-positive cytotoxic T-cell phenotype. <i>Journal of Investigative Dermatology</i> , 1997 , 109, 636-40	4.3	110
224	TCR gamma delta+ cells expressing Vgamma 9V delta2, which normally predominate the blood, are found in the spleens of patients with hairy cell leukemia. <i>Leukemia</i> , 1997 , 11, 106-9	10.7	16
223	Allogeneic bone marrow transplantation for acute leukemia in patients over the age of 40 years. Acute Leukemia Working Party of the European Group for Bone Marrow Transplantation (EBMT). <i>Leukemia</i> , 1997 , 11, 416-9	10.7	63
222	Mosaicism of the 5q deletion as assessed by interphase FISH is a common phenomenon in MDS and restricted to myeloid cells. <i>Leukemia</i> , 1997 , 11, 519-23	10.7	24
221	Quality of life in patients with acute myelogenous leukemia in prolonged first complete remission after bone marrow transplantation (allogeneic or autologous) or chemotherapy: a cross-sectional study of the EORTC-GIMEMA AML 8A trial. <i>Bone Marrow Transplantation</i> , 1997 , 20, 307-15	4.4	92
220	Problems and possible solutions in finding an unrelated bone marrow donor. Results of consecutive searches for 240 Dutch patients. <i>Bone Marrow Transplantation</i> , 1997 , 20, 1011-7	4.4	36
219	Pentoxifylline inhibits human T-cell adhesion to dermal endothelial cells. <i>Archives of Dermatological Research</i> , 1997 , 289, 189-93	3.3	18
218	The role of cytokines and hematopoietic growth factors in the autocrine/paracrine regulation of inducible hematopoiesis. <i>Annals of Hematology</i> , 1997 , 75, 27-31	3	19
217	Absence of mutations in the RET gene in acute myeloid leukemia. <i>Annals of Hematology</i> , 1997 , 75, 87-90	3	11
216	Extramedullary hematopoietic growth factor and cytokine gene expression indicate endocrine regulation of hematopoiesis in patients with acute appendicitis. <i>Annals of Hematology</i> , 1997 , 75, 201-5	3	3
215	Leukemia inhibitory factor induces in vivo expansion of bone marrow progenitor cells that accelerate hematopoietic reconstitution but do not enhance radioprotection in lethally irradiated mice. <i>Stem Cells</i> , 1997 , 15, 50-5	5.8	6

214	T cells selectively infiltrate bone marrow areas with residual haemopoiesis of patients with acquired aplastic anaemia. <i>British Journal of Haematology</i> , 1997 , 99, 517-9	4.5	25
213	Analysis of T-cell clonality in bone marrow of patients with acquired aplastic anaemia. <i>British Journal of Haematology</i> , 1997 , 96, 85-91	4.5	23
212	HOXC4, HOXC5, and HOXC6 expression in primary cutaneous lymphoid lesions. High expression of HOXC5 in anaplastic large-cell lymphomas. <i>American Journal of Pathology</i> , 1997 , 151, 1067-74	5.8	12
211	Immunophenotyping and gene rearrangement analysis provide additional criteria to differentiate between cutaneous T-cell lymphomas and pseudo-T-cell lymphomas. <i>American Journal of Pathology</i> , 1997 , 150, 1941-9	5.8	47
210	EORTC classification for primary cutaneous lymphomas: a proposal from the Cutaneous Lymphoma Study Group of the European Organization for Research and Treatment of Cancer. <i>Blood</i> , 1997 , 90, 354-71 ²	7.1 ²	802
209	Molecular remission in PML/RAR alpha-positive acute promyelocytic leukemia by combined all-trans retinoic acid and idarubicin (AIDA) therapy. Gruppo Italiano-Malattie Ematologiche Maligne dell'Adulto and Associazione Italiana di Ematologia ed Oncologia Pediatrica Cooperative Groups. <i>Blood</i> , 1997 , 89, 1811-21	2.2	348
208	Characterization of a novel malignant B cell line with t(14;18) and t(4;11) established from a patient with acute monoblastic leukemia. <i>Experimental Hematology</i> , 1997 , 25, 1062-8	3.1	3
207	HOXC4, HOXC5, and HOXC6 expression in non-Hodgkin's lymphoma: preferential expression of the HOXC5 gene in primary cutaneous anaplastic T-cell and oro-gastrointestinal tract mucosa-associated B-cell lymphomas. <i>Blood</i> , 1997 , 90, 4116-25	2.2	5
206	Is mycosis fungoides exacerbated by fluoxetine?. <i>Journal of the American Academy of Dermatology</i> , 1996 , 35, 635-6	4.5	14
205	The role of the CD40 antigen on malignant B cells. <i>Leukemia and Lymphoma</i> , 1996 , 22, 229-35	1.9	9
204	Treatment of primary cutaneous B-cell lymphomas of follicle center cell origin: a clinical follow-up study of 55 patients treated with radiotherapy or polychemotherapy. <i>Journal of Clinical Oncology</i> , 1996 , 14, 549-55	2.2	116
203	Detection of partial cDNA sequences differentially expressed in patients with myelodysplasia. <i>Annals of Hematology</i> , 1996 , 72, 231-6	3	3
202	Interleukin-10, interleukin-12, and tumor necrosis factor-alpha differentially influence the proliferation of human CD8+ and CD4+ T-cell clones. <i>Annals of Hematology</i> , 1996 , 72, 245-52	3	14
201	Umbilical cord blood contains normal frequencies of cytotoxic T-lymphocyte precursors (ctlp) and helper T-lymphocyte precursors against noninherited maternal antigens and noninherited paternal antigens. <i>Annals of Hematology</i> , 1996 , 72, 260-4	3	19
200	Accelerated platelet reconstitution following transplantation of bone marrow cells derived from IL-6-treated donor mice. <i>Annals of Hematology</i> , 1996 , 73, 239-45	3	8
199	Aplastic anaemia patients with clonal X-chromosome inactivation pattern in haemopoietic cells exhibit polyclonal TCRgamma and IgH gene rearrangements. <i>British Journal of Haematology</i> , 1996 , 93, 326-32	4.5	7
198	Haemopoietic growth factor tyrosine kinase receptor expression profiles in normal haemopoiesis. <i>British Journal of Haematology</i> , 1996 , 94, 236-41	4.5	6
197	Proliferation of precursor B-lineage acute lymphoblastic leukaemia by activating the CD40 antigen. <i>British Journal of Haematology</i> , 1996 , 95, 319-26	4.5	21

196	Colony growth of human hematopoietic progenitor cells in the absence of serum is supported by a proteinase inhibitor identified as antileukoproteinase. <i>Journal of Experimental Medicine</i> , 1996 , 184, 1305-12	16.6	16
195	Persistent clonal excess and skewed T-cell repertoire in T cells from patients with hairy cell leukemia. <i>Blood</i> , 1996 , 87, 3795-802	2.2	9
194	Proliferation of B cell malignancies in all stages of differentiation upon stimulation in the 'CD40 system'. <i>Leukemia</i> , 1996 , 10, 488-93	10.7	31
193	Anti-CD45 and anti-CD52 (Campath) monoclonal antibodies effectively eliminate systematically disseminated human non-Hodgkin's lymphoma B cells in Scid mice. <i>Experimental Hematology</i> , 1996 , 24, 919-26	3.1	6
192	Direct display of hematopoietic tyrosine kinase receptor expression profiles in KG1 cells by PCR using degenerate primers. <i>Leukemia</i> , 1996 , 10, 1395-9	10.7	3
191	Mycosis fungoides with extracutaneous localization in the breast. <i>British Journal of Dermatology</i> , 1996 , 134, 1125-30	4	1
190	Unrelated bone marrow transplantation as a rescue procedure following inadvertent loss of an autologous bone marrow graft. <i>Bone Marrow Transplantation</i> , 1996 , 18, 461-3	4.4	3
189	Improved survival of lethally irradiated recipient mice transplanted with circulating progenitor cells mobilized by IL-8 after pretreatment with stem cell factor. <i>Experimental Hematology</i> , 1996 , 24, 1387-93	3.1	28
188	Primary cutaneous large B-cell lymphomas of the legs. A distinct type of cutaneous B-cell lymphoma with an intermediate prognosis. Dutch Cutaneous Lymphoma Working Group. <i>Archives of Dermatology</i> , 1996 , 132, 1304-8		22
187	In vitro-induced resistance to the deoxycytidine analogues cytarabine (AraC) and 5-aza-2'-deoxycytidine (DAC) in a rat model for acute myeloid leukemia is mediated by mutations in the deoxycytidine kinase (dck) gene. <i>Annals of Hematology</i> , 1995 , 71, 41-7	3	31
186	High-dose Ara-C for remission induction and consolidation of previously untreated adults with ALL or lymphoblastic lymphoma. <i>Annals of Hematology</i> , 1995 , 70, 71-4	3	17
185	Are most primary cutaneous B-cell lymphomas 'marginal cell lymphomas'?. <i>British Journal of Dermatology</i> , 1995 , 133, 950-2; discussion 953-4	4	29
184	Pentoxifylline inhibits T-cell adherence to keratinocytes. <i>Journal of Investigative Dermatology</i> , 1995 , 104, 1004-7	4.3	13
183	Autologous or allogeneic bone marrow transplantation compared with intensive chemotherapy in acute myelogenous leukemia. European Organization for Research and Treatment of Cancer (EORTC) and the Gruppo Italiano Malattie Ematologiche Maligne dell'Adulto (GIMEMA) Leukemia Cooperative Group. <i>Journal of Clinical Oncology</i> , 1995 , 13, 202-17	59.2	719
182	Modification of rhodamine staining allows identification of hematopoietic stem cells with preferential short-term or long-term bone marrow-repopulating ability. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995 , 92, 8901-5	11.5	92
181	The diagnostic value of morphometry on blood lymphocytes in erythrodermic actinic reticuloid. <i>Archives of Dermatology</i> , 1995 , 131, 1298-303		1
180	De novo induced mutations in the deoxycytidine kinase (dck) gene in rat leukemic clonal cell lines confer resistance to cytarabine (AraC) and 5-aza-2'-deoxycytidine (DAC). <i>Leukemia</i> , 1995 , 9, 1032-8	10.7	20
179	CD30 expression in normal and neoplastic lymphoid tissue: biological aspects and clinical implications. <i>Leukemia</i> , 1995 , 9, 1620-7	10.7	40

178	Accelerated reconstitution of platelets and erythrocytes after syngeneic transplantation of bone marrow cells derived from thrombopoietin pretreated donor mice. <i>Blood</i> , 1995 , 86, 3308-13	2.2	11
177	Multiple minor histocompatibility antigen-specific cytotoxic T lymphocyte clones can be generated during graft rejection after HLA-identical bone marrow transplantation. <i>Bone Marrow Transplantation</i> , 1995 , 16, 125-32	4.4	36
176	Constitutive in vivo cytokine and hematopoietic growth factor gene expression in the bone marrow and peripheral blood of healthy individuals. <i>Blood</i> , 1995 , 85, 2038-44	2.2	2
175	Recognition of clonogenic leukemic cells, remission bone marrow and HLA-identical donor bone marrow by CD8+ or CD4+ minor histocompatibility antigen-specific cytotoxic T lymphocytes. <i>Journal of Clinical Investigation</i> , 1995 , 96, 877-83	15.9	65
174	High-dose Ara-C for remission induction and consolidation of previously untreated adults with ALL or lymphoblastic lymphoma 1995 , 70, 71		1
173	Primary cutaneous B-cell lymphomas. <i>Leukemia and Lymphoma</i> , 1994 , 14, 213-8	1.9	17
172	IL-4 down-regulates IL-2-, IL-3-, and GM-CSF-induced cytokine gene expression in peripheral blood monocytes. <i>Annals of Hematology</i> , 1994 , 68, 293-8	3	9
171	Human growth factor (huGRO), interleukin-8 (IL-8) and interferon-gamma-inducible protein (gamma-IP-10) gene expression in cultured normal human keratinocytes. <i>Archives of Dermatological Research</i> , 1994 , 286, 471-5	3.3	30
170	Classification of primary cutaneous T-cell lymphomas. <i>Histopathology</i> , 1994 , 24, 405-15	7.3	105
169	Cloning of the Dck gene encoding rat deoxycytidine kinase. <i>Gene</i> , 1994 , 150, 351-4	3.8	5
168	Granzyme B-expressing peripheral T-cell lymphomas: neoplastic equivalents of activated cytotoxic T cells with preference for mucosa-associated lymphoid tissue localization. <i>Blood</i> , 1994 , 84, 3785-91	2.2	21
167	Abnormally activated T lymphocytes in the spleen of patients with hairy-cell leukemia. <i>Leukemia</i> , 1994 , 8, 2095-101	10.7	13
166	Cutaneous pseudolymphomas: classification and differential diagnosis. <i>Seminars in Dermatology</i> , 1994 , 13, 187-96		43
165	Classification of primary cutaneous large cell lymphomas. <i>Dermatologic Clinics</i> , 1994 , 12, 361-73	4.2	2
164	Homing and antigenic characterization of a human non-Hodgkin's lymphoma B cell line in severe combined immunodeficient (SCID) mice. <i>Leukemia</i> , 1994 , 8, 1385-91	10.7	7
163	Effect of calcitriol on growth, differentiation, chemokine mRNA expression of cultured keratinocytes and on keratinocyte-T cell binding. <i>Acta Dermato-venereologica Supplementum</i> , 1994 , 186, 52-4		
162	Presence of Epstein-Barr virus in extranodal T-cell lymphomas: differences in relation to site. <i>Blood</i> , 1994 , 83, 1612-8	2.2	12
161	Multiple minor histocompatibility antigen disparities between a recipient and four HLA-identical potential sibling donors for bone marrow transplantation. <i>Human Immunology</i> , 1993 , 37, 221-8	2.3	7

160	Spectrum of primary cutaneous CD30 (Ki-1)-positive lymphoproliferative disorders. A proposal for classification and guidelines for management and treatment. <i>Journal of the American Academy of Dermatology</i> , 1993 , 28, 973-80	4.5	298
159	Primary cutaneous CD30-positive large cell lymphoma: definition of a new type of cutaneous lymphoma with a favorable prognosis. A European Multicenter Study of 47 patients. <i>Cancer</i> , 1993 , 71, 2097-104	6.4	288
158	The human interleukin-6 receptor alpha chain gene is localized on chromosome 1 band q21. <i>Human Genetics</i> , 1993 , 90, 542-4	6.3	17
157	CD44 antibody stimulates adhesion of peripheral blood T cells to keratinocytes through the leukocyte function-associated antigen-1/intercellular adhesion molecule-1 pathway. <i>Journal of Investigative Dermatology</i> , 1993 , 100, 424-8	4.3	21
156	Cutaneous immunocytomas: a clinicopathologic study of 26 cases. <i>Histopathology</i> , 1993 , 23, 117-25	7.3	134
155	Differences in clinical behaviour and immunophenotype between primary cutaneous and primary nodal anaplastic large cell lymphoma of T-cell or null cell phenotype. <i>Histopathology</i> , 1993 , 23, 127-35	7.3	152
154	Biology and treatment of myelodysplastic syndromes--developments in the past decade. <i>Annals of Hematology</i> , 1993 , 66, 107-15	3	21
153	Separation, enrichment, and characterization of human hematopoietic progenitor cells from umbilical cord blood. <i>Annals of Hematology</i> , 1993 , 67, 231-6	3	13
152	Order of human hematopoietic growth factor and receptor genes on the long arm of chromosome 5, as determined by fluorescence in situ hybridization. <i>Annals of Hematology</i> , 1993 , 66, 15-20	3	7
151	Interleukin-9 stimulates the proliferation of enriched human erythroid progenitor cells: additive effect with GM-CSF. <i>Annals of Hematology</i> , 1993 , 66, 45-9	3	13
150	Preliminary results with 5-aza-2'-deoxycytidine (DAC)-containing chemotherapy in patients with relapsed or refractory acute leukemia. The EORTC Leukemia Cooperative Group. <i>Leukemia</i> , 1993 , 7 Suppl 1, 49-50	10.7	14
149	Role of deoxycytidine kinase in an in vitro model for AraC- and DAC-resistance: substrate-enzyme interactions with deoxycytidine, 1-beta-D-arabinofuranosylcytosine and 5-aza-2'-deoxycytidine. <i>Leukemia</i> , 1993 , 7, 1005-11	10.7	23
148	Regulatory effects of T cell lymphokines on cytokine gene expression in monocytes. <i>Lymphokine and Cytokine Research</i> , 1993 , 12, 457-64		14
147	Continuous infusion of interleukin-6 in sublethally irradiated mice accelerates platelet reconstitution and the recovery of myeloid but not of megakaryocytic progenitor cells in bone marrow. <i>Experimental Hematology</i> , 1993 , 21, 1621-7	3.1	10
146	Involvement of the complement system in the pathogenesis of pulmonary leukostasis in experimental myelocytic leukemia. <i>Leukemia</i> , 1993 , 7, 1608-14	10.7	9
145	Inhibitory effect of interleukin-4 on the proliferation of acute myeloid leukemia cells with myelo-monocytic differentiation (AML-M4/M5); the role of interleukin-6. <i>Leukemia</i> , 1993 , 7, 643-5	10.7	8
144	The prognosis of patients with lymphomatoid papulosis associated with malignant lymphomas. <i>British Journal of Dermatology</i> , 1992 , 126, 596-602	4	119
143	Differential expression of the HECA-452 antigen (cutaneous lymphocyte associated antigen, CLA) in cutaneous and non-cutaneous T-cell lymphomas. <i>Histopathology</i> , 1992 , 21, 59-64	7.3	33

142	Response to interferon alfa-2b in a patient with systemic mastocytosis. <i>New England Journal of Medicine</i> , 1992 , 326, 619-23	59.2	192
141	GM-CSF and IL-3 in Myelodysplastic Syndromes at Low Risk of Developing Acute Leukemia (MDS-LR). <i>Leukemia and Lymphoma</i> , 1992 , 7, 64-64	1.9	1
140	Interleukin 6 is a permissive factor for monocytic colony formation by human hematopoietic progenitor cells. <i>Journal of Experimental Medicine</i> , 1992 , 175, 1151-4	16.6	67
139	Generation of leukemia-reactive cytotoxic T lymphocyte clones from the HLA-identical bone marrow donor of a patient with leukemia. <i>Journal of Experimental Medicine</i> , 1992 , 176, 1283-9	16.6	133
138	Prevention of viridans-group streptococcal septicemia in oncohematologic patients: a controlled comparative study on the effect of penicillin G and cotrimoxazole. <i>Annals of Hematology</i> , 1992 , 64, 260-3		26
137	Quantitative skin cultures of <i>Pityrosporum</i> yeasts in patients seropositive for the human immunodeficiency virus with and without seborrheic dermatitis. <i>Journal of the American Academy of Dermatology</i> , 1992 , 27, 37-9	4.5	41
136	Induction of ICAM-1 expression by epidermal keratinocytes via a paracrine pathway possibly involving dermal dendritic cells. <i>Archives of Dermatological Research</i> , 1992 , 284, 250-2	3.3	10
135	Demonstration of clonal immunoglobulin gene rearrangements in cutaneous B-cell lymphomas and pseudo-B-cell lymphomas: differential diagnostic and pathogenetic aspects. <i>Journal of Investigative Dermatology</i> , 1992 , 99, 749-54	4.3	52
134	Oral etoposide in the treatment of cutaneous large-cell lymphomas. A preliminary report of four cases. <i>British Journal of Dermatology</i> , 1992 , 127, 524-8	4	21
133	A randomized phase-I/II multicenter study of recombinant human granulocyte-macrophage colony-stimulating factor (GM-CSF) therapy for patients with myelodysplastic syndromes and a relatively low risk of acute leukemia. EORTC Leukemia Cooperative Group. <i>Annals of Hematology</i> , 1992 , 64, 173-80	3	44
132	Increased risk of second cancers in managing Hodgkins disease: the 20-year Leiden experience. <i>Annals of Hematology</i> , 1992 , 65, 213-8	3	24
131	Increase in Ara-C sensitivity in Ara-C sensitive and -resistant leukemia by stimulation of the salvage and inhibition of the de novo pathway. <i>Annals of Hematology</i> , 1992 , 65, 26-32	3	16
130	Cutaneous pseudo-T-cell lymphomas. A clinicopathologic study of 20 patients. <i>Cancer</i> , 1992 , 69, 717-24	6.4	78
129	Cytomegalovirus antigenemia assay or PCR can be used to monitor ganciclovir treatment in bone marrow transplant recipients. <i>Bone Marrow Transplantation</i> , 1992 , 9, 247-53	4.4	24
128	Sustained engraftment of mice transplanted with IL-1-primed blood-derived stem cells. <i>Journal of Immunology</i> , 1992 , 148, 417-21	5.3	45
127	Frequency and prognostic significance of clonal T-cell receptor beta-gene rearrangements in the peripheral blood of patients with mycosis fungoides. <i>Archives of Dermatology</i> , 1992 , 128, 1602-7		3
126	Production of tumor necrosis factor-alpha by normal and malignant B lymphocytes in response to interferon-alpha, interferon-gamma and interleukin-4. <i>Leukemia</i> , 1992 , 6, 116-9	10.7	24
125	Development of pulmonary leukostasis in experimental myelocytic leukemia in the Brown-Norway rat. <i>Leukemia</i> , 1992 , 6, 142-9	10.7	13

124	Human interleukin for DA cells (HILDA) does not affect the proliferation and differentiation of hematopoietic progenitor cells in human long-term bone marrow cultures. <i>Experimental Hematology</i> , 1992 , 20, 6-10	3.1	9
123	In vivo use of Campath-1G to prevent graft-versus-host disease and graft rejection after bone marrow transplantation. <i>Bone Marrow Transplantation</i> , 1992 , 9, 255-61	4.4	25
122	Serum monocyte colony-stimulating factor, erythropoietin and interleukin-6 in relation to pancytopenia in hairy cell leukemia. <i>Leukemia</i> , 1992 , 6, 735-7	10.7	3
121	Differential diagnosis of cutaneous large cell lymphomas using monoclonal antibodies reactive in paraffin-embedded skin biopsy specimens. <i>American Journal of Dermatopathology</i> , 1991 , 13, 342-9	0.9	10
120	The antileukaemic activity of 5-Aza-2 deoxycytidine (Aza-dC) in patients with relapsed and resistant leukaemia. <i>British Journal of Cancer</i> , 1991 , 64, 144-8	8.7	68
119	Diagnostic value of T-cell receptor beta gene rearrangement analysis on peripheral blood lymphocytes of patients with erythroderma. <i>Journal of Investigative Dermatology</i> , 1991 , 97, 782-6	4.3	35
118	Effect of subcutaneously administered human recombinant erythropoietin on erythropoiesis in patients with myelodysplasia. <i>British Journal of Haematology</i> , 1991 , 78, 488-93	4.5	48
117	Neutropenic enterocolitis following treatment with cytosine arabinoside-containing regimens for hematological malignancies: a potentiating role for amsacrine. <i>Annals of Hematology</i> , 1991 , 62, 129-34	3	24
116	Bone marrow transplantation or chemotherapy as post-remission treatment of adult acute myelogenous leukemia. <i>Annals of Hematology</i> , 1991 , 62, 59-63	3	10
115	Correction of neutropenia associated with chronic lymphocytic leukaemia following treatment with granulocyte-macrophage colony-stimulating factor. <i>Annals of Hematology</i> , 1991 , 62, 32-4	3	20
114	The role of interleukin-1 in hematopoiesis. <i>Acta Haematologica</i> , 1991 , 86, 148-54	2.7	28
113	Studies on clonality by PCR analysis of the PGK-1 gene. <i>Nucleic Acids Research</i> , 1991 , 19, 2794	20.1	40
112	Repeat polymorphisms in the interleukin-4 gene (IL4). <i>Nucleic Acids Research</i> , 1991 , 19, 3763	20.1	97
111	Growth inhibition of clonogenic leukemic precursor cells by minor histocompatibility antigen-specific cytotoxic T lymphocytes. <i>Journal of Experimental Medicine</i> , 1991 , 174, 27-33	16.6	104
110	Minor histocompatibility antigen-specific cytotoxic T cell lines, capable of lysing human hematopoietic progenitor cells, can be generated in vitro by stimulation with HLA-identical bone marrow cells. <i>Journal of Experimental Medicine</i> , 1991 , 173, 101-9	16.6	33
109	Mycosis fungoides-like lesions associated with phenytoin and carbamazepine therapy. <i>Journal of the American Academy of Dermatology</i> , 1991 , 24, 216-20	4.5	70
108	Clonal hematopoiesis in patients with acquired aplastic anemia. <i>Blood</i> , 1991 , 78, 3209-14	2.2	8
107	Interleukin 4 down-regulates the expression of CD14 and the production of interleukin 6 in acute myeloid leukemia cells. <i>Lymphokine and Cytokine Research</i> , 1991 , 10, 457-61		4

106	The prognostic significance of chromosomal findings in patients with acute myeloid leukemia in a study comparing the efficacy of autologous and allogeneic bone marrow transplantation. <i>Bone Marrow Transplantation</i> , 1991 , 8, 377-81	4.4	7
105	Cytokine production by the bladder carcinoma cell line 5637: rapid analysis of mRNA expression levels using a cDNA-PCR procedure. <i>Lymphokine and Cytokine Research</i> , 1991 , 10, 231-5		38
104	Experimental myelocytic leukemia in the Brown-Norway rat as a model for pulmonary leukostasis. <i>American Journal of Pathology</i> , 1991 , 138, 777-80	5.8	3
103	Stage is a better prognostic indicator than morphologic subtype in primary noncutaneous T-cell lymphoma. <i>American Journal of Clinical Pathology</i> , 1990 , 93, 49-57	1.9	41
102	The significance of Leu 8 negative T cells in lymphoid skin infiltrates: malignant transformation, selective homing or T-cell activation?. <i>British Journal of Dermatology</i> , 1990 , 123, 587-93	4	10
101	Characterization of the dermal infiltrates in Jessner's lymphocytic infiltrate of the skin, polymorphous light eruption and cutaneous lupus erythematosus: differential diagnostic and pathogenetic aspects. <i>Journal of Cutaneous Pathology</i> , 1990 , 17, 2-8	1.7	36
100	Deoxycytidine kinase, thymidine kinase and cytidine deaminase and the formation of Ara-CTP in leukemic cells in different phases of the cell cycle. <i>Leukemia Research</i> , 1990 , 14, 363-9	2.7	15
99	Respiratory failure elicited by streptococcal septicaemia in patients treated with cytosine arabinoside, and its prevention by penicillin. <i>Infection</i> , 1990 , 18, 131-7	5.8	43
98	Interleukin-4. A regulatory protein. <i>Blut</i> , 1990 , 60, 269-74		33
97	A simplified assay for measurement of cytosine arabinoside incorporation into DNA in Ara-C-sensitive and -resistant leukemic cells. <i>Cancer Chemotherapy and Pharmacology</i> , 1990 , 27, 151-6	3.5	10
96	Differentiation between lymphadenosis benigna cutis and primary cutaneous follicular center cell lymphomas. A comparative clinicopathologic study of 57 patients. <i>Cancer</i> , 1990 , 65, 2301-6	6.4	68
95	Autologous bone marrow transplantation in acute myeloid leukemia in first remission: results of a Dutch prospective study. <i>Journal of Clinical Oncology</i> , 1990 , 8, 287-94	2.2	100
94	Gastrointestinal complications of cytosine-arabinoside chemotherapy: findings on plain abdominal radiographs. <i>American Journal of Roentgenology</i> , 1990 , 154, 95-8	5.4	19
93	Pentostatin (2'-deoxycoformycin) for the treatment of lymphoid neoplasms. <i>Cancer Treatment Reviews</i> , 1990 , 17, 213-5	14.4	12
92	Rejection of bone-marrow graft by recipient-derived cytotoxic T lymphocytes against minor histocompatibility antigens. <i>Lancet, The</i> , 1990 , 335, 131-4	4.0	92
91	Spectrophotometric quantification of cell-cell adherence by an enzyme-linked immuno-cell adhesion assay. <i>Journal of Immunological Methods</i> , 1990 , 132, 51-6	2.5	16
90	Substrate-specific deoxycytidine kinase deficiency in 1-beta-D-arabinofuranosylcytosine-resistant leukemic cells. <i>Cancer Research</i> , 1990 , 50, 6515-9	10.1	16
89	Response to pentostatin in hairy-cell leukemia refractory to interferon-alpha. The European Organization for Research and Treatment of Cancer Leukemia Cooperative Group. <i>Journal of Clinical Oncology</i> , 1989 , 7, 1533-8	2.2	69

88	Inhibition of human macrophage colony formation by interleukin 4. <i>Journal of Experimental Medicine</i> , 1989 , 170, 577-82	16.6	129
87	The hematopoietic activities of interleukin-1. <i>Biotherapy (Dordrecht, Netherlands)</i> , 1989 , 1, 263-71		13
86	The biological activities of interleukin-1. <i>Blut</i> , 1989 , 59, 147-56		36
85	Primary gastrointestinal non-Hodgkin's lymphoma in a population-based registry. <i>British Journal of Cancer</i> , 1989 , 60, 745-50	8.7	94
84	Primary extranodal and nodal non-Hodgkin's lymphoma. A survey of a population-based registry. <i>European Journal of Cancer & Clinical Oncology</i> , 1989 , 25, 1203-10		127
83	Interleukin 1 and poly(rI).poly(rC) induce production of granulocyte CSF, macrophage CSF, and granulocyte-macrophage CSF by human endothelial cells. <i>Experimental Hematology</i> , 1989 , 17, 229-34	3.1	42
82	Jessner's lymphocytic infiltration of the skin. A clinical study of 100 patients. <i>Archives of Dermatology</i> , 1989 , 125, 1525-30		5
81	Post remission treatment in acute myelogenous leukaemia in first remission. The EORTC experience. EORTC Leukemia Cooperative Group. <i>Bone Marrow Transplantation</i> , 1989 , 4 Suppl 3, 72	4.4	
80	Comparison of intensive consolidation therapy and autologous bone marrow transplantation in acute myeloid leukaemia. The Leiden experience. <i>Bone Marrow Transplantation</i> , 1989 , 4 Suppl 3, 73	4.4	
79	Prognostic significance of CD30 (Ki-1/Ber-H2) expression in primary cutaneous large-cell lymphomas of T-cell origin. A clinicopathologic and immunohistochemical study in 20 patients. <i>American Journal of Pathology</i> , 1989 , 135, 1169-78	5.8	113
78	Factors influencing release of granulocyte-macrophage colony-stimulating activity from human mononuclear phagocytes. <i>European Journal of Haematology</i> , 1988 , 41, 352-8	3.8	8
77	Short-term intensive treatment (V.A.A.P.) of adult acute lymphoblastic leukemia and lymphoblastic lymphoma. <i>European Journal of Haematology</i> , 1988 , 41, 489-95	3.8	11
76	Results of induction and consolidation treatment with intermediate and high-dose cytosine arabinoside and m-Amsa of patients with poor-risk acute myelogenous leukaemia. <i>European Journal of Haematology</i> , 1988 , 40, 198-204	3.8	16
75	Evolution of acquired severe aplastic anaemia to myelodysplasia and subsequent leukaemia in adults. <i>British Journal of Haematology</i> , 1988 , 70, 55-62	4.5	107
74	Day-6 bone marrow aspirate for the prediction of response to remission induction therapy for acute myelogenous leukaemia. <i>Blut</i> , 1988 , 57, 91-5		6
73	High-dose cytosine arabinoside: pharmacological and clinical aspects. <i>Blut</i> , 1988 , 56, 1-11		31
72	Leucostasis, an underestimated cause of death in leukaemia. <i>Blut</i> , 1988 , 56, 39-44		30
71	Comparison of the antileukaemic activity of 5 aza-2-deoxycytidine and arabinofuranosyl-cytosine in rats with myelocytic leukaemia. <i>British Journal of Cancer</i> , 1988 , 58, 730-3	8.7	18

70	Radioresistant pseudo-colony formation in the PHA-leukocyte feeder colony assay. <i>Leukemia Research</i> , 1988 , 12, 917-22	2.7	1
69	Intermediate and high dose Ara-C and m-AMSA for remission induction and consolidation treatment of patients with acute myeloid leukemia: an EORTC Leukemia Cooperative Group phase II study. <i>European Journal of Cancer & Clinical Oncology</i> , 1988 , 24, 1721-5		19
68	Activation of the c-myc oncogene in a precursor-B-cell blast crisis of follicular lymphoma, presenting as composite lymphoma. <i>New England Journal of Medicine</i> , 1988 , 318, 1373-8	59.2	218
67	Alopecia mucinosa progressing into mycosis fungoides. A long-term follow-up study of two patients. <i>American Journal of Dermatopathology</i> , 1988 , 10, 478-86	0.9	33
66	Immunohistochemical studies of lymphadenosis benigna cutis occurring in a tattoo. <i>American Journal of Dermatopathology</i> , 1988 , 10, 518-23	0.9	43
65	Deoxycoformycin in therapy of refractory lymphoid neoplasms. <i>Oncology Research and Treatment</i> , 1988 , 11, 35-7	2.8	1
64	Gall bladder perforation in response to chemotherapy in a patient with a histiocytic lymphoma. <i>Netherlands Journal of Medicine</i> , 1988 , 33, 78-82	0.5	2
63	The unrelated bone marrow donor. <i>Bone Marrow Transplantation</i> , 1988 , 3, 371-7	4.4	0
62	Fournier's gangrene or fulminant necrotizing fasciitis of the scrotum and penis as a complication of granulocytopenia in a patient with acute myelogenous leukaemia (AML). <i>Netherlands Journal of Medicine</i> , 1988 , 32, 235-9	0.5	8
61	Colony growth of normal and neoplastic cells in various concentrations of methylcellulose. <i>Experimental Hematology</i> , 1988 , 16, 922-8	3.1	4
60	Pulmonary complications of cytosine-arabioside therapy: radiographic findings. <i>American Journal of Roentgenology</i> , 1987 , 149, 23-7	5.4	32
59	B-cell colony growth of malignant and normal B-cells. <i>British Journal of Cancer</i> , 1987 , 55, 397-405	8.7	5
58	Preliminary results of consolidation therapy with high-dose cytosine arabinoside for patients with bad-risk or relapsed acute leukemia or lymphoblastic non-Hodgkin's lymphoma. <i>European Journal of Cancer & Clinical Oncology</i> , 1987 , 23, 401-5		1
57	Recent developments in the early diagnosis of mycosis fungoides and Sjöary's syndrome. <i>European Journal of Cancer & Clinical Oncology</i> , 1987 , 23, 1581-3		1
56	Primary cutaneous large cell lymphomas of follicular center cell origin. A clinical follow-up study of nineteen patients. <i>Journal of the American Academy of Dermatology</i> , 1987 , 16, 518-26	4.5	152
55	Human B cell colony assays. <i>Blut</i> , 1987 , 55, 189-98		3
54	Percentage of S-phase cells in bone marrow aspirates, biopsy specimens and bone marrow aspirates corrected for blood dilution from patients with acute leukemia. <i>Leukemia Research</i> , 1987 , 11, 209-13	2.7	11
53	Hairy cell leukemia: in-vitro proliferation and pseudo-colony formation. <i>Leukemia Research</i> , 1987 , 11, 911-21	2.7	3

52	Angiotropic (intravascular) large cell lymphoma of the skin previously classified as malignant angioendotheliomatosis. <i>British Journal of Dermatology</i> , 1987 , 116, 393-9	4	48
51	Side effects of intermediate- and high-dose cytosine arabinoside in the treatment of refractory or relapsed acute leukaemia and non-Hodgkin's lymphoma. <i>Netherlands Journal of Medicine</i> , 1987 , 30, 64-74	0.5	26
50	Diffuse large cell lymphomas of follicular center cell origin presenting in the skin. A clinicopathologic and immunologic study of 16 patients. <i>American Journal of Pathology</i> , 1987 , 126, 325-33	5.8	70
49	A histologic study of lymph nodes from patients with the Sézary syndrome. <i>Cancer</i> , 1986 , 57, 2375-80	6.4	35
48	A randomized comparison of maintenance treatment with androgens, immunotherapy, and chemotherapy in adult acute myelogenous leukemia. A Leukemia-Lymphoma Group Trial of the EORTC. <i>Cancer</i> , 1986 , 58, 617-23	6.4	36
47	Peripheral T-cell lymphomas: a clinicopathological and immunological study of 10 cases. <i>Histopathology</i> , 1986 , 10, 235-49	7.3	34
46	Neurological complications in patients with cutaneous T-cell lymphoma. <i>British Journal of Dermatology</i> , 1986 , 114, 359-66	4	13
45	Histopathologic studies in Sézary syndrome and erythrodermic mycosis fungoides: a comparison with benign forms of erythroderma. <i>Journal of the American Academy of Dermatology</i> , 1986 , 15, 1217-26	4.5	80
44	Effect of the interval between high dose 1-beta-D-arabinofuranosylcytosine injections on leukemic cell load, intestinal toxicity, and normal hematopoietic stem cells in a rat model for acute myelogenous leukemia. <i>Cancer Research</i> , 1986 , 46, 3825-7	10.1	8
43	Reduction and repopulation of T-lymphocytes after cytoreductive therapy with or without autologous bone marrow rescue. <i>Experimental Hematology</i> , 1986 , 14, 173-7	3.1	8
42	Thrombocytopenic purpura, coeliac disease and IgA deficiency. <i>Netherlands Journal of Medicine</i> , 1986 , 29, 165-6	0.5	6
41	Quantitative electron microscopy in the early diagnosis of cutaneous T-cell lymphomas. A long-term follow-up study of 77 patients. <i>American Journal of Pathology</i> , 1986 , 123, 166-73	5.8	13
40	Intermediate and high-dose ARA-C and m-AMSA (or daunorubicin) as remission and consolidation treatment for patients with relapsed acute leukaemia and lymphoblastic non-Hodgkin lymphoma. <i>Scandinavian Journal of Haematology</i> , 1985 , 34, 83-7		29
39	A comparison of two culture techniques: an in vitro & an in vivo tumour colony-forming assay. <i>British Journal of Cancer</i> , 1985 , 52, 713-7	8.7	2
38	Mycosis fungoides simulating acanthosis nigricans. <i>American Journal of Dermatopathology</i> , 1985 , 7, 367-71	19	24
37	Analysis of major histocompatibility antigens and the mononuclear cell infiltrate in halo nevi. <i>Journal of Investigative Dermatology</i> , 1985 , 85, 25-9	4.3	35
36	Intercellular distribution of HLA-DR and T6 antigens in the epidermis of various dermatoses. <i>British Journal of Dermatology</i> , 1985 , 113, 802-4	4	9
35	Prediction of response of acute non-lymphocytic leukaemia to therapy with 'high dose' cytosine arabinoside. <i>British Journal of Haematology</i> , 1985 , 59, 733-4	4.5	4

34	Palmar-plantar skin changes and cytarabine. <i>Annals of Internal Medicine</i> , 1985 , 103, 805	8	18
33	An in vivo clonogenic assay for human tumors using plasma clot diffusion chambers implanted in mice. <i>European Journal of Cancer & Clinical Oncology</i> , 1985 , 21, 127-34		1
32	Clinical and histologic differentiation between lymphomatoid papulosis and pityriasis lichenoides. <i>Journal of the American Academy of Dermatology</i> , 1985 , 13, 418-28	4.5	51
31	Amsacrine with high-dose cytarabine in acute leukemia. <i>Cancer Treatment Reports</i> , 1985 , 69, 1447-8		23
30	Monoclonal antibodies in the histopathological diagnosis of non-Hodgkin lymphomas. <i>Netherlands Journal of Medicine</i> , 1985 , 28, 138-41	0.5	2
29	Lymphomatoid papulosis. <i>Dermatologic Clinics</i> , 1985 , 3, 735-47	4.2	3
28	Analysis of helper activity on pokeweed mitogen- and interleukin 2-driven immunoglobulin synthesis by neoplastic T4+ cells. <i>Journal of Clinical Investigation</i> , 1985 , 76, 2139-43	15.9	9
27	Immunohistochemical studies using monoclonal antibodies on lymph nodes from patients with mycosis fungoides and Szary's syndrome. <i>American Journal of Pathology</i> , 1985 , 120, 46-54	5.8	19
26	Lymphocytic infiltration of the skin (Jessner): a T-cell lymphoproliferative disease. <i>British Journal of Dermatology</i> , 1984 , 110, 523-9	4	44
25	Regulatory activity of neoplastic T cells in Sezary syndrome on in vitro immunoglobulin production. <i>Leukemia Research</i> , 1984 , 8, 873-84	2.7	10
24	Histiocytic sarcoma (true histiocytic lymphoma): a clinicopathological study of 20 cases. <i>Histopathology</i> , 1984 , 8, 105-23	7.3	31
23	Immunohistochemical studies in lymphocytic infiltration of the skin (Jessner) and discoid lupus erythematosus. A comparative study. <i>Journal of the American Academy of Dermatology</i> , 1984 , 11, 832-40 ^{4.5}	4.5	40
22	Long-term survival in adolescent and adult acute lymphoblastic leukemia. <i>Cancer</i> , 1983 , 52, 30-4	6.4	9
21	Lymphomatoid papulosis and Hodgkin's disease: are they related?. <i>Archives of Dermatological Research</i> , 1983 , 275, 159-67	3.3	42
20	Primary plasma cell leukaemia. A case report and a review of the literature. <i>Acta Haematologica</i> , 1983 , 70, 122-9	2.7	30
19	Experience with intermediate and high dose cytosine arabinoside in refractory acute leukaemia. <i>Oncology Research and Treatment</i> , 1983 , 6, 200-4	2.8	14
18	Immunological, cytochemical and ultrastructural studies in lymphomatoid papulosis. <i>British Journal of Dermatology</i> , 1983 , 108, 381-94	4	54
17	The cell population of cutaneous B-cell lymphomas. <i>British Journal of Dermatology</i> , 1983 , 108, 395-409	4	34

16	Diagnostic criteria in Szary's syndrome: a multiparameter study of peripheral blood lymphocytes in 32 patients with erythroderma. <i>Journal of Investigative Dermatology</i> , 1983 , 81, 392-7	4.3	64
15	Characterization of T-cell subpopulations in skin and peripheral blood of patients with cutaneous T-cell lymphomas and benign inflammatory dermatoses. <i>Journal of Investigative Dermatology</i> , 1983 , 80, 60-6	4.3	103
14	E-rosette positive acute lymphoblastic leukaemia in adolescents and adults. <i>British Journal of Haematology</i> , 1983 , 55, 295-304	4.5	3
13	Experience with intermediate- and high-dose cytosine arabinoside in relapsed and refractory acute leukaemia. <i>Netherlands Journal of Medicine</i> , 1983 , 26, 215-9	0.5	14
12	High dose cytosine arabinoside in the management of refractory acute leukaemia. <i>Scandinavian Journal of Haematology</i> , 1982 , 29, 141-6		56
11	Treatment of refractory acute leukaemia with high dose cytosine arabinoside. <i>British Journal of Haematology</i> , 1982 , 51, 497-8	4.5	12
10	Reticulum cell sarcomas (large cell lymphomas) presenting in the skin. High frequency of true histiocytic lymphoma. <i>Cancer</i> , 1982 , 50, 1367-79	6.4	30
9	The clinical and histological spectrum of lymphomatoid papulosis. <i>British Journal of Dermatology</i> , 1982 , 107, 131-44	4	182
8	Results of a multicentre study of the treatment of acute lymphoblastic leukaemia in adolescents and adults (1978-1981). <i>Netherlands Journal of Medicine</i> , 1982 , 25, 303-7	0.5	3
7	Solitary plasmocytoma of the vagina. <i>Acta Haematologica</i> , 1981 , 66, 140-4	2.7	5
6	The in vivo diffusion chamber technique for bone marrow or blood cell culture. <i>Blut</i> , 1981 , 42, 59-67		2
5	Primary lymphoblastic non-Hodgkin's lymphoma of the skin. <i>British Journal of Dermatology</i> , 1981 , 104, 333-8	4	3
4	Diffuse cutaneous mastocytosis with multiple cutaneous mastocytomas. Report of a case with clinical, histopathological and ultrastructural aspects. <i>British Journal of Dermatology</i> , 1980 , 102, 601-7	4	30
3	Megakaryoblastic leukaemia (acute myelofibrosis): a report of three cases. <i>British Journal of Haematology</i> , 1979 , 42, 9-20	4.5	117
2	Marrow culture in diffusion chambers in rabbits: III. Effect of endotoxin and leukocyte products on cell production. <i>American Journal of Hematology</i> , 1979 , 7, 33-43	7.1	8
1	The morphology of dyserythropoiesis in a patient with acute erythroleukaemia associated with multiple myeloma. <i>Scandinavian Journal of Haematology</i> , 1976 , 17, 353-68		7