

Torsten Lauirtzen

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4587932/torsten-lauirtzen-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

80
papers

2,901
citations

26
h-index

53
g-index

82
ext. papers

3,532
ext. citations

6.8
avg, IF

4.48
L-index

#	Paper	IF	Citations
80	Changes in type 2 diabetes incidence and mortality associated with introduction of HbA1c as diagnostic option: A Danish 24-year population-based study.. <i>Lancet Regional Health - Europe, The</i> , 2022 , 14, 100291		1
79	Role of fasting duration and weekday in incretin and glucose regulation. <i>Endocrine Connections</i> , 2021 , 10, X2-X3	3.5	78
78	Lung function in adults and future burden of obstructive lung diseases in a long-term follow-up. <i>Npj Primary Care Respiratory Medicine</i> , 2020 , 30, 10	3.2	2
77	Alanine Aminotransferase and 20-Year Risk of Major Chronic Diseases and Death in a Healthy Cohort Aged 30 to 49 Years. <i>Clinical Epidemiology</i> , 2020 , 12, 345-351	5.9	3
76	Role of fasting duration and weekday in incretin and glucose regulation. <i>Endocrine Connections</i> , 2020 , 9, 279-288	3.5	2
75	The effect of training GPs in motivational interviewing on incident cardiovascular disease and mortality in people with screen-detected diabetes. Results from the ADDITION-Denmark randomised trial. <i>BJGP Open</i> , 2020 , 4,	3.1	1
74	Abdominal visceral and subcutaneous adipose tissue and associations with cardiometabolic risk in Inuit, Africans and Europeans: a cross-sectional study. <i>BMJ Open</i> , 2020 , 10, e038071	3	8
73	Greater glucagon-like peptide-1 responses to oral glucose are associated with lower central and peripheral blood pressures. <i>Cardiovascular Diabetology</i> , 2019 , 18, 130	8.7	0
72	Validity of Danish register diagnoses of myocardial infarction and stroke against experts in people with screen-detected diabetes. <i>BMC Public Health</i> , 2019 , 19, 228	4.1	4
71	A Randomised Trial Examining Cardiovascular Morbidity and All-Cause Mortality 24 years Following General Health Checks: the Ebeltoft Health Promotion Project (EHPP). <i>BMJ Open</i> , 2019 , 9, e030400	3	1
70	Habitual physical activity is associated with lower fasting and greater glucose-induced GLP-1 response in men. <i>Endocrine Connections</i> , 2019 , 8, 1607-1617	3.5	1
69	Long-term effects of intensive multifactorial therapy in individuals with screen-detected type 2 diabetes in primary care: 10-year follow-up of the ADDITION-Europe cluster-randomised trial. <i>Lancet Diabetes and Endocrinology,the</i> , 2019 , 7, 925-937	18.1	18
68	Impact of Geriatrician-Performed Comprehensive Geriatric Care on Medication Use and Cognitive Function in Older Adults Referred to a Non-Hospital-Based Rehabilitation Unit. <i>American Journal of Medicine</i> , 2019 , 132, 93-102.e2	2.4	1
67	Risk Factors for Incident Diabetic Polyneuropathy in a Cohort With Screen-Detected Type 2 Diabetes Followed for 13 Years: ADDITION-Denmark. <i>Diabetes Care</i> , 2018 , 41, 1068-1075	14.6	96
66	Geriatrician-performed comprehensive geriatric care in older adults referred to an outpatient community rehabilitation unit: A randomized controlled trial. <i>European Journal of Internal Medicine</i> , 2018 , 51, 18-24	3.9	6
65	Evidence of a liver-alpha cell axis in humans: hepatic insulin resistance attenuates relationship between fasting plasma glucagon and glucagonotropic amino acids. <i>Diabetologia</i> , 2018 , 61, 671-680	10.3	41
64	Early-onset type 2 diabetes: Age gradient in clinical and behavioural risk factors in 5115 persons with newly diagnosed type 2 diabetes-Results from the DD2 study. <i>Diabetes/Metabolism Research and Reviews</i> , 2018 , 34, e2968	7.5	20

63	Risk-Factor Trajectories Preceding Diabetic Polyneuropathy: ADDITION-Denmark. <i>Diabetes Care</i> , 2018 , 41, 1955-1962	14.6	20
62	Risk Factors for the Presence and Progression of Cardiovascular Autonomic Neuropathy in Type 2 Diabetes: ADDITION-Denmark. <i>Diabetes Care</i> , 2018 , 41, 2586-2594	14.6	40
61	Response to Comment on Andersen et al. Risk-Factor Trajectories Preceding Diabetic Polyneuropathy: ADDITION-Denmark. <i>Diabetes Care</i> 2018;41:1955-1962. <i>Diabetes Care</i> , 2018 , 41, e148-e149	14.6	149
60	The role of physical activity in the development of first cardiovascular disease event: a tree-structured survival analysis of the Danish ADDITION-PRO cohort. <i>Cardiovascular Diabetology</i> , 2018 , 17, 126	8.7	12
59	Short-term telemedical home blood pressure monitoring does not improve blood pressure in uncomplicated hypertensive patients. <i>Journal of Human Hypertension</i> , 2017 , 31, 93-98	2.6	7
58	Cardiovascular risk factors and incident albuminuria in screen-detected type 2 diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2017 , 33, e2877	7.5	2
57	Ethnic differences in anthropometric measures and abdominal fat distribution: a cross-sectional pooled study in Inuit, Africans and Europeans. <i>Journal of Epidemiology and Community Health</i> , 2017 , 71, 536-543	5.1	17
56	Does training of general practitioners for intensive treatment of people with screen-detected diabetes have a spillover effect on mortality and cardiovascular morbidity in high risk individuals with normoglycaemia? Results from the ADDITION-Denmark cluster-randomised controlled trial. <i>Diabetologia</i> , 2017 , 60, 1816-1821	10.3	5
55	A Low-Frequency Inactivating Variant Enriched in the Finnish Population Is Associated With Fasting Insulin Levels and Type 2 Diabetes Risk. <i>Diabetes</i> , 2017 , 66, 2019-2032	0.9	29
54	and Loci Identified through Large-Scale Exome Chip Analysis Regulate Kidney Development and Function. <i>Journal of the American Society of Nephrology: JASN</i> , 2017 , 28, 981-994	12.7	30
53	Exome-wide association study of plasma lipids in >300,000 individuals. <i>Nature Genetics</i> , 2017 , 49, 1758-1766	17.6	310
52	The obesity-associated risk of cardiovascular disease and all-cause mortality is not lower in Inuit compared to Europeans: A cohort study of Greenlandic Inuit, Nunavik Inuit and Danes. <i>Atherosclerosis</i> , 2017 , 265, 207-214	3.1	10
51	Effect of screening for type 2 diabetes on risk of cardiovascular disease and mortality: a controlled trial among 139,075 individuals diagnosed with diabetes in Denmark between 2001 and 2009. <i>Diabetologia</i> , 2017 , 60, 2192-2199	10.3	37
50	Effect of population screening for type 2 diabetes and cardiovascular risk factors on mortality rate and cardiovascular events: a controlled trial among 1,912,392 Danish adults. <i>Diabetologia</i> , 2017 , 60, 2183-2191	19.1	28
49	Alma-Ata to Berlin: diabetes prevention and treatment to achieve healthy living. <i>Diabetic Medicine</i> , 2017 , 34, 1169-1170	3.5	2
48	Sequence data and association statistics from 12,940 type 2 diabetes cases and controls. <i>Scientific Data</i> , 2017 , 4, 170179	8.2	22
47	Does Training and Support of General Practitioners in Intensive Treatment of People with Screen-Detected Diabetes Improve Medication, Morbidity and Mortality in People with Clinically-Diagnosed Diabetes? Investigation of a Spill-Over Effect in a Cluster RCT. <i>PLoS ONE</i> , 2017 , 12, e0170697	3.7	1
46	Methylglyoxal is associated with changes in kidney function among individuals with screen-detected Type 2 diabetes mellitus. <i>Diabetic Medicine</i> , 2016 , 33, 1625-1631	3.5	29

45	High Prevalence of Hypertension in a Danish Population Telemedical Home Measurement of Blood Pressure in Citizens Aged 55-64 Years in Holstebro County. <i>American Journal of Hypertension</i> , 2016 , 29, 439-47	2.3	9
44	Glucose-Dependent Insulinotropic Polypeptide Is Associated With Lower Low-Density Lipoprotein But Unhealthy Fat Distribution, Independent of Insulin: The ADDITION-PRO Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016 , 101, 485-93	5.6	34
43	A randomised trial of the effect and cost-effectiveness of early intensive multifactorial therapy on 5-year cardiovascular outcomes in individuals with screen-detected type 2 diabetes: the Anglo-Danish-Dutch Study of Intensive Treatment in People with Screen-Detected Diabetes in Primary Care (ADDITION-Europe) study. <i>Health Technology Assessment</i> , 2016 , 20, 1-86	4.4	11
42	Response to the Letter: Comment on "Abdominal Fat Distribution and Cardiovascular Risk in Men and Women With Different Levels of Glucose Tolerance" by Scheuer S.H., et al. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016 , 101, L13-4	5.6	
41	SP096TREATMENT OF HYPERTENSION USING TELEMEDICAL HOME BLOOD PRESSURE MEASUREMENTS. <i>Nephrology Dialysis Transplantation</i> , 2016 , 31, i117-i117	4.3	
40	The genetic architecture of type 2 diabetes. <i>Nature</i> , 2016 , 536, 41-47	50.4	704
39	Functional and genetic epidemiological characterisation of the FFAR4 (GPR120) p.R270H variant in the Danish population. <i>Journal of Medical Genetics</i> , 2016 , 53, 616-23	5.8	14
38	Incidence of register-based diabetes 10 years after a stepwise diabetes screening programme: the ADDITION-Denmark study. <i>Diabetologia</i> , 2016 , 59, 989-97	10.3	10
37	Insulin Resistance Is Accompanied by Increased Fasting Glucagon and Delayed Glucagon Suppression in Individuals With Normal and Impaired Glucose Regulation. <i>Diabetes</i> , 2016 , 65, 3473-3481	0.9	93
36	Association of self-perceived body image with body mass index and type 2 diabetes-The ADDITION-PRO study. <i>Preventive Medicine</i> , 2015 , 75, 64-9	4.3	8
35	Impact of intensive treatment on serum methylglyoxal levels among individuals with screen-detected type 2 diabetes: the ADDITION-Denmark study. <i>Acta Diabetologica</i> , 2015 , 52, 929-36	3.9	8
34	The role of serum methylglyoxal on diabetic peripheral and cardiovascular autonomic neuropathy: the ADDITION Denmark study. <i>Diabetic Medicine</i> , 2015 , 32, 778-85	3.5	30
33	Physical activity energy expenditure vs cardiorespiratory fitness level in impaired glucose metabolism. <i>Diabetologia</i> , 2015 , 58, 2709-17	10.3	10
32	Abdominal Fat Distribution and Cardiovascular Risk in Men and Women With Different Levels of Glucose Tolerance. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015 , 100, 3340-7	5.6	24
31	Cost-effectiveness of intensive multifactorial treatment compared with routine care for individuals with screen-detected Type 2 diabetes: analysis of the ADDITION-UK cluster-randomized controlled trial. <i>Diabetic Medicine</i> , 2015 , 32, 907-19	3.5	14
30	Intensive multifactorial treatment modifies the effect of family history of diabetes on glycaemic control in people with Type 2 diabetes: a post hoc analysis of the ADDITION-Denmark randomized controlled trial. <i>Diabetic Medicine</i> , 2015 , 32, 1085-9	3.5	3
29	FP090LESS THAN HALF OF CITIZENS, AGED 55-64 YEARS, HAD A NORMAL BLOOD PRESSURE IN A DANISH POPULATION PREVALENCE OF HYPERTENSION IN HOLSTEBRO COUNTY USING TELEMEDICALLY TRANSMITTED HOME BLOOD PRESSURE MEASUREMENT. <i>Nephrology Dialysis Transplantation</i> , 2015 , 30, iii96-iii97	4.3	
28	Impact of PTBP1 rs11085226 on glucose-stimulated insulin release in adult Danes. <i>BMC Medical Genetics</i> , 2015 , 16, 17	2.1	4

27	Early Detection and Treatment of Type 2 Diabetes Reduce Cardiovascular Morbidity and Mortality: A Simulation of the Results of the Anglo-Danish-Dutch Study of Intensive Treatment in People With Screen-Detected Diabetes in Primary Care (ADDITION-Europe). <i>Diabetes Care</i> , 2015 , 38, 1449-55	14.6	154
26	Associations between ultrasound measures of abdominal fat distribution and indices of glucose metabolism in a population at high risk of type 2 diabetes: the ADDITION-PRO study. <i>PLoS ONE</i> , 2015 , 10, e0123062	3.7	22
25	Identification of low-frequency and rare sequence variants associated with elevated or reduced risk of type 2 diabetes. <i>Nature Genetics</i> , 2014 , 46, 294-8	36.3	241
24	Diabetes-specific quality of life but not health status is independently associated with glycaemic control among patients with type 2 diabetes: a cross-sectional analysis of the ADDITION-Europe trial cohort. <i>Diabetes Research and Clinical Practice</i> , 2014 , 104, 281-7	7.4	35
23	Impact of moderate interval exercise versus supine rest on the pharmacokinetics and pharmacodynamic profiles of subcutaneously administered growth hormone in adult growth hormone deficient patients. <i>Growth Hormone and IGF Research</i> , 2014 , 24, 198-204	2	3
22	General health checks may work. <i>BMJ, The</i> , 2014 , 349, g4697	5.9	5
21	Periodontal status among patients with diabetes in Nuuk, Greenland. <i>International Journal of Circumpolar Health</i> , 2014 , 73, 26093	1.7	4
20	A combined analysis of 48 type 2 diabetes genetic risk variants shows no discriminative value to predict time to first prescription of a glucose lowering drug in Danish patients with screen detected type 2 diabetes. <i>PLoS ONE</i> , 2014 , 9, e104837	3.7	9
19	Short-term effects of NPH insulin, insulin detemir, and insulin glargine on the GH-IGF1-IGFBP axis in patients with type 1 diabetes. <i>European Journal of Endocrinology</i> , 2014 , 171, 471-9	6.5	8
18	Effect of early multifactorial therapy compared with routine care on microvascular outcomes at 5 years in people with screen-detected diabetes: a randomized controlled trial: the ADDITION-Europe Study. <i>Diabetes Care</i> , 2014 , 37, 2015-23	14.6	48
17	Socioeconomic position and cardiovascular risk factors among people with screen-detected Type 2 DM: six-year follow-up of the ADDITION-Denmark trial. <i>Primary Care Diabetes</i> , 2014 , 8, 322-9	2.4	8
16	All-cause mortality and pharmacological treatment intensity following a high risk screening program for diabetes. A 6.6 year follow-up of the ADDITION study, Denmark. <i>Primary Care Diabetes</i> , 2012 , 6, 193-200	2.4	13
15	HbA1c and cardiovascular risk score identify people who may benefit from preventive interventions: a 7 year follow-up of a high-risk screening programme for diabetes in primary care (ADDITION), Denmark. <i>Diabetologia</i> , 2011 , 54, 1318-26	10.3	27
14	Health tests and health consultations reduced cardiovascular risk without psychological strain, increased healthcare utilization or increased costs. An overview of the results from a 5-year randomized trial in primary care. The Ebeltoft Health Promotion Project (EHPP). <i>Scandinavian Journal of Public Health</i> , 2008 , 36, 650-61	3	31
13	Is prevention of Type-2 diabetes feasible and efficient in primary care? A systematic PubMed review. <i>Primary Care Diabetes</i> , 2007 , 1, 5-11	2.4	27
12	How do general practitioners, pharmacists and patients evaluate the substitution system for prescription in Denmark?. <i>European Journal of General Practice</i> , 2002 , 8, 13-17	2.8	2
11	The ADDITION study: proposed trial of the cost-effectiveness of an intensive multifactorial intervention on morbidity and mortality among people with Type 2 diabetes detected by screening. <i>International Journal of Obesity</i> , 2000 , 24 Suppl 3, S6-11	5.5	200
10	Near-Patient Test for C-Reactive Protein in General Practice: Assessment of Clinical, Organizational, and Economic Outcomes. <i>Clinical Chemistry</i> , 1999 , 45, 478-485	5.5	50

9	From science to everyday clinical practice. Need for systematic evaluation of research findings. <i>Scandinavian Journal of Primary Health Care</i> , 1999 , 17, 6-10	2.7	3
8	Audiometry in general practice: validation of a pragmatic pure-tone audiometry method. <i>Scandinavian Audiology</i> , 1998 , 27, 137-42		10
7	Do general practitioners want guidelines? Attitudes toward a county-based and a national college-based approach. <i>Scandinavian Journal of Primary Health Care</i> , 1997 , 15, 141-5	2.7	13
6	C-reactive protein in general practice--how commonly is it used and why?. <i>Scandinavian Journal of Primary Health Care</i> , 1997 , 15, 35-8	2.7	10
5	Evaluation of a near-patient test for C-reactive protein used in daily routine in primary healthcare by use of difference plots. <i>Clinical Chemistry</i> , 1997 , 43, 2064-2075	5.5	33
4	Validity of self-reported smoking habits. <i>Scandinavian Journal of Primary Health Care</i> , 1995 , 13, 236-7	2.7	19
3	Ebeltoft project: baseline data from a five-year randomized, controlled, prospective health promotion study in a Danish population. <i>British Journal of General Practice</i> , 1995 , 45, 542-7	1.6	34
2	The ADDITION study: proposed trial of the cost-effectiveness of an intensive multifactorial intervention on morbidity and mortality among people with Type 2 diabetes detected by screening. <i>International Journal of Obesity</i> , 24 , S6-S11	5.5	26
1	Review of prandial glucose regulation with repaglinide: a solution to the problem of hypoglycaemia in the treatment of Type 2 diabetes?. <i>International Journal of Obesity</i> , 24 , S21-S31	5.5	5