

# Davy Vancampfort

## List of Publications by Year in descending order

Source: [//exaly.com/author-pdf/4594280/publications.pdf](https://exaly.com/author-pdf/4594280/publications.pdf)

Version: 2025-02-01

420  
papers

30,029  
citations

3139

87  
h-index

4771

159  
g-index

453  
all docs

453  
docs citations

453  
times ranked

26924  
citing authors

#	ARTICLE	IF	CITATIONS
1	The feasibility of a stepping exergame prototype for older adults with major neurocognitive disorder residing in a long-term care facility: a mixed methods pilot study. <i>Disability and Rehabilitation</i> , 2024, 46, 896-910.	2.7	4
2	Physical activity, sedentary behaviour and mental health outcomes in firefighters: A cross-sectional study. <i>Journal of Workplace Behavioral Health</i> , 2024, 39, 88-105.	1.6	3
3	Dropout From Exercise Interventions in Adults With Fibromyalgia: A Systematic Review and Meta-analysis. <i>Archives of Physical Medicine and Rehabilitation</i> , 2024, 105, 571-579.	2.2	5
4	Levels of sedentary behaviour in people with fibromyalgia: a systematic review and meta-analysis. <i>Disability and Rehabilitation</i> , 2024, 46, 1921-1927.	2.7	3
5	The Effectiveness of Interventions for Improving Chronic Pain Symptoms Among People With Mental Illness: A Systematic Review. <i>Journal of Pain</i> , 2024, 25, 104421.	1.4	0
6	Dropout From Exercise Interventions in Adults With Knee or Hip Osteoarthritis: A Systematic Review and Meta-analysis. <i>Archives of Physical Medicine and Rehabilitation</i> , 2024, 105, 2317-2326.	2.2	4
7	Global and risk-group stratified well-being and mental health during the COVID-19 pandemic in adults: Results from the international COH-FIT Study. <i>Psychiatry Research</i> , 2024, 342, 115972.	3.4	0
8	The Daily Mile reduces depressive and anxiety symptoms in school-going Ugandan adolescents aged 16-17. <i>Pan African Medical Journal</i> , 2024, 48, .	0.7	0
9	Physical activity counselling in people with suicidal ideation: a secondary analysis of a pilot study in Ugandan primary care settings. <i>Pan African Medical Journal</i> , 2024, 48, .	0.7	0
10	The association of cannabis use with fast-food consumption, overweight, and obesity among adolescents aged 12-15 years from 28 countries. <i>Journal of Substance Use</i> , 2023, 28, 952-961.	1.1	1
11	Efficacy of basic body awareness therapy on functional outcomes: A systematic review and meta-analysis of randomized controlled trials. <i>Physiotherapy Research International</i> , 2023, 28, .	1.6	2
12	Physical activity correlates in people with fibromyalgia: a systematic review. <i>Disability and Rehabilitation</i> , 2023, 45, 4165-4174.	2.7	5
13	Autonomous Motivation for Exercise Is Key to an Active Lifestyle in Firefighters. <i>Workplace Health and Safety</i> , 2023, 71, 238-244.	1.8	2
14	Basic body awareness therapy within physiotherapy: More clarity about its' core concepts and more scientific evidence is needed. <i>Physiotherapy Research International</i> , 2023, 28, .	1.6	0
15	Efficacy of an 8-hour education intervention on dementia knowledge, attitude and skills in healthcare professionals in regional hospitals: a nation-wide study from Uganda. <i>Pan African Medical Journal</i> , 2023, 44, .	0.7	1
16	Exercise as medicine for depressive symptoms? A systematic review and meta-analysis with meta-regression. <i>British Journal of Sports Medicine</i> , 2023, 57, 1049-1057.	7.7	144
17	Correlates of sedentary behavior in people with fibromyalgia: A systematic review. <i>International Journal of Rheumatic Diseases</i> , 2023, 26, 841-849.	1.7	3
18	Sticking with it? Factors associated with exercise adherence in people with alcohol use disorder. <i>Addictive Behaviors</i> , 2023, 144, 107730.	3.2	7

#	ARTICLE	IF	CITATIONS
19	Physical activity levels among people with fibromyalgia: A systematic review and meta-analysis. <i>Musculoskeletal Care</i> , 2023, 21, 623-632.	1.6	3
20	Exergaming for people with major neurocognitive disorder: a qualitative study. <i>Disability and Rehabilitation</i> , 2022, 44, 2044-2052.	2.7	13
21	Exergames in people with major neurocognitive disorder: a systematic review. <i>Disability and Rehabilitation: Assistive Technology</i> , 2022, 17, 376-389.	3.2	22
22	Correlates of physical activity stages of change in people living with HIV in a Ugandan community. <i>Disability and Rehabilitation</i> , 2022, 44, 443-448.	2.7	4
23	Physical activity correlates in children and adolescents, adults, and older adults with an intellectual disability: a systematic review. <i>Disability and Rehabilitation</i> , 2022, 44, 4189-4200.	2.7	8
24	The collaborative outcomes study on health and functioning during infection times in adults (COH-FIT-Adults): Design and methods of an international online survey targeting physical and mental health effects of the COVID-19 pandemic. <i>Journal of Affective Disorders</i> , 2022, 299, 393-407.	4.7	24
25	Physical activity correlates in children and adolescents with autism spectrum disorder: a systematic review. <i>Disability and Rehabilitation</i> , 2022, 44, 6539-6550.	2.7	8
26	Can high-intensity interval training improve mental health outcomes in the general population and those with physical illnesses? A systematic review and meta-analysis. <i>British Journal of Sports Medicine</i> , 2022, 56, 279-291.	7.7	35
27	Physical and mental health impact of COVID-19 on children, adolescents, and their families: The Collaborative Outcomes study on Health and Functioning during Infection Times - Children and Adolescents (COH-FIT-C&A). <i>Journal of Affective Disorders</i> , 2022, 299, 367-376.	4.7	47
28	Prevalence of diabetes in people with intellectual disabilities and age- and gender-matched controls: A meta-analysis. <i>Journal of Applied Research in Intellectual Disabilities</i> , 2022, 35, 301-311.	2.2	7
29	Prospective Associations of Leisure-Time Physical Activity With Psychological Distress and Well-Being: A 12-Year Cohort Study. <i>Psychosomatic Medicine</i> , 2022, 84, 116-122.	2.3	8
30	A large-scale meta-analytic atlas of mental health problems prevalence during the COVID-19 early pandemic. <i>Journal of Medical Virology</i> , 2022, 94, 1935-1949.	3.8	182
31	Mental contrasting and implementation of physical activity intentions in Ugandan primary care patients with mental health problems: A real-world intervention involving support partners. <i>Psychiatry Research</i> , 2022, 307, 114335.	3.4	6
32	Osteoporosis referral and treatment among people with severe mental illness: A ten-year data linkage study. <i>Journal of Psychiatric Research</i> , 2022, 147, 94-102.	3.1	1
33	Prospective associations between multiple lifestyle behaviors and depressive symptoms. <i>Journal of Affective Disorders</i> , 2022, 301, 233-239.	4.7	15
34	Multimorbidity among Latinx-Americans and Asian American/Pacific Islanders. <i>Journal of Ageing and Longevity</i> , 2022, 2, 26-33.	1.0	2
35	Less time spent walking and depressive symptoms are associated with more self-perceived disability in primary care patients: a cross-sectional study from Uganda. <i>Pan African Medical Journal</i> , 2022, 41, .	0.7	0
36	Exercise to spot the differences: a framework for the effect of exercise on hippocampal pattern separation in humans. <i>Reviews in the Neurosciences</i> , 2022, 33, 555-582.	4.0	4

#	ARTICLE	IF	CITATIONS
37	Predictors of falls and fractures leading to hospitalisation in 36%101 people with affective disorders: a large representative cohort study. <i>BMJ Open</i> , 2022, 12, e055070.	2.0	4
38	Exercise and PTSD Symptoms in Emergency Service and Frontline Medical Workers: A Systematic Review. <i>Translational Journal of the American College of Sports Medicine</i> , 2022, 7, .	0.6	1
39	Mental health outcomes of the Daily Mile in elementary school children: a single-arm pilot study. <i>Child and Adolescent Mental Health</i> , 2022, 27, 361-368.	4.9	5
40	Mental health and lifestyle in mental health nurses: a cross-sectional, nation-wide study from Uganda during COVID-19 times. <i>Pan African Medical Journal</i> , 2022, 42, .	0.7	4
41	Precision-based exercise in people with anxiety and stress related disorders: Are there interindividual differences in anxiolytic effects? An ancillary meta-analysis of randomized controlled trials. <i>Psychiatry Research</i> , 2022, 317, 114803.	3.4	4
42	Associations between mild depression and anxiety and physical inactivity among school-going Ugandan adolescents aged 14 to 17 years. <i>Journal of Child and Adolescent Mental Health</i> , 2022, 34, 115-125.	0.1	2
43	Functioning in schizophrenia: a Delphi study covering the perspective of physiotherapists. <i>Disability and Rehabilitation</i> , 2021, 43, 3733-3740.	2.7	9
44	The efficacy of meditation-based mind-body interventions for mental disorders: A meta-review of 17 meta-analyses of randomized controlled trials. <i>Journal of Psychiatric Research</i> , 2021, 134, 181-191.	3.1	55
45	The efficacy of a lay health workers led physical activity counselling program in patients with HIV and mental health problems: a real-world intervention from Uganda. <i>AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV</i> , 2021, 33, 1189-1195.	1.5	14
46	Food Insecurity With Hunger and Sexual Behavior Among Adolescents From 53 Countries. <i>International Journal of Sexual Health</i> , 2021, 33, 88-98.	1.7	11
47	Test-retest reliability, concurrent validity and correlates of the two-minute walk test in outpatients with alcohol use disorder. <i>Alcohol</i> , 2021, 90, 74-79.	0.9	4
48	Test-retest reliability and correlates of the 6-min walk test in people with alcohol use disorders. <i>Physiotherapy Research International</i> , 2021, 26, .	1.6	0
49	The efficacy of exergaming in people with major neurocognitive disorder residing in long-term care facilities: a pilot randomized controlled trial. <i>Alzheimer's Research and Therapy</i> , 2021, 13, .	6.7	39
50	Physical activity, physical fitness and quality of life in outpatients with a psychotic disorder versus healthy matched controls in a low-income country. <i>Schizophrenia Research</i> , 2021, 229, 1-2.	2.4	3
51	Dropout from exercise randomized controlled trials among people with anxiety and stress-related disorders: A meta-analysis and meta-regression. <i>Journal of Affective Disorders</i> , 2021, 282, 996-1004.	4.7	40
52	Correlates of missed HIV appointments in low-resource settings: a study from Uganda. <i>African Journal of AIDS Research</i> , 2021, 20, 125-131.	0.9	0
53	Effects of acute exercise on craving, mood and anxiety in non-treatment seeking adults with alcohol use disorder: An exploratory study. <i>Drug and Alcohol Dependence</i> , 2021, 220, 108506.	3.1	21
54	Reliability and validity of physical fitness tests in people with mental disorders: A systematic review and meta-analysis. <i>Physiotherapy Research International</i> , 2021, 26, .	1.6	4

#	ARTICLE	IF	CITATIONS
55	A Mental Healthâ€œInformed Physical Activity Intervention for First Responders and Their Partners Delivered Using Facebook: Mixed Methods Pilot Study. JMIR Formative Research, 2021, 5, e23432.	2.3	16
56	The Relationship between Motor Skills and Intelligence in Children with Autism Spectrum Disorder. Journal of Autism and Developmental Disorders, 2021, 52, 1189-1199.	2.3	14
57	Moderate to vigorous physical activity and sedentary behavior changes in self-isolating adults during the COVID-19 pandemic in Brazil: a cross-sectional survey exploring correlates. Sport Sciences for Health, 2021, 18, 155-163.	1.1	45
58	The anxiolytic effects of exercise for people with anxiety and related disorders: An update of the available meta-analytic evidence. Psychiatry Research, 2021, 302, 114046.	3.4	60
59	Influence of weekday of admission and level of distress on length of hospital stay in patients with low back pain: a retrospective cohort study. BMC Musculoskeletal Disorders, 2021, 22, .	2.2	1
60	Cross-sectional associations of leisure and transport related physical activity with depression and anxiety. Journal of Psychiatric Research, 2021, 140, 228-234.	3.1	24
61	Multimorbidity and obesity in older adults from six low- and middle-income countries. Preventive Medicine, 2021, 153, 106816.	2.9	19
62	Endothelial dysfunction in people with depressive disorders: A systematic review and meta-analysis. Journal of Psychiatric Research, 2021, 141, 152-159.	3.1	21
63	Risk of hospitalised falls and hip fractures in working age adults receiving mental health care. General Hospital Psychiatry, 2021, 72, 81-87.	2.5	8
64	Evaluation of the Developmental Coordination Questionnaire (DCDQ) as a Screening Instrument for Co-occurring Motor Problems in Children with Autism Spectrum Disorder. Journal of Autism and Developmental Disorders, 2021, 52, 4079-4088.	2.3	14
65	Changes in craving following acute aerobic exercise in adults with alcohol use disorder. Journal of Psychiatric Research, 2021, 142, 243-249.	3.1	14
66	The complex and dynamic interplay between self-esteem, belongingness and physical activity in daily life: An experience sampling study in adolescence and young adulthood. Mental Health and Physical Activity, 2021, 21, 100413.	2.2	20
67	Objectively measured physical activity and symptoms of psychopathology in general population adolescents from the SIGMA cohort. Mental Health and Physical Activity, 2021, 21, 100416.	2.2	4
68	Association between physical activity and leisure-time sedentary behavior among 140,808 adolescents aged 12 to 15 from 47 low- and middle-income countries. Public Health, 2021, 199, 1-9.	2.7	6
69	Physical activity, physical fitness and quality of life in outpatients with major depressive disorder versus matched healthy controls: Data from a low-income country. Journal of Affective Disorders, 2021, 294, 802-804.	4.7	6
70	Increased rates of respiratory disease in schizophrenia: A systematic review and meta-analysis including 619,214 individuals with schizophrenia and 52,159,551 controls. Schizophrenia Research, 2021, 237, 131-140.	2.4	41
71	Does being overweight play a role in the reduced inhibitory control of patients receiving treatment for substance use disorder?. Physiology and Behavior, 2021, 241, 113587.	2.4	1
72	Physical activity, exercise, and mental disorders: it is time to move on. Trends in Psychiatry and Psychotherapy, 2021, , .	0.7	56

#	ARTICLE	IF	CITATIONS
73	Physical Activity, Sedentary Behavior, and Cardiorespiratory Fitness in Hazardous and Non-Hazardous Alcohol Consumers. <i>American Journal of Health Promotion</i> , 2021, 35, 669-678.	2.6	17
74	The Impact of Pharmacological and Non-Pharmacological Interventions to Improve Physical Health Outcomes in People With Schizophrenia: A Meta-Review of Meta-Analyses of Randomized Controlled Trials. <i>Focus (American Psychiatric Publishing)</i> , 2021, 19, 116-128.	1.3	10
75	Independent and combined associations of sugar-sweetened beverage consumption, TV viewing, and physical activity with severe depressive symptoms among 59,402 adults. <i>Revista Brasileira De Psiquiatria</i> , 2021, 43, 574-583.	1.1	10
76	Exercise for the Treatment of Depression. <i>The Open Complementary Medicine Journal</i> , 2021, 1, 78-83.	2.8	191
77	Internalised HIV-related stigma associated with physical inactivity in people with HIV and AIDS: A cross-sectional study from Uganda. <i>African Journal of AIDS Research</i> , 2021, 20, 238-243.	0.9	5
78	Moderate to severe generalized anxiety disorder symptoms are associated with physical inactivity in people with HIV/AIDS: a study from Uganda. <i>International Journal of STD and AIDS</i> , 2021, 32, 170-175.	1.3	11
79	The VITAAL Stepping Exergame Prototype for Older Adults With Major Neurocognitive Disorder: A Usability Study. <i>Frontiers in Aging Neuroscience</i> , 2021, 13, .	4.1	10
80	Psychotic experiences and subjective cognitive complaints among 224 842 people in 48 low- and middle-income countries. <i>Epidemiology and Psychiatric Sciences</i> , 2020, 29, .	4.4	24
81	Cross-sectional and prospective relationships of passive and mentally active sedentary behaviours and physical activity with depression – CORRIGENDUM. <i>British Journal of Psychiatry</i> , 2020, 217, 459-459.	2.3	8
82	From impact factors to real impact: translating evidence on lifestyle interventions into routine mental health care. <i>Translational Behavioral Medicine</i> , 2020, 10, 1070-1073.	2.3	43
83	Cross-sectional and prospective relationships of passive and mentally active sedentary behaviours and physical activity with depression. <i>British Journal of Psychiatry</i> , 2020, 217, 413-419.	2.3	77
84	Sexual behavior and suicide attempts among adolescents aged 12–15 years from 38 countries: A global perspective. <i>Psychiatry Research</i> , 2020, 287, 112564.	3.4	41
85	Functional exercise capacity in inpatients with alcohol use disorder versus healthy controls: A pilot study. <i>Alcohol</i> , 2020, 82, 47-52.	0.9	9
86	Association of bullying victimization with overweight and obesity among adolescents from 41 low- and middle-income countries. <i>Pediatric Obesity</i> , 2020, 15, .	2.7	34
87	The efficacy of physical activity counseling in Ugandan patients with HIV and a co-morbid mental disorder: a pilot study. <i>AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV</i> , 2020, 32, 758-761.	1.5	6
88	Sedentary behavior and depression among community-dwelling adults aged ≥50 years: Results from the Irish Longitudinal Study on Ageing. <i>Journal of Affective Disorders</i> , 2020, 262, 389-396.	4.7	33
89	Does physical activity reduce the risk of psychosis? A systematic review and meta-analysis of prospective studies. <i>Psychiatry Research</i> , 2020, 284, 112675.	3.4	31
90	Handgrip strength is associated with suicidal thoughts in men: Cross-sectional analyses from NHANES. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2020, 30, 92-99.	3.4	25

#	ARTICLE	IF	CITATIONS
91	Prospective relationships of mentally passive sedentary behaviors with depression: Mediation by sleep problems. <i>Journal of Affective Disorders</i> , 2020, 265, 538-544.	4.7	27
92	Perceived barriers, benefits and correlates of physical activity in outpatients with Major Depressive Disorder: A study from Brazil. <i>Psychiatry Research</i> , 2020, 284, 112751.	3.4	11
93	Test-retest reliability, validity, and correlates of the 2-min walk test in outpatients with depression. <i>Physiotherapy Research International</i> , 2020, 25, .	1.6	8
94	Handgrip Strength Is Associated With Hippocampal Volume and White Matter Hyperintensities in Major Depression and Healthy Controls: A UK Biobank Study. <i>Psychosomatic Medicine</i> , 2020, 82, 39-46.	2.3	48
95	Metabolic syndrome and its components in people with intellectual disability: a meta-analysis. <i>Journal of Intellectual Disability Research</i> , 2020, 64, 804-815.	2.1	15
96	Associations of moderate to vigorous physical activity and sedentary behavior with depressive and anxiety symptoms in self-isolating people during the COVID-19 pandemic: A cross-sectional survey in Brazil. <i>Psychiatry Research</i> , 2020, 292, 113339.	3.4	189
97	A meta-review of "lifestyle psychiatry": the role of exercise, smoking, diet and sleep in the prevention and treatment of mental disorders. <i>World Psychiatry</i> , 2020, 19, 360-380.	12.0	575
98	Impact of Psychotropic Medication Effects on Obesity and the Metabolic Syndrome in People With Serious Mental Illness. <i>Frontiers in Endocrinology</i> , 2020, 11, .	4.0	101
99	Associations of interruptions to leisure-time sedentary behaviour with symptoms of depression and anxiety. <i>Translational Psychiatry</i> , 2020, 10, .	5.7	43
100	The Association Between Sedentary Behavior and Sarcopenia Among Adults Aged ≥65 Years in Low- and Middle-Income Countries. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 1708.	3.1	53
101	Associations of sedentary behavior in leisure and occupational contexts with symptoms of depression and anxiety. <i>Preventive Medicine</i> , 2020, 133, 106021.	2.9	42
102	The Impact of Pharmacologic and Nonpharmacologic Interventions to Improve Physical Health Outcomes in People With Dementia: A Meta-Review of Meta-Analyses of Randomized Controlled Trials. <i>Journal of the American Medical Directors Association</i> , 2020, 21, 1410-1414.e2.	2.7	20
103	Informal Caregiving, Chronic Physical Conditions, and Physical Multimorbidity in 48 Low- and Middle-Income Countries. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2020, 75, 1572-1578.	3.6	23
104	Sex difference in the association between physical activity and suicide attempts among adolescents from 48 countries: A global perspective. <i>Journal of Affective Disorders</i> , 2020, 266, 311-318.	4.7	28
105	Joint association of ultra-processed food and sedentary behavior with anxiety-induced sleep disturbance among Brazilian adolescents. <i>Journal of Affective Disorders</i> , 2020, 266, 135-142.	4.7	30
106	Informal caregiving and physical activity among 204,315 adults in 38 low- and middle-income countries: A cross-sectional study. <i>Preventive Medicine</i> , 2020, 132, 106007.	2.9	11
107	Risk of Hospitalized Falls and Hip Fractures in 22,103 Older Adults Receiving Mental Health Care vs 161,603 Controls: A Large Cohort Study. <i>Journal of the American Medical Directors Association</i> , 2020, 21, 1893-1899.	2.7	20
108	Predictors of physical activity recording in routine mental healthcare. <i>Mental Health and Physical Activity</i> , 2020, 18, 100329.	2.2	2

#	ARTICLE	IF	CITATIONS
109	Lifestyle behaviors among 4,343 Brazilian adults with severe mental illness and 55,859 general population controls: data from the Brazilian National Health Survey. <i>Revista Brasileira De Psiquiatria</i> , 2020, 42, 245-249.	1.1	7
110	Physical activity participation is associated with higher quality of life scores in men with alcohol use disorders: a study from Uganda. <i>African Health Sciences</i> , 2020, 20, 1407-1415.	0.8	5
111	Test-retest reliability and correlates of the Simple Physical Activity Questionnaire in Ugandan out-patients with psychosis. <i>African Health Sciences</i> , 2020, 20, 1438-1445.	0.8	6
112	Physical fitness and physical activity levels in people with alcohol use disorder versus matched healthy controls: A pilot study. <i>Alcohol</i> , 2019, 76, 73-79.	0.9	23
113	Cannabis use and leisure-time sedentary behavior among 94,035 adolescents aged 12-15 years from 24 low- and middle-income countries. <i>Addictive Behaviors</i> , 2019, 99, 106104.	3.2	3
114	The Lancet Psychiatry Commission: a blueprint for protecting physical health in people with mental illness. <i>Lancet Psychiatry</i> , 2019, 6, 675-712.	9.0	928
115	Physical activity and loneliness among adults aged 50 years or older in six low- and middle-income countries. <i>International Journal of Geriatric Psychiatry</i> , 2019, 34, 1855-1864.	2.4	27
116	Test-retest reliability, concurrent validity and correlates of the two-minute walk test in outpatients with psychosis. <i>Psychiatry Research</i> , 2019, 282, 112619.	3.4	6
117	Motives for physical activity in the adoption and maintenance of physical activity in middle-aged and old age outpatients with a mental disorder: A cross-sectional study from a low-income country. <i>Psychiatry Research</i> , 2019, 282, 112620.	3.4	6
118	A quantitative assessment of the views of mental health professionals on exercise for people with mental illness: perspectives from a low-resource setting. <i>African Health Sciences</i> , 2019, 19, 2172.	0.8	13
119	Cannabis use and physical activity among 89,777 adolescents aged 12-15 years from 21 low- and middle-income countries. <i>Drug and Alcohol Dependence</i> , 2019, 205, 107584.	3.1	10
120	The efficacy and safety of nutrient supplements in the treatment of mental disorders: a meta-analysis of randomized-controlled trials. <i>World Psychiatry</i> , 2019, 18, 308-324.	12.0	147
121	The experience of mental health service users in health system strengthening: lessons from Uganda. <i>International Journal of Mental Health Systems</i> , 2019, 13, .	3.2	12
122	Functional exercise capacity is associated with global functioning in patients with alcohol use disorder. <i>Archives of Psychiatric Nursing</i> , 2019, 33, 144-148.	2.0	5
123	Physical fitness levels and moderators in people with epilepsy: A systematic review and meta-analysis. <i>Epilepsy and Behavior</i> , 2019, 99, 106448.	2.0	6
124	Which clinical and biochemical predictors should be used to screen for diabetes in patients with serious mental illness receiving antipsychotic medication? A large observational study. <i>PLoS ONE</i> , 2019, 14, e0210674.	2.5	3
125	Correlates of physical activity among 142,118 adolescents aged 12-15 years from 48 low- and middle-income countries. <i>Preventive Medicine</i> , 2019, 127, 105819.	2.9	27
126	Associations between physical inactivity, major depressive disorder, and alcohol use disorder in people living with HIV in a Ugandan fishing community. <i>International Journal of STD and AIDS</i> , 2019, 30, 1177-1184.	1.3	6

#	ARTICLE	IF	CITATIONS
127	Physical activity and sedentary levels among people living with epilepsy: A systematic review and meta-analysis. <i>Epilepsy and Behavior</i> , 2019, 99, 106390.	2.0	10
128	Food insecurity (hunger) and suicide attempts among 179,771 adolescents attending school from 9 high-income, 31 middle-income, and 4 low-income countries: A cross-sectional study. <i>Journal of Affective Disorders</i> , 2019, 248, 91-98.	4.7	76
129	Associations between handgrip strength and mild cognitive impairment in middle-aged and older adults in six low- and middle-income countries. <i>International Journal of Geriatric Psychiatry</i> , 2019, 34, 609-616.	2.4	38
130	Physical activity protects from incident anxiety: A meta-analysis of prospective cohort studies. <i>Depression and Anxiety</i> , 2019, 36, 846-858.	3.9	257
131	The "online brain" how the Internet may be changing our cognition. <i>World Psychiatry</i> , 2019, 18, 119-129.	12.0	317
132	Associations of the Built Environment With Physical Activity and Sedentary Time in Ugandan Outpatients With Mental Health Problems. <i>Journal of Physical Activity and Health</i> , 2019, 16, 243-250.	3.0	12
133	Association of leisure-time sedentary behavior with fast food and carbonated soft drink consumption among 133,555 adolescents aged 12-15 years in 44 low- and middle-income countries. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2019, 16, .	4.5	39
134	Visual impairment and objectively measured physical activity and sedentary behaviour in US adolescents and adults: a cross-sectional study. <i>BMJ Open</i> , 2019, 9, e027267.	2.0	31
135	Promoting physical activity in vulnerable adults "at risk" of homelessness: a randomised controlled trial protocol. <i>BMJ Open</i> , 2019, 9, e026466.	2.0	4
136	Sedentary behavior and anxiety-induced sleep disturbance among 181,093 adolescents from 67 countries: a global perspective. <i>Sleep Medicine</i> , 2019, 58, 19-26.	1.4	35
137	Leisure-time sedentary behavior and suicide attempt among 126,392 adolescents in 43 countries. <i>Journal of Affective Disorders</i> , 2019, 250, 346-353.	4.7	40
138	Does loneliness contribute to mild cognitive impairment and dementia? A systematic review and meta-analysis of longitudinal studies. <i>Ageing Research Reviews</i> , 2019, 52, 7-16.	12.1	214
139	Bullying Victimization and Suicide Attempt Among Adolescents Aged 12-15 Years From 48 Countries. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2019, 58, 907-918.e4.	0.6	163
140	Leisure-time Sedentary Behavior and Obesity Among 116,762 Adolescents Aged 12-15 Years from 41 Low- and Middle-income Countries. <i>Obesity</i> , 2019, 27, 830-836.	4.3	28
141	Leisure-time sedentary behavior and loneliness among 148,045 adolescents aged 12-15 years from 52 low- and middle-income countries. <i>Journal of Affective Disorders</i> , 2019, 251, 149-155.	4.7	45
142	Food Insecurity Is Associated with Mild Cognitive Impairment among Middle-Aged and Older Adults in South Africa: Findings from a Nationally Representative Survey. <i>Nutrients</i> , 2019, 11, 749.	4.6	41
143	Health care professionals' perspectives on physical activity within the Ugandan mental health care system. <i>Mental Health and Physical Activity</i> , 2019, 16, 1-7.	2.2	22
144	Low Physical Activity and Cardiorespiratory Fitness in People With Schizophrenia: A Comparison With Matched Healthy Controls and Associations With Mental and Physical Health. <i>Frontiers in Psychiatry</i> , 2019, 10, .	2.7	45

#	ARTICLE	IF	CITATIONS
145	Mental health informed physical activity for first responders and their support partner: a protocol for a stepped-wedge evaluation of an online, codesigned intervention. <i>BMJ Open</i> , 2019, 9, e030668.	2.0	12
146	Correlates of leisure-time sedentary behavior among 181,793 adolescents aged 12-15 years from 66 low- and middle-income countries. <i>PLoS ONE</i> , 2019, 14, e0224339.	2.5	20
147	Yoga as part of a package of care versus non-standard care for schizophrenia. <i>The Cochrane Library</i> , 2019, 2019, .	2.4	6
148	The impact of pharmacological and non-pharmacological interventions to improve physical health outcomes in people with schizophrenia: a meta-review of meta-analyses of randomized controlled trials. <i>World Psychiatry</i> , 2019, 18, 53-66.	12.0	167
149	Do reductions in ghrelin contribute towards antipsychotic-induced weight gain?. <i>Schizophrenia Research</i> , 2019, 210, 301-302.	2.4	9
150	Cannabis use and suicide attempts among 86,254 adolescents aged 12-15 years from 21 low- and middle-income countries. <i>European Psychiatry</i> , 2019, 56, 8-13.	0.3	78
151	Correlates of sedentary behavior among community-dwelling adults with anxiety in six low- and middle-income countries. <i>Psychiatry Research</i> , 2019, 273, 501-508.	3.4	8
152	Test-retest reliability and clinical correlates of the Eurofit test battery in people with alcohol use disorders. <i>Psychiatry Research</i> , 2019, 271, 208-213.	3.4	13
153	Does exercise improve sleep quality in individuals with mental illness? A systematic review and meta-analysis. <i>Journal of Psychiatric Research</i> , 2019, 109, 96-106.	3.1	95
154	Prospective associations between physical activity and clinician diagnosed major depressive disorder in adults: A 13-year cohort study. <i>Preventive Medicine</i> , 2019, 118, 38-43.	2.9	18
155	Correlates of sedentary behavior in middle-aged and old age people with mild cognitive impairment: a multinational study. <i>International Psychogeriatrics</i> , 2019, 31, 579-589.	1.0	12
156	Handgrip strength, chronic physical conditions and physical multimorbidity in middle-aged and older adults in six low- and middle income countries. <i>European Journal of Internal Medicine</i> , 2019, 61, 96-102.	2.6	29
157	Handgrip strength and depression among 34,129 adults aged 50 years and older in six low- and middle-income countries. <i>Journal of Affective Disorders</i> , 2019, 243, 448-454.	4.7	65
158	Validity and correlates of the International Physical Activity Questionnaire in first-episode psychosis. <i>Microbial Biotechnology</i> , 2019, 13, 562-567.	2.2	12
159	Exercise as Medicine for Mental and Substance Use Disorders: A Meta-review of the Benefits for Neuropsychiatric and Cognitive Outcomes. <i>Sports Medicine</i> , 2019, 50, 151-170.	6.7	260
160	Barriers, attitudes, confidence, and knowledge of nurses regarding metabolic health screening and intervention in people with mental illness: a pilot study from Uganda. <i>African Health Sciences</i> , 2019, 19, 2546-2554.	0.8	9
161	Correlates of sedentary behavior in 2,375 people with depression from 6 low- and middle-income countries. <i>Journal of Affective Disorders</i> , 2018, 234, 97-104.	4.7	23
162	Leisure time physical activity and future psychological distress: A thirteen year longitudinal population-based study. <i>Journal of Psychiatric Research</i> , 2018, 101, 50-56.	3.1	31

#	ARTICLE	IF	CITATIONS
163	Grip Strength Is Associated With Cognitive Performance in Schizophrenia and the General Population: A UK Biobank Study of 476559 Participants. <i>Schizophrenia Bulletin</i> , 2018, 44, 728-736.	4.3	76
164	Physical Activity and Incident Depression: A Meta-Analysis of Prospective Cohort Studies. <i>American Journal of Psychiatry</i> , 2018, 175, 631-648.	10.5	1,012
165	Association Between Muscular Strength and Cognition in People With Major Depression or Bipolar Disorder and Healthy Controls. <i>JAMA Psychiatry</i> , 2018, 75, 740.	14.1	64
166	Associations between TV viewing and depressive symptoms among 60,202 Brazilian adults: The Brazilian national health survey. <i>Journal of Affective Disorders</i> , 2018, 236, 23-30.	4.7	30
167	Motives for physical activity in the adoption and maintenance of physical activity in men with alcohol use disorders. <i>Psychiatry Research</i> , 2018, 261, 522-526.	3.4	8
168	Physical activity correlates in people with mild cognitive impairment: findings from six low- and middle-income countries. <i>Public Health</i> , 2018, 156, 15-25.	2.7	11
169	Chronic Physical Conditions, Multimorbidity, and Mild Cognitive Impairment in Low- and Middle-Income Countries. <i>Journal of the American Geriatrics Society</i> , 2018, 66, 721-727.	2.8	89
170	Adherence to physical activity recommendations and physical and mental health risk in people with severe mental illness in Uganda. <i>Psychiatry Research</i> , 2018, 260, 236-240.	3.4	13
171	Relationship between sedentary behavior and depression: A mediation analysis of influential factors across the lifespan among 42,469 people in low- and middle-income countries. <i>Journal of Affective Disorders</i> , 2018, 229, 231-238.	4.7	110
172	Correlates of sedentary behaviour among adults with hazardous drinking habits in six low- and middle-income countries. <i>Psychiatry Research</i> , 2018, 261, 406-413.	3.4	8
173	Nutritional Deficiencies and Clinical Correlates in First-Episode Psychosis: A Systematic Review and Meta-analysis. <i>Schizophrenia Bulletin</i> , 2018, 44, 1275-1292.	4.3	62
174	Mild cognitive impairment and sedentary behavior: A multinational study. <i>Experimental Gerontology</i> , 2018, 108, 174-180.	3.8	25
175	Is autonomous motivation the key to maintaining an active lifestyle in first-episode psychosis?. <i>Microbial Biotechnology</i> , 2018, 12, 821-827.	2.2	29
176	Global physical activity levels among people living with HIV: a systematic review and meta-analysis. <i>Disability and Rehabilitation</i> , 2018, 40, 388-397.	2.7	100
177	Physical activity correlates in people living with HIV/AIDS: a systematic review of 45 studies. <i>Disability and Rehabilitation</i> , 2018, 40, 1618-1629.	2.7	67
178	Physical activity correlates among people with psychosis: Data from 47 low- and middle-income countries. <i>Schizophrenia Research</i> , 2018, 193, 412-417.	2.4	28
179	Multimorbidity and perceived stress: a population-based cross-sectional study among older adults across six low- and middle-income countries. <i>Maturitas</i> , 2018, 107, 84-91.	1.7	44
180	The Validity and Value of Self-reported Physical Activity and Accelerometry in People With Schizophrenia: A Population-Scale Study of the UK Biobank. <i>Schizophrenia Bulletin</i> , 2018, 44, 1293-1300.	4.3	67

#	ARTICLE	IF	CITATIONS
181	Sedentary behavior and anxiety: Association and influential factors among 42,469 community-dwelling adults in six low- and middle-income countries. <i>General Hospital Psychiatry</i> , 2018, 50, 26-32.	2.5	36
182	Correlates of physical activity among depressed older people in six low-income and middle-income countries: A community-based cross-sectional study. <i>International Journal of Geriatric Psychiatry</i> , 2018, 33, .	2.4	16
183	Physical activity and suicidal ideation: A systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2018, 225, 438-448.	4.7	159
184	Exercise self-efficacy correlates in people with psychosis. <i>Psychiatry Research</i> , 2018, 262, 359-362.	3.4	7
185	Correlates of low physical activity across 46 low- and middle-income countries: A cross-sectional analysis of community-based data. <i>Preventive Medicine</i> , 2018, 106, 107-113.	2.9	31
186	Gender differences in motives for physical activity across the stages of change in Ugandan outpatients with psychosis. <i>Schizophrenia Research</i> , 2018, 197, 568-569.	2.4	4
187	Effect of aerobic exercise on hippocampal volume in humans: A systematic review and meta-analysis. <i>NeuroImage</i> , 2018, 166, 230-238.	4.8	360
188	Correlates of physical activity among community-dwelling individuals aged 65 years or older with anxiety in six low- and middle-income countries. <i>International Psychogeriatrics</i> , 2018, 30, 705-714.	1.0	10
189	Corrigendum to "Physical activity and sedentary behavior in people with major depressive disorder: A systematic review and meta-analysis" [J. Affect. Disord. 210 (2017) 139-150]. <i>Journal of Affective Disorders</i> , 2018, 225, 79.	4.7	4
190	EPA guidance on physical activity as a treatment for severe mental illness: a meta-review of the evidence and Position Statement from the European Psychiatric Association (EPA), supported by the International Organization of Physical Therapists in Mental Health (IOPTMH). <i>European Psychiatry</i> , 2018, 54, 124-144.	0.3	425
191	Physical activity and mental health. <i>Lancet Psychiatry</i> , the, 2018, 5, 873.	9.0	45
192	Physical activity and generalized anxiety disorder: results from The Irish Longitudinal Study on Ageing (TILDA). <i>International Journal of Epidemiology</i> , 2018, 47, 1443-1453.	5.2	54
193	<i>Schizophrenia and Exercise.</i> , 2018, , 65-82.		0
194	Physical activity and HIV in sub-Saharan Africa: a systematic review of correlates and levels. <i>African Health Sciences</i> , 2018, 18, .	0.8	15
195	<i>Bipolar Disorder and Physical Activity.</i> , 2018, , 53-63.		1
196	<i>Research and Evaluation in Exercise and Mental Health.</i> , 2018, , 301-317.		0
197	Cardiorespiratory fitness and response to exercise treatment in depression. <i>BJPsych Open</i> , 2018, 4, 346-351.	1.0	31
198	Associations between active travel and physical multi-morbidity in six low- and middle-income countries among community-dwelling older adults: A cross-sectional study. <i>PLoS ONE</i> , 2018, 13, e0203277.	2.5	21

#	ARTICLE	IF	CITATIONS
199	Predictors of falls and fractures leading to hospitalization in people with schizophrenia spectrum disorder: A large representative cohort study. <i>Schizophrenia Research</i> , 2018, 201, 70-78.	2.4	28
200	Pain and severe sleep disturbance in the general population: Primary data and meta-analysis from 240,820 people across 45 low- and middle-income countries. <i>General Hospital Psychiatry</i> , 2018, 53, 52-58.	2.5	16
201	Association between depression and smoking: A global perspective from 48 low- and middle-income countries. <i>Journal of Psychiatric Research</i> , 2018, 103, 142-149.	3.1	55
202	Physical activity and sleep problems in 38 low- and middle-income countries. <i>Sleep Medicine</i> , 2018, 48, 140-147.	1.4	16
203	Sedentary behaviour and sleep problems among 42,489 community-dwelling adults in six low- and middle-income countries. <i>Journal of Sleep Research</i> , 2018, 27, .	4.2	37
204	Cultural responses towards the aftermath of suicide among the Acholi in Northern Uganda. <i>International Journal of Social Psychiatry</i> , 2018, 64, 545-553.	2.5	5
205	Sibship size, birth order and psychotic experiences: Evidence from 43 low- and middle-income countries. <i>Schizophrenia Research</i> , 2018, 201, 406-412.	2.4	1
206	Is it possible for people with severe mental illness to sit less and move more? A systematic review of interventions to increase physical activity or reduce sedentary behaviour. <i>Schizophrenia Research</i> , 2018, 202, 3-16.	2.4	60
207	Association Between Gait Speed With Mortality, Cardiovascular Disease and Cancer: A Systematic Review and Meta-analysis of Prospective Cohort Studies. <i>Journal of the American Medical Directors Association</i> , 2018, 19, 981-988.e7.	2.7	139
208	Moving to Beat Anxiety: Epidemiology and Therapeutic Issues with Physical Activity for Anxiety. <i>Current Psychiatry Reports</i> , 2018, 20, .	6.2	155
209	Hand grip strength and cognitive function among elderly cancer survivors. <i>PLoS ONE</i> , 2018, 13, e0197909.	2.5	34
210	The association of depression and all-cause and cause-specific mortality: an umbrella review of systematic reviews and meta-analyses. <i>BMC Medicine</i> , 2018, 16, .	7.4	165
211	Physiotherapy for people with mental health problems in Sub-Saharan African countries: a systematic review. <i>Archives of Physiotherapy</i> , 2018, 8, .	2.3	11
212	Predictors of Falls and Fractures Leading to Hospitalization in People With Dementia: A Representative Cohort Study. <i>Journal of the American Medical Directors Association</i> , 2018, 19, 607-612.	2.7	47
213	Correlates of sedentary behavior in the general population: A cross-sectional study using nationally representative data from six low- and middle-income countries. <i>PLoS ONE</i> , 2018, 13, e0202222.	2.5	57
214	Psychotic experiences as an independent risk factor for angina pectoris in 48 low- and middle-income countries. <i>World Psychiatry</i> , 2018, 17, 232-234.	12.0	6
215	Passive and mentally-active sedentary behaviors and incident major depressive disorder: A 13-year cohort study. <i>Journal of Affective Disorders</i> , 2018, 241, 579-585.	4.7	94
216	New steps for treating alcohol use disorder: the emerging importance of physical exercise. <i>Psychopharmacology</i> , 2018, 235, 2771-2773.	3.0	9

#	ARTICLE	IF	CITATIONS
217	Associations between TV viewing, sitting time, physical activity and insomnia among 100,839 Brazilian adolescents. <i>Psychiatry Research</i> , 2018, 269, 700-706.	3.4	26
218	Sedentary behavior and depressive symptoms among 67,077 adolescents aged 12-15 years from 30 low- and middle-income countries. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2018, 15, .	4.5	105
219	Sedentary behavior and perceived stress among adults aged ≥50 years in six low- and middle-income countries. <i>Maturitas</i> , 2018, 116, 100-107.	1.7	21
220	Depressive symptoms, handgrip strength, and weight status in US older adults. <i>Journal of Affective Disorders</i> , 2018, 238, 305-310.	4.7	39
221	Environmental risk factors and nonpharmacological and nonsurgical interventions for obesity: An umbrella review of meta-analyses of cohort studies and randomized controlled trials. <i>European Journal of Clinical Investigation</i> , 2018, 48, .	3.2	55
222	Different patterns of alcohol consumption and the incidence and persistence of depressive and anxiety symptoms among older adults in Ireland: A prospective community-based study. <i>Journal of Affective Disorders</i> , 2018, 238, 651-658.	4.7	15
223	Sedentary Behavior and Quality of Life in People with Psychotic Disorders from a Low Income Country: A Study from Uganda. <i>Community Mental Health Journal</i> , 2018, 55, 714-720.	1.3	1
224	Yoga as part of a package of care versus standard care for schizophrenia. <i>The Cochrane Library</i> , 2018, 2018, .	2.4	15
225	Exercise and Posttraumatic Stress Disorder. , 2018, , 375-387.		2
226	Evaluating correlations between physical activity, psychological mediators of physical activity, and negative symptoms in individuals living with psychosis and diabetes.. <i>Psychiatric Rehabilitation Journal</i> , 2018, 41, 153-156.	1.5	3
227	The intriguing relationship between coronary heart disease and mental disorders. <i>Dialogues in Clinical Neuroscience</i> , 2018, 20, 31-40.	5.6	326
228	Physical Activity Levels and Psychosis: A Mediation Analysis of Factors Influencing Physical Activity Target Achievement Among 204 186 People Across 46 Low- and Middle-Income Countries. <i>Schizophrenia Bulletin</i> , 2017, , sbw111.	4.3	53
229	Aerobic Exercise Improves Cognitive Functioning in People With Schizophrenia: A Systematic Review and Meta-Analysis. <i>Schizophrenia Bulletin</i> , 2017, , sbw115.	4.3	277
230	Chronic physical conditions, multimorbidity and physical activity across 46 low- and middle-income countries. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2017, 14, .	4.5	114
231	An examination of the anxiolytic effects of exercise for people with anxiety and stress-related disorders: A meta-analysis. <i>Psychiatry Research</i> , 2017, 249, 102-108.	3.4	418
232	Are people with schizophrenia adherent to diabetes medication? A comparative meta-analysis. <i>Psychiatry Research</i> , 2017, 250, 17-24.	3.4	21
233	Exercise as treatment for alcohol use disorders: systematic review and meta-analysis. <i>British Journal of Sports Medicine</i> , 2017, 51, 1058-1064.	7.7	126
234	Depression and physical health multimorbidity: primary data and country-wide meta-analysis of population data from 190 593 people across 43 low- and middle-income countries. <i>Psychological Medicine</i> , 2017, 47, 2107-2117.	4.6	113

#	ARTICLE	IF	CITATIONS
235	The effect of active video games on cognitive functioning in clinical and non-clinical populations: A meta-analysis of randomized controlled trials. <i>Neuroscience and Biobehavioral Reviews</i> , 2017, 78, 34-43.	7.3	278
236	Prevalence, incidence and mortality from cardiovascular disease in patients with pooled and specific severe mental illness: a large-scale meta-analysis of 3,211,768 patients and 113,383,368 controls. <i>World Psychiatry</i> , 2017, 16, 163-180.	12.0	1,206
237	Physical activity and sedentary behavior in people with major depressive disorder: A systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2017, 210, 139-150.	4.7	452
238	Perceived Stress and Its Relationship With Chronic Medical Conditions and Multimorbidity Among 229,293 Community-Dwelling Adults in 44 Low- and Middle-Income Countries. <i>American Journal of Epidemiology</i> , 2017, 186, 979-989.	3.6	68
239	Higher cardio-respiratory fitness is associated with increased mental and physical quality of life in people with bipolar disorder: A controlled pilot study. <i>Psychiatry Research</i> , 2017, 256, 219-224.	3.4	20
240	Interest, competence, appearance, fitness and social relatedness as motives for physical activity in Ugandan outpatients with psychosis. <i>Mental Health and Physical Activity</i> , 2017, 13, 94-99.	2.2	8
241	Physical activity correlates among 24,230 people with depression across 46 low- and middle-income countries. <i>Journal of Affective Disorders</i> , 2017, 221, 81-88.	4.7	23
242	Lower cardiorespiratory fitness is associated with more time spent sedentary in first episode psychosis: A pilot study. <i>Psychiatry Research</i> , 2017, 253, 13-17.	3.4	10
243	Bidirectional longitudinal relationship between leisure-time physical activity and psychotropic medication usage: A register linked follow-up study. <i>Psychiatry Research</i> , 2017, 247, 208-213.	3.4	13
244	Sedentary behavior and physical activity levels in people with schizophrenia, bipolar disorder and major depressive disorder: a global systematic review and meta-analysis. <i>World Psychiatry</i> , 2017, 16, 308-315.	12.0	664
245	Yoga versus non-standard care for schizophrenia. <i>The Cochrane Library</i> , 2017, 2017, .	2.4	22
246	Physical activity is associated with the physical, psychological, social and environmental quality of life in people with mental health problems in a low resource setting. <i>Psychiatry Research</i> , 2017, 258, 250-254.	3.4	26
247	Physical activity correlates in heavy episodic drinkers: Data from 46 low- and middle-income countries. <i>Mental Health and Physical Activity</i> , 2017, 13, 163-170.	2.2	7
248	Lifetime self-reported arthritis is associated with elevated levels of mental health burden: A multi-national cross sectional study across 46 low- and middle-income countries. <i>Scientific Reports</i> , 2017, 7, .	3.7	16
249	Mild cognitive impairment and physical activity in the general population: Findings from six low- and middle-income countries. <i>Experimental Gerontology</i> , 2017, 100, 100-105.	3.8	44
250	Physical activity correlates in people with anxiety: Data from 46 low- and middle-income countries. <i>General Hospital Psychiatry</i> , 2017, 49, 26-31.	2.5	14
251	Depression and pain: primary data and meta-analysis among 237,952 people across 47 low- and middle-income countries. <i>Psychological Medicine</i> , 2017, 47, 2906-2917.	4.6	70
252	Physical health policies and metabolic screening in mental health care systems of sub-Saharan African countries: a systematic review. <i>International Journal of Mental Health Systems</i> , 2017, 11, .	3.2	14

#	ARTICLE	IF	CITATIONS
253	Treatment Guidelines for Depression: Greater Emphasis on Physical Activity is Needed. <i>European Psychiatry</i> , 2017, 40, 1-3.	0.3	59
254	The relationship between chronic physical conditions, multimorbidity and anxiety in the general population: A global perspective across 42 countries. <i>General Hospital Psychiatry</i> , 2017, 45, 1-6.	2.5	78
255	Physical activity and anxiety: A perspective from the World Health Survey. <i>Journal of Affective Disorders</i> , 2017, 208, 545-552.	4.7	220
256	Dropout from physical activity interventions in people living with HIV: a systematic review and meta-analysis. <i>AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV</i> , 2017, 29, 636-643.	1.5	40
257	Physical activity and metabolic disease among people with affective disorders: Prevention, management and implementation. <i>Journal of Affective Disorders</i> , 2017, 224, 87-94.	4.7	17
258	Physical fitness in people with posttraumatic stress disorder: a systematic review. <i>Disability and Rehabilitation</i> , 2017, 39, 2461-2467.	2.7	35
259	More Reasons to Move: Exercise in the Treatment of Alcohol Use Disorders. <i>Frontiers in Psychiatry</i> , 2017, 8, .	2.7	27
260	A systematic review of physical activity policy recommendations and interventions for people with mental health problems in Sub-Saharan African countries. <i>Pan African Medical Journal</i> , 2017, 26, .	0.7	20
261	Correlates of physical activity among community-dwelling adults aged 50 or over in six low- and middle-income countries. <i>PLoS ONE</i> , 2017, 12, e0186992.	2.5	29
262	Depression comorbid with tuberculosis and its impact on health status: cross-sectional analysis of community-based data from 48 low- and middle-income countries. <i>BMC Medicine</i> , 2017, 15, .	7.4	92
263	Physical chronic conditions, multimorbidity and sedentary behavior amongst middle-aged and older adults in six low- and middle-income countries. <i>International Journal of Behavioral Nutrition and Physical Activity</i> , 2017, 14, .	4.5	75
264	Prevalence of suicidal ideation and attempt: associations with psychiatric disorders and HIV/AIDS in post-conflict Northern Uganda. <i>African Health Sciences</i> , 2017, 16, 1027.	0.8	18
265	Exercise and Posttraumatic Stress Disorder. , 2017, , 1-13.		0
266	Yoga vs Standard Care for Schizophrenia: Table 1.. <i>Schizophrenia Bulletin</i> , 2016, , sbv165.	4.3	3
267	Prevalence and clinical features associated with bipolar disorder polypharmacy: a systematic review. <i>Neuropsychiatric Disease and Treatment</i> , 2016, , 719.	2.5	77
268	Exercise is medicine for depression: even when the &ldquo;pill&rdquo; is small. <i>Neuropsychiatric Disease and Treatment</i> , 2016, Volume 12, 2715-2721.	2.5	16
269	Type 2 Diabetes Mellitus. , 2016, , 255-278.		3
270	Exercise for depression in older adults: a meta-analysis of randomized controlled trials adjusting for publication bias. <i>Revista Brasileira De Psiquiatria</i> , 2016, 38, 247-254.	1.1	161

#	ARTICLE	IF	CITATIONS
271	Depression and Reduced Bone Mineral Density at the Hip and Lumbar Spine. <i>Psychosomatic Medicine</i> , 2016, 78, 492-500.	2.3	20
272	Type 2 Diabetes Among People With Posttraumatic Stress Disorder. <i>Psychosomatic Medicine</i> , 2016, 78, 465-473.	2.3	77
273	The association between smoking prevalence and eating disorders: a systematic review and meta-analysis. <i>Addiction</i> , 2016, 111, 1914-1922.	6.1	76
274	Among Inpatients, Posttraumatic Stress Disorder Symptom Severity Is Negatively Associated With Time Spent Walking. <i>Journal of Nervous and Mental Disease</i> , 2016, 204, 15-19.	1.1	14
275	How can we increase physical activity and exercise among youth experiencing first episode psychosis? A systematic review of intervention variables. <i>Microbial Biotechnology</i> , 2016, 10, 435-440.	2.2	21
276	Motivating factors and barriers towards exercise in severe mental illness: a systematic review and meta-analysis. <i>Psychological Medicine</i> , 2016, 46, 2869-2881.	4.6	371
277	Sitting time, physical fitness impairments and metabolic abnormalities in people with bipolar disorder: An exploratory study. <i>Psychiatry Research</i> , 2016, 242, 7-12.	3.4	19
278	Autonomous motivation is associated with the maintenance stage of behaviour change in people with affective disorders. <i>Psychiatry Research</i> , 2016, 240, 267-271.	3.4	32
279	Physical activity and sedentary behavior in people with bipolar disorder: A systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2016, 201, 145-152.	4.7	117
280	Cardiorespiratory fitness in outpatients with bipolar disorder versus matched controls: An exploratory study. <i>Journal of Affective Disorders</i> , 2016, 199, 1-5.	4.7	22
281	Depressive symptoms and muscular fitness contribute independently to the ability to perform daily life activities in people with bipolar disorder. <i>Nordic Journal of Psychiatry</i> , 2016, 70, 477-482.	1.8	15
282	Exercise improves physical and psychological quality of life in people with depression: A meta-analysis including the evaluation of control group response. <i>Psychiatry Research</i> , 2016, 241, 47-54.	3.4	123
283	Is Pain Perception Altered in People With Depression? A Systematic Review and Meta-Analysis of Experimental Pain Research. <i>Journal of Pain</i> , 2016, 17, 1257-1272.	1.4	117
284	The epidemiology of back pain and its relationship with depression, psychosis, anxiety, sleep disturbances, and stress sensitivity: Data from 43 low- and middle-income countries. <i>General Hospital Psychiatry</i> , 2016, 43, 63-70.	2.5	126
285	Physical activity as a vital sign in patients with bipolar disorder. <i>Psychiatry Research</i> , 2016, 246, 218-222.	3.4	18
286	Dropout from physical activity interventions in children and adolescents with attention deficit hyperactivity disorder: A systematic review and meta-analysis. <i>Mental Health and Physical Activity</i> , 2016, 11, 46-52.	2.2	11
287	Cardiorespiratory fitness levels and moderators in people with HIV: A systematic review and meta-analysis. <i>Preventive Medicine</i> , 2016, 93, 106-114.	2.9	36
288	Exercise improves depressive symptoms in older adults: An umbrella review of systematic reviews and meta-analyses. <i>Psychiatry Research</i> , 2016, 244, 202-209.	3.4	139

#	ARTICLE	IF	CITATIONS
289	Diabetes mellitus in people with schizophrenia, bipolar disorder and major depressive disorder: a systematic review and large scale meta-analysis. <i>World Psychiatry</i> , 2016, 15, 166-174.	12.0	496
290	Is pain sensitivity altered in people with Alzheimer's disease? A systematic review and meta-analysis of experimental pain research. <i>Experimental Gerontology</i> , 2016, 82, 30-38.	3.8	25
291	Physical multimorbidity and psychosis: comprehensive cross sectional analysis including 242,952 people across 48 low- and middle-income countries. <i>BMC Medicine</i> , 2016, 14, .	7.4	126
292	Lower Bone Mineral Density at the Hip and Lumbar Spine in People with Psychosis Versus Controls: a Comprehensive Review and Skeletal Site-Specific Meta-analysis. <i>Current Osteoporosis Reports</i> , 2016, 14, 249-259.	4.9	28
293	The prevalence of diabetes mellitus type 2 in people with alcohol use disorders: a systematic review and large scale meta-analysis. <i>Psychiatry Research</i> , 2016, 246, 394-400.	3.4	30
294	Moving beyond the weight-loss paradigm of exercise interventions for mental illness. <i>Psychiatry Research</i> , 2016, 246, 392-393.	3.4	4
295	Are lower levels of cardiorespiratory fitness associated with incident depression? A systematic review of prospective cohort studies. <i>Preventive Medicine</i> , 2016, 93, 159-165.	2.9	89
296	The Functional Exercise Capacity Is Associated With Global Functioning in People With Bipolar Disorder. <i>Journal of Nervous and Mental Disease</i> , 2016, 204, 673-677.	1.1	9
297	The Prevalence of Metabolic Syndrome in Alcohol Use Disorders: A Systematic Review and Meta-analysis. <i>Alcohol and Alcoholism</i> , 2016, 51, 515-521.	1.9	80
298	Impact of antipsychotic medication on physical activity and physical fitness in adolescents: An exploratory study. <i>Psychiatry Research</i> , 2016, 242, 192-197.	3.4	24
299	How much physical activity do people with schizophrenia engage in? A systematic review, comparative meta-analysis and meta-regression. <i>Schizophrenia Research</i> , 2016, 176, 431-440.	2.4	288
300	What are the top 10 physical activity research questions in schizophrenia?. <i>Disability and Rehabilitation</i> , 2016, 38, 2235-2243.	2.7	30
301	Type 2 Diabetes Mellitus in Youth Exposed to Antipsychotics. <i>JAMA Psychiatry</i> , 2016, 73, 247.	14.1	191
302	Prevalence and predictors of treatment dropout from physical activity interventions in schizophrenia: a meta-analysis. <i>General Hospital Psychiatry</i> , 2016, 39, 15-23.	2.5	174
303	Physical activity as a vital sign in patients with schizophrenia: Evidence and clinical recommendations. <i>Schizophrenia Research</i> , 2016, 170, 336-340.	2.4	35
304	Top 10 research questions to promote physical activity in bipolar disorders: A consensus statement from the International Organization of Physical Therapists in Mental Health. <i>Journal of Affective Disorders</i> , 2016, 195, 82-87.	4.7	19
305	Concurrent validity of the international physical activity questionnaire in outpatients with bipolar disorder: Comparison with the Sensewear Armband. <i>Psychiatry Research</i> , 2016, 237, 122-126.	3.4	15
306	How common is bipolar disorder in general primary care attendees? A systematic review and meta-analysis investigating prevalence determined according to structured clinical assessments. <i>Australian and New Zealand Journal of Psychiatry</i> , 2016, 50, 631-639.	2.7	30

#	ARTICLE	IF	CITATIONS
307	The prevalence and predictors of obstructive sleep apnea in major depressive disorder, bipolar disorder and schizophrenia: A systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2016, 197, 259-267.	4.7	131
308	Exercise as a treatment for depression: A meta-analysis adjusting for publication bias. <i>Journal of Psychiatric Research</i> , 2016, 77, 42-51.	3.1	992
309	A comparison of physical fitness in patients with bipolar disorder, schizophrenia and healthy controls. <i>Disability and Rehabilitation</i> , 2016, 38, 2047-2051.	2.7	26
310	Top 10 research questions to promote physical activity research in people with binge eating disorder. <i>Eating Disorders</i> , 2016, 24, 326-337.	4.0	4
311	A narrative synthesis investigating the use and value of social support to promote physical activity among individuals with schizophrenia. <i>Disability and Rehabilitation</i> , 2016, 38, 123-150.	2.7	36
312	Dropout from exercise randomized controlled trials among people with depression: A meta-analysis and meta regression. <i>Journal of Affective Disorders</i> , 2016, 190, 457-466.	4.7	217
313	Exercise improves cardiorespiratory fitness in people with depression: A meta-analysis of randomized control trials. <i>Journal of Affective Disorders</i> , 2016, 190, 249-253.	4.7	140
314	Are leptin levels increased among people with schizophrenia versus controls? A systematic review and comparative meta-analysis. <i>Psychoneuroendocrinology</i> , 2016, 63, 144-154.	2.5	56
315	Considering the Role of Physical Therapists Within the Treatment and Rehabilitation of Individuals With Eating Disorders: An International Survey of Expert Clinicians. <i>Physiotherapy Research International</i> , 2016, 21, 237-246.	1.6	8
316	Cardiorespiratory Fitness in Severe Mental Illness: A Systematic Review and Meta-analysis. <i>Sports Medicine</i> , 2016, 47, 343-352.	6.7	176
317	Validity of the 6min walk test in outpatients with bipolar disorder. <i>Psychiatry Research</i> , 2015, 230, 664-667.	3.4	9
318	Decreased pain sensitivity among people with schizophrenia. <i>Pain</i> , 2015, 156, 2121-2131.	4.6	104
319	Yoga versus standard care for schizophrenia. <i>The Cochrane Library</i> , 2015, 2015, .	2.4	32
320	TYPE 2 DIABETES IN PATIENTS WITH MAJOR DEPRESSIVE DISORDER: A META-ANALYSIS OF PREVALENCE ESTIMATES AND PREDICTORS. <i>Depression and Anxiety</i> , 2015, 32, 763-773.	3.9	139
321	Health-related quality of life and aerobic fitness in people with schizophrenia. <i>International Journal of Mental Health Nursing</i> , 2015, 24, 394-402.	3.6	35
322	The Psychosocial Consequences of Sports Participation for Individuals with Severe Mental Illness: A Metasynthesis Review. <i>Advances in Psychiatry</i> , 2015, 2015, 1-8.	0.7	18
323	Risk of metabolic syndrome and its components in people with schizophrenia and related psychotic disorders, bipolar disorder and major depressive disorder: a systematic review and meta-analysis. <i>World Psychiatry</i> , 2015, 14, 339-347.	12.0	889
324	No evidence of a control group response in exercise randomised controlled trials in people with schizophrenia: A systematic review and meta-analysis. <i>Psychiatry Research</i> , 2015, 229, 840-843.	3.4	3

#	ARTICLE	IF	CITATIONS
325	Exercise improves cardiorespiratory fitness in people with schizophrenia: A systematic review and meta-analysis. <i>Schizophrenia Research</i> , 2015, 169, 453-457.	2.4	91
326	Integrating physical activity as medicine in the care of people with severe mental illness. <i>Australian and New Zealand Journal of Psychiatry</i> , 2015, 49, 681-682.	2.7	68
327	Six minutes walk test for individuals with schizophrenia. <i>Disability and Rehabilitation</i> , 2015, 37, 921-927.	2.7	33
328	Schizophrenia and the risk of fractures: a systematic review and comparative meta-analysis. <i>General Hospital Psychiatry</i> , 2015, 37, 126-133.	2.5	79
329	A Systematic Review of Physical Activity Correlates in Alcohol Use Disorders. <i>Archives of Psychiatric Nursing</i> , 2015, 29, 196-201.	2.0	38
330	Do people with mental illness receive adequate smoking cessation advice? A systematic review and meta-analysis. <i>General Hospital Psychiatry</i> , 2015, 37, 14-23.	2.5	73
331	A systematic review of the benefits of physical therapy within a multidisciplinary care approach for people with schizophrenia: An update. <i>Psychiatry Research</i> , 2015, 229, 828-839.	3.4	50
332	Could autonomous motivation hold the key to successfully implementing lifestyle changes in affective disorders? A multicentre cross sectional study. <i>Psychiatry Research</i> , 2015, 228, 100-106.	3.4	50
333	Test-retest reliability, feasibility and clinical correlates of the Eurofit test battery in people with bipolar disorder. <i>Psychiatry Research</i> , 2015, 228, 620-625.	3.4	23
334	The functional exercise capacity in patients with bipolar disorder versus healthy controls: A pilot study. <i>Psychiatry Research</i> , 2015, 229, 194-199.	3.4	20
335	Health-related physical fitness in patients with bipolar disorder vs. healthy controls: An exploratory study. <i>Journal of Affective Disorders</i> , 2015, 177, 22-27.	4.7	25
336	Why moving more should be promoted for severe mental illness. <i>Lancet Psychiatry</i> , 2015, 2, 295.	9.0	39
337	Aerobic exercise capacity: an important correlate of psychosocial function in first episode psychosis. <i>Acta Psychiatrica Scandinavica</i> , 2015, 131, 234-234.	4.7	18
338	Associations Between Metabolic and Aerobic Fitness Parameters in Patients With Schizophrenia. <i>Journal of Nervous and Mental Disease</i> , 2015, 203, 23-27.	1.1	18
339	Identifying the facilitators and processes which influence recovery in individuals with schizophrenia: a systematic review and thematic synthesis. <i>Journal of Mental Health</i> , 2015, 24, 103-110.	3.0	63
340	Physical activity in the treatment of Post-traumatic stress disorder: A systematic review and meta-analysis. <i>Psychiatry Research</i> , 2015, 230, 130-136.	3.4	236
341	Aerobic capacity is associated with global functioning in people with schizophrenia. <i>Journal of Mental Health</i> , 2015, 24, 214-218.	3.0	17
342	The prevalence and risk of metabolic syndrome and its components among people with posttraumatic stress disorder: a systematic review and meta-analysis. <i>Metabolism: Clinical and Experimental</i> , 2015, 64, 926-933.	5.0	184

#	ARTICLE	IF	CITATIONS
343	Adopting and maintaining physical activity behaviours in people with severe mental illness: The importance of autonomous motivation. <i>Preventive Medicine</i> , 2015, 81, 216-220.	2.9	91
344	Negative symptoms are associated with lower autonomous motivation towards physical activity in people with schizophrenia. <i>Comprehensive Psychiatry</i> , 2015, 56, 128-132.	4.4	75
345	Exercise therapy improves both mental and physical health in patients with major depression. <i>Disability and Rehabilitation</i> , 2015, 37, 1490-1495.	2.7	257
346	The functional exercise capacity and its correlates in obese treatment-seeking people with binge eating disorder: an exploratory study. <i>Disability and Rehabilitation</i> , 2015, 37, 777-782.	2.7	20
347	Challenges Establishing the Efficacy of Exercise as an Antidepressant Treatment: A Systematic Review and Meta-Analysis of Control Group Responses in Exercise Randomised Controlled Trials. <i>Sports Medicine</i> , 2015, 46, 699-713.	6.7	46
348	Prevalence and Predictors of Type 2 Diabetes Mellitus in People With Bipolar Disorder. <i>Journal of Clinical Psychiatry</i> , 2015, 76, 1490-1499.	2.9	91
349	The Assessment, Benefits and Delivery of Physical Activity in People with Schizophrenia: A Survey of Members of the International Organization of Physical Therapists in Mental Health. <i>Physiotherapy Research International</i> , 2014, 19, 248-256.	1.6	14
350	Metabolic syndrome and metabolic abnormalities in patients with major depressive disorder: a meta-analysis of prevalences and moderating variables. <i>Psychological Medicine</i> , 2014, 44, 2017-2028.	4.6	237
351	The value of social support to encourage people with schizophrenia to engage in physical activity: an international insight from specialist mental health physiotherapists. <i>Journal of Mental Health</i> , 2014, 23, 256-260.	3.0	27
352	Physical Activity Correlates in Persons with Binge Eating Disorder: A Systematic Review. <i>European Eating Disorders Review</i> , 2014, 22, 1-8.	4.2	36
353	Reliability and clinical correlates of the Astrand "Rhyming sub-maximal exercise test in patients with schizophrenia or schizoaffective disorder. <i>Psychiatry Research</i> , 2014, 220, 778-783.	3.4	27
354	The prevalence and moderators of clinical pain in people with schizophrenia: A systematic review and large scale meta-analysis. <i>Schizophrenia Research</i> , 2014, 160, 1-8.	2.4	63
355	Self-determination and stage of readiness to change physical activity behaviour in schizophrenia. <i>Mental Health and Physical Activity</i> , 2014, 7, 171-176.	2.2	41
356	Understanding the role of physiotherapists in schizophrenia: an international perspective from members of the International Organisation of Physical Therapists in Mental Health (IOPTMH). <i>Journal of Mental Health</i> , 2014, 23, 125-129.	3.0	35
357	Barriers to and Facilitators of Physical Activity Among Persons With Schizophrenia: A Survey of Physical Therapists. <i>Psychiatric Services</i> , 2014, 65, 693-696.	2.4	90
358	Physiotherapists can help implement physical activity programmes in clinical practice. <i>British Journal of Psychiatry</i> , 2014, 204, 164-164.	2.3	52
359	A Clinical Review of the Treatment of Catatonia. <i>Frontiers in Psychiatry</i> , 2014, 5, .	2.7	213
360	A systematic review of physical therapy interventions for patients with anorexia and bulimia nervosa. <i>Disability and Rehabilitation</i> , 2014, 36, 628-634.	2.7	90

#	ARTICLE	IF	CITATIONS
361	The importance of physiotherapy within the multidisciplinary treatment of eating disorders. <i>Fisioterapia</i> , 2014, 36, 55-57.	0.1	1
362	Selection, Use and Psychometric Properties of Physical Activity Measures to Assess Individuals with Severe Mental Illness: A Narrative Synthesis. <i>Archives of Psychiatric Nursing</i> , 2014, 28, 135-151.	2.0	120
363	Neurobiological effects of physical exercise in schizophrenia: a systematic review. <i>Disability and Rehabilitation</i> , 2014, 36, 1749-1754.	2.7	61
364	Markers of inflammation in schizophrenia: association vs. causation. <i>World Psychiatry</i> , 2014, 13, 189-192.	12.0	54
365	What are the factors associated with physical activity (PA) participation in community dwelling adults with dementia? A systematic review of PA correlates. <i>Archives of Gerontology and Geriatrics</i> , 2014, 59, 195-203.	3.5	64
366	The transcending benefits of physical activity for individuals with schizophrenia: A systematic review and meta-ethnography. <i>Psychiatry Research</i> , 2014, 220, 11-19.	3.4	92
367	Metabolic syndrome and lung function in schizophrenia: A pilot study. <i>Psychiatry Research</i> , 2014, 220, 58-62.	3.4	12
368	Health related quality of life, physical fitness and physical activity participation in treatment-seeking obese persons with and without binge eating disorder. <i>Psychiatry Research</i> , 2014, 216, 97-102.	3.4	38
369	Associations between perceived neighbourhood environmental attributes and self-reported sitting time in patients with schizophrenia: A pilot study. <i>Psychiatry Research</i> , 2014, 215, 33-38.	3.4	24
370	Associations between expiratory spirometry parameters and limitations in daily life activities in patients with schizophrenia. <i>General Hospital Psychiatry</i> , 2014, 36, 172-176.	2.5	15
371	Addressing the disparity in physical health provision for people with schizophrenia: an important role for physiotherapists. <i>Physiotherapy</i> , 2014, 100, 185-186.	0.6	15
372	Changes in physical activity, physical fitness, self-perception and quality of life following a 6-month physical activity counseling and cognitive behavioral therapy program in outpatients with binge eating disorder. <i>Psychiatry Research</i> , 2014, 219, 361-366.	3.4	32
373	The benefits of walking for individuals with schizophrenia spectrum disorders: A systematic review. <i>International Journal of Therapy and Rehabilitation</i> , 2014, 21, 410-420.	0.4	30
374	Physical activity and serious mental illness: A multidisciplinary call to action. <i>Mental Health and Physical Activity</i> , 2014, 7, 153-154.	2.2	22
375	An impaired health related muscular fitness contributes to a reduced walking capacity in patients with schizophrenia: a cross-sectional study. <i>BMC Psychiatry</i> , 2013, 13, .	3.4	27
376	SOCIAL ANXIETY IN PHYSICAL ACTIVITY PARTICIPATION IN PATIENTS WITH MENTAL ILLNESS: A CROSS-SECTIONAL MULTICENTER STUDY. <i>Depression and Anxiety</i> , 2013, 30, 757-762.	3.9	19
377	The importance of self-determined motivation towards physical activity in patients with schizophrenia. <i>Psychiatry Research</i> , 2013, 210, 812-818.	3.4	128
378	Neurocognition in clinical high risk young adults who did or did not convert to a first schizophrenic psychosis: A meta-analysis. <i>Schizophrenia Research</i> , 2013, 149, 48-55.	2.4	96

#	ARTICLE	IF	CITATIONS
379	A systematic review on physical therapy interventions for patients with binge eating disorder. <i>Disability and Rehabilitation</i> , 2013, 35, 2191-2196.	2.7	59
380	Progressive muscle relaxation in persons with schizophrenia: a systematic review of randomized controlled trials. <i>Clinical Rehabilitation</i> , 2013, 27, 291-298.	3.2	44
381	A review of physical activity correlates in patients with bipolar disorder. <i>Journal of Affective Disorders</i> , 2013, 145, 285-291.	4.7	102
382	Relationships between physical fitness, physical activity, smoking and metabolic and mental health parameters in people with schizophrenia. <i>Psychiatry Research</i> , 2013, 207, 25-32.	3.4	127
383	Associations between physical activity and the built environment in patients with schizophrenia: a multi-centre study. <i>General Hospital Psychiatry</i> , 2013, 35, 653-658.	2.5	35
384	There is a need for physiotherapists in the multidisciplinary treatment of schizophrenia. <i>Fisioterapia</i> , 2013, 35, 239-242.	0.1	1
385	Response to Bartoli et al.. <i>American Journal of Psychiatry</i> , 2013, 170, 928-929.	10.5	2
386	Is the Prevalence of Metabolic Syndrome and Metabolic Abnormalities Increased in Early Schizophrenia? A Comparative Meta-Analysis of First Episode, Untreated and Treated Patients. <i>Schizophrenia Bulletin</i> , 2013, 39, 295-305.	4.3	262
387	Physiotherapy for patients with anorexia nervosa. <i>Advances in Eating Disorders (Abingdon, England)</i> , 2013, 1, 224-238.	0.7	18
388	A meta-analysis of cardio-metabolic abnormalities in drug naïve, first-episode and multi-episode patients with schizophrenia versus general population controls. <i>World Psychiatry</i> , 2013, 12, 240-250.	12.0	224
389	Metabolic Syndrome and Metabolic Abnormalities in Bipolar Disorder: A Meta-Analysis of Prevalence Rates and Moderators. <i>American Journal of Psychiatry</i> , 2013, 170, 265-274.	10.5	328
390	Diabetes, physical activity participation and exercise capacity in patients with schizophrenia. <i>Psychiatry and Clinical Neurosciences</i> , 2013, 67, 451-456.	3.2	26
391	Prevalence of Metabolic Syndrome and Metabolic Abnormalities in Schizophrenia and Related Disorders—A Systematic Review and Meta-Analysis. <i>Schizophrenia Bulletin</i> , 2013, 39, 306-318.	4.3	821
392	Physical activity and sedentary behaviour in outpatients with schizophrenia: A systematic review and meta-analysis. <i>International Journal of Therapy and Rehabilitation</i> , 2013, 20, 588-595.	0.4	73
393	Systematic Review of the Benefits of Physical Therapy Within a Multidisciplinary Care Approach for People With Schizophrenia. <i>Physical Therapy</i> , 2012, 92, 11-23.	2.7	111
394	Guideline concordant monitoring of metabolic risk in people treated with antipsychotic medication: systematic review and meta-analysis of screening practices. <i>Psychological Medicine</i> , 2012, 42, 125-147.	4.6	276
395	A Cognitive- Behavioral Therapeutic Program for Patients With Obesity and Binge Eating Disorder. <i>Behavior Modification</i> , 2012, 36, 670-686.	2.5	25
396	Authors' reply. <i>British Journal of Psychiatry</i> , 2012, 201, 76-77.	2.3	1

#	ARTICLE	IF	CITATIONS
397	International Organization of Physical Therapy in Mental Health consensus on physical activity within multidisciplinary rehabilitation programmes for minimising cardio-metabolic risk in patients with schizophrenia. <i>Disability and Rehabilitation</i> , 2012, 34, 1-12.	2.7	121
398	Eurofit test battery in patients with schizophrenia or schizoaffective disorder: Reliability and clinical correlates. <i>European Psychiatry</i> , 2012, 27, 416-421.	0.3	28
399	Associations between sedentary behaviour and metabolic parameters in patients with schizophrenia. <i>Psychiatry Research</i> , 2012, 200, 73-78.	3.4	113
400	State anxiety, psychological stress and positive well-being responses to yoga and aerobic exercise in people with schizophrenia: a pilot study. <i>Disability and Rehabilitation</i> , 2011, 33, 684-689.	2.7	91
401	A systematic evaluation and comparison of the guidelines for screening and monitoring of cardiometabolic risk in people with schizophrenia. <i>European Psychiatry</i> , 2011, 26, 2189-2189.	0.3	1
402	Lack of physical activity during leisure time contributes to an impaired health related quality of life in patients with schizophrenia. <i>Schizophrenia Research</i> , 2011, 129, 122-127.	2.4	87
403	Association of the metabolic syndrome with physical activity performance in patients with schizophrenia. <i>Diabetes and Metabolism</i> , 2011, 37, 318-323.	3.5	51
404	Reliability, minimal detectable changes, practice effects and correlates of the 6-min walk test in patients with schizophrenia. <i>Psychiatry Research</i> , 2011, 187, 62-67.	3.4	48
405	Effects of progressive muscle relaxation on state anxiety and subjective well-being in people with schizophrenia: a randomized controlled trial. <i>Clinical Rehabilitation</i> , 2011, 25, 567-575.	3.2	52
406	Guidelines for screening and monitoring of cardiometabolic risk in schizophrenia: systematic evaluation. <i>British Journal of Psychiatry</i> , 2011, 199, 99-105.	2.3	167
407	Physical activity participation, functional exercise capacity and self-esteem in patients with schizophrenia. <i>International Journal of Therapy and Rehabilitation</i> , 2011, 18, 222-229.	0.4	6
408	Quality Assessment of Physical Activity Recommendations Within Clinical Practice Guidelines for the Prevention and Treatment of Cardio-metabolic Risk Factors in People With Schizophrenia. <i>Community Mental Health Journal</i> , 2011, 47, 703-710.	1.3	21
409	De eetbuistoornis. <i>Bijblijven (Amsterdam, Netherlands)</i> , 2010, 26, 43-48.	0.0	0
410	De psychotherapeutische behandeling van patiënten met de eetbuistoornis: een literatuuroverzicht. <i>Dth</i> , 2010, 30, 203-221.	0.2	0
411	Considering a frame of reference for physical activity research related to the cardiometabolic risk profile in schizophrenia. <i>Psychiatry Research</i> , 2010, 177, 271-279.	3.4	105
412	Cardiometabolic effects of physical activity interventions for people with schizophrenia. <i>Physical Therapy Reviews</i> , 2009, 14, 388-398.	1.1	47
413	State anxiety and subjective well-being responses to acute bouts of aerobic exercise in patients with depressive and anxiety disorders. <i>British Journal of Sports Medicine</i> , 2009, 43, 756-759.	7.7	50
414	Een cognitief-gedragstherapeutisch programma voor patiënten met obesitas en binge eating disorder: kortetermijnresultaten. <i>Dth</i> , 2009, 29, 112-126.	0.2	2

#	ARTICLE	IF	CITATIONS
415	Metabolic syndrome in people with schizophrenia: a review. <i>World Psychiatry</i> , 2009, 8, 15-22.	12.0	480
416	Yoga as part of a package of care versus non-standard care for schizophrenia. <i>The Cochrane Library</i> , 0, , .	2.4	2
417	Yoga for schizophrenia. <i>The Cochrane Library</i> , 0, , .	2.4	1
418	A 10-hours workshop improves physical activity prescription for mental illness knowledge and confidence in health care professionals: a nation-wide multicentre study from Uganda. <i>Disability and Rehabilitation</i> , 0, , 1-6.	2.7	6
419	Interaction styles of health care professionals supporting patients with chronic conditions in general hospitals towards self-management: one-year follow-up data from a brief educational intervention. <i>International Journal of Health Promotion and Education</i> , 0, , 1-12.	1.2	1
420	Barriers and facilitators of physical activity participation in adolescents with autism. <i>Children's Health Care</i> , 0, , 1-44.	0.8	4