

# David Moscatelli

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4610930/publications.pdf>

Version: 2024-02-01

54  
papers

4,360  
citations

201385

27  
h-index

197535

49  
g-index

56  
all docs

56  
docs citations

56  
times ranked

2958  
citing authors

#	ARTICLE	IF	CITATIONS
1	The Fgf Family of Growth Factors and Oncogenes. <i>Advances in Cancer Research</i> , 1992, 59, 115-165.	1.9	1,016
2	High and low affinity binding sites for basic fibroblast growth factor on cultured cells: Absence of a role for low affinity binding in the stimulation of plasminogen activator production by bovine capillary endothelial cells. <i>Journal of Cellular Physiology</i> , 1987, 131, 123-130.	2.0	636
3	Proximal location of mouse prostate epithelial stem cells. <i>Journal of Cell Biology</i> , 2002, 157, 1257-1265.	2.3	298
4	Sca-1 expression identifies stem cells in the proximal region of prostatic ducts with high capacity to reconstitute prostatic tissue. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 7180-7185.	3.3	249
5	Both normal and tumor cells produce basic fibroblast growth factor. <i>Journal of Cellular Physiology</i> , 1986, 129, 273-276.	2.0	234
6	Role of extracellular matrix in the action of basic fibroblast growth factor: Matrix as a source of growth factor for long-term stimulation of plasminogen activator production and DNA synthesis. <i>Journal of Cellular Physiology</i> , 1989, 140, 75-81.	2.0	229
7	Membrane and matrix localization of proteinases: a common theme in tumor cell invasion and angiogenesis. <i>Biochimica Et Biophysica Acta: Reviews on Cancer</i> , 1988, 948, 67-85.	3.3	138
8	High Aldehyde Dehydrogenase Activity: A Novel Functional Marker of Murine Prostate Stem/Progenitor Cells. <i>Stem Cells</i> , 2009, 27, 2220-2228.	1.4	128
9	TGF- $\beta$ 2 maintains dormancy of prostatic stem cells in the proximal region of ducts. <i>Journal of Cell Biology</i> , 2005, 170, 81-90.	2.3	124
10	A form of human basic fibroblast growth factor with an extended amino terminus. <i>Biochemical and Biophysical Research Communications</i> , 1987, 144, 543-550.	1.0	112
11	Lack of ERK activation and cell migration in FGF-2 deficient endothelial cells. <i>FASEB Journal</i> , 2002, 16, 598-600.	0.2	106
12	Inflammatory Mediators Regulate Cathepsin S in Macrophages and Microglia: A Role in Attenuating Heparan Sulfate Interactions. <i>Molecular Medicine</i> , 1999, 5, 320-333.	1.9	82
13	Proximal Prostatic Stem Cells Are Programmed to Regenerate a Proximal-Distal Ductal Axis. <i>Stem Cells</i> , 2006, 24, 1859-1868.	1.4	81
14	Molecular Signatures of Prostate Stem Cells Reveal Novel Signaling Pathways and Provide Insights into Prostate Cancer. <i>PLoS ONE</i> , 2009, 4, e5722.	1.1	64
15	Increased hyaluronic acid production on stimulation of DNA synthesis in chick embryo fibroblasts. <i>Nature</i> , 1975, 254, 65-66.	13.7	55
16	Multiple forms of an angiogenesis factor: basic fibroblast growth factor. <i>Biochimie</i> , 1988, 70, 83-87.	1.3	54
17	Fibroblast Growth Factor-2 Can Mediate Cell Attachment by Linking Receptors and Heparan Sulfate Proteoglycans on Neighboring Cells. <i>Journal of Biological Chemistry</i> , 1995, 270, 24188-24196.	1.6	52
18	Hormonal control of hyaluronic acid production in fibroblasts and its relation to nucleic acid and protein synthesis. <i>Journal of Cellular Physiology</i> , 1977, 91, 79-88.	2.0	50

#	ARTICLE	IF	CITATIONS
19	Urokinase-type and tissue-type plasminogen activators have different distributions in cultured bovine capillary endothelial cells. <i>Journal of Cellular Biochemistry</i> , 1986, 30, 19-29.	1.2	45
20	Induction of stromelysin-1 (MMP-3) by fibroblast growth factor-2 (FGF-2) in FGF-2 microvascular endothelial cells requires prolonged activation of extracellular signal-regulated kinases-1 and -2 (ERK-1/2). <i>Journal of Cellular Biochemistry</i> , 2003, 90, 1015-1025.	1.2	40
21	The development of a quantitative RIA for basic fibroblast growth factor using polyclonal antibodies against the 157 amino acid form of human bFGF. <i>Journal of Immunological Methods</i> , 1988, 110, 183-192.	0.6	39
22	An amino-terminally extended and post-translationally modified form of a 25kD basic fibroblast growth factor. <i>Biochemical and Biophysical Research Communications</i> , 1989, 160, 1267-1274.	1.0	38
23	Expression of fibroblast growth factors and their receptors in acquired immunodeficiency syndrome-associated Kaposi sarcoma tissue and derived cells. <i>Cancer</i> , 1993, 72, 2253-2259.	2.0	38
24	Effects of depletion of K <sup>+</sup> , Na <sup>+</sup> , or Ca <sup>2+</sup> on DNA synthesis and cell cation content in chick embryo fibroblasts. <i>Journal of Cellular Physiology</i> , 1979, 101, 117-128.	2.0	35
25	Generation of active TGF- $\beta$ by prostatic cell cocultures using novel basal and luminal prostatic epithelial cell lines. <i>Journal of Cellular Physiology</i> , 2000, 184, 70-79.	2.0	31
26	Vascular endothelial growth factor and angiopoietin are required for prostate regeneration. <i>Prostate</i> , 2007, 67, 485-499.	1.2	31
27	Androgens modulate the balance between VEGF and angiopoietin expression in prostate epithelial and smooth muscle cells. <i>Prostate</i> , 2002, 50, 83-91.	1.2	30
28	Studies on FGF-2: Nuclear localization and function of high molecular weight forms and receptor binding in the absence of heparin. <i>Molecular Reproduction and Development</i> , 1994, 39, 102-105.	1.0	27
29	Transforming growth factor- $\beta$ is an autocrine mitogen for a novel androgen-responsive murine prostatic smooth muscle cell line, PSMC1. <i>Journal of Cellular Physiology</i> , 2000, 185, 416-424.	2.0	24
30	Turnover of Functional Basic Fibroblast Growth Factor Receptors on the Surface of BHK and NIH 3T3 Cells. <i>Growth Factors</i> , 1990, 3, 25-33.	0.5	23
31	Molecular Signatures of the Primitive Prostate Stem Cell Niche Reveal Novel Mesenchymal-Epithelial Signaling Pathways. <i>PLoS ONE</i> , 2010, 5, e13024.	1.1	23
32	New Observations on the Intracellular Localization and Release of bFGF. <i>Annals of the New York Academy of Sciences</i> , 1991, 638, 204-206.	1.8	21
33	Induction of Urokinase-type Plasminogen Activator by Fibroblast Growth Factor (FGF)-2 Is Dependent on Expression of FGF Receptors and Does Not Require Activation of Phospholipase C $\beta$ 1. <i>Journal of Biological Chemistry</i> , 1996, 271, 31154-31159.	1.6	21
34	Proteases and Angiogenesis: Production of Plasminogen Activator and Collagenase by Endothelial Cells. , 1982, , 191-197.		20
35	Interaction of Basic Fibroblast Growth Factor with Extracellular Matrix and Receptors. <i>Annals of the New York Academy of Sciences</i> , 1991, 638, 177-181.	1.8	19
36	Autocrine downregulation of fibroblast growth factor receptors in F9 teratocarcinoma cells. <i>Journal of Cellular Physiology</i> , 1994, 160, 555-562.	2.0	18

#	ARTICLE	IF	CITATIONS
37	Involvement of the conserved acidic amino acid domain of FGF receptor 1 in ligand-receptor interaction. <i>Journal of Cellular Physiology</i> , 1993, 157, 209-216.	2.0	15
38	Lipopolysaccharide inhibits activation of latent transforming growth factor- $\beta$ in bovine endothelial cells. <i>Journal of Cellular Physiology</i> , 1995, 163, 210-219.	2.0	14
39	Fibroblast Growth Factor (FGF)-2 Mediates Cell Attachment through Interactions with Two FGF Receptor-1 Isoforms and Extracellular Matrix or Cell-Associated Heparan Sulfate Proteoglycans. <i>Biochemical and Biophysical Research Communications</i> , 2000, 276, 399-405.	1.0	14
40	Retinal blood vessels develop in response to local VEGF-A signals in the absence of blood flow. <i>Experimental Eye Research</i> , 2005, 81, 147-158.	1.2	14
41	Vascular density is highest in the proximal region of the mouse prostate. <i>Prostate</i> , 2007, 67, 968-975.	1.2	14
42	Prostate cells express two isoforms of fibroblast growth factor receptor 1 with different affinities for fibroblast growth factor-2. <i>Prostate</i> , 2007, 67, 115-124.	1.2	12
43	Differentiation and stromal-induced growth promotion of murine prostatic tumors. <i>Prostate</i> , 2002, 51, 175-188.	1.2	11
44	Mechanisms Controlling the Extracellular Activity of Basic Fibroblast Growth Factor and Transforming Growth Factor $\beta$ 2a. <i>Annals of the New York Academy of Sciences</i> , 1991, 614, 250-258.	1.8	9
45	Stromal/epithelial interactions of murine prostatic cell lines in vivo: A model for benign prostatic hyperplasia and the effect of doxazosin on tissue size. <i>Prostate</i> , 2003, 54, 17-24.	1.2	8
46	Endothelial cells support the growth of prostate tissue in vivo. <i>Prostate</i> , 2008, 68, 893-901.	1.2	8
47	PINing Down the Origin of Prostate Cancer. <i>Science Translational Medicine</i> , 2010, 2, 43ps38.	5.8	6
48	Synthesis of collagenase and plasminogen activator by endothelial cells. <i>Developments in Cardiovascular Medicine</i> , 1984, , 429-437.	0.1	2
49	Transforming growth factor $\beta$ 2 is an autocrine mitogen for a novel androgen-responsive murine prostatic smooth muscle cell line, PSMC1. <i>Journal of Cellular Physiology</i> , 2000, 185, 416-424.	2.0	1
50	Bmi-1, stem cells and prostate carcinogenesis. <i>Asian Journal of Andrology</i> , 2011, 13, 353-354.	0.8	1
51	Basic FGF (FGF-2) is responsible for endothelial cell repair after mechanical damage: a genetic evidence. <i>Journal of the American College of Surgeons</i> , 2000, 191, S77-S78.	0.2	0
52	Effects of phosphodiesterase inhibitors on vascular development: Implications for retinopathy of prematurity. <i>Early Human Development</i> , 2008, 84, S81-S82.	0.8	0
53	BONE MARROW CELLS ARE ABLE TO GENERATE PROSTATIC EPITHELIAL AND STROMAL CELLS. <i>Journal of Urology</i> , 2009, 181, 41-42.	0.2	0
54	The Prostate Stem Cell Niche. , 2013, , 91-109.		0