

# E-Jean Tan

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4612736/publications.pdf>

Version: 2024-02-01

13  
papers

894  
citations

933447

10  
h-index

1199594

12  
g-index

14  
all docs

14  
docs citations

14  
times ranked

1586  
citing authors

#	ARTICLE	IF	CITATIONS
1	Cell-lineage controlled epigenetic regulation in glioblastoma stem cells determines functionally distinct subgroups and predicts patient survival. <i>Nature Communications</i> , 2022, 13, 2236.	12.8	7
2	A molecularly distinct subset of glioblastoma requires serum-containing media to establish sustainable bona fide glioblastoma stem cell cultures. <i>Glia</i> , 2020, 68, 1228-1240.	4.9	12
3	LGR5 promotes tumorigenicity and invasion of glioblastoma stem-like cells and is a potential therapeutic target for a subset of glioblastoma patients. <i>Journal of Pathology</i> , 2019, 247, 228-240.	4.5	19
4	Snail regulates BMP and TGF $\beta$ 2 pathways to control the differentiation status of glioma-initiating cells. <i>Oncogene</i> , 2018, 37, 2515-2531.	5.9	46
5	Epithelial-Mesenchymal Transition. , 2018, , .		0
6	Reprogramming during epithelial to mesenchymal transition under the control of TGF $\beta$ 2. <i>Cell Adhesion and Migration</i> , 2015, 9, 233-246.	2.7	82
7	The high mobility group A2 protein epigenetically silences the Cdh1 gene during epithelial-to-mesenchymal transition. <i>Nucleic Acids Research</i> , 2015, 43, 162-178.	14.5	69
8	p53 regulates epithelial-mesenchymal transition induced by transforming growth factor $\beta$ 2. <i>Journal of Cellular Physiology</i> , 2013, 228, 801-813.	4.1	37
9	Regulation of Transcription Factor Twist Expression by the DNA Architectural Protein High Mobility Group A2 during Epithelial-to-Mesenchymal Transition. <i>Journal of Biological Chemistry</i> , 2012, 287, 7134-7145.	3.4	94
10	PIG3: A novel link between oxidative stress and DNA damage response in cancer. <i>Cancer Letters</i> , 2012, 327, 97-102.	7.2	50
11	HMGA2 and Smads Co-regulate SNAIL1 Expression during Induction of Epithelial-to-Mesenchymal Transition. <i>Journal of Biological Chemistry</i> , 2008, 283, 33437-33446.	3.4	310
12	Functional interactions between phosphatase POPX2 and mDia modulate RhoA pathways. <i>Journal of Cell Science</i> , 2008, 121, 514-521.	2.0	31
13	The p21-Activated Kinase PAK Is Negatively Regulated by POPX1 and POPX2, a Pair of Serine/Threonine Phosphatases of the PP2C Family. <i>Current Biology</i> , 2002, 12, 317-321.	3.9	137