

# ValÃ©rie Lamantia

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4618689/publications.pdf>

Version: 2024-02-01

8  
papers

119  
citations

1306789

7  
h-index

1588620

8  
g-index

8  
all docs

8  
docs citations

8  
times ranked

170  
citing authors

#	ARTICLE	IF	CITATIONS
1	Lower plasma PCSK9 in normocholesterolemic subjects is associated with upregulated adipose tissue surface expression of LDLR and CD36 and NLRP3 inflammasome. <i>Physiological Reports</i> , 2021, 9, e14721.	0.7	15
2	White Adipose Tissue Surface Expression of LDLR and CD36 is Associated with Risk Factors for Type 2 Diabetes in Adults with Obesity. <i>Obesity</i> , 2020, 28, 2357-2367.	1.5	14
3	The Association of Polyunsaturated Fatty Acid $\Delta^5$ -Desaturase Activity with Risk Factors for Type 2 Diabetes Is Dependent on Plasma ApoB-Lipoproteins in Overweight and Obese Adults. <i>Journal of Nutrition</i> , 2019, 149, 57-67.	1.3	9
4	High plasma apolipoprotein B identifies obese subjects who best ameliorate white adipose tissue dysfunction and glucose-induced hyperinsulinemia after a hypocaloric diet. <i>American Journal of Clinical Nutrition</i> , 2018, 108, 62-76.	2.2	13
5	ApoB-lipoproteins and dysfunctional white adipose tissue: Relation to risk factors for type 2 diabetes in humans. <i>Journal of Clinical Lipidology</i> , 2017, 11, 34-45.e2.	0.6	18
6	Nutritional management of hyperapoB. <i>Nutrition Research Reviews</i> , 2016, 29, 202-233.	2.1	22
7	WAT apoC-I secretion: role in delayed chylomicron clearance in vivo and ex vivo in WAT in obese subjects. <i>Journal of Lipid Research</i> , 2016, 57, 1074-1085.	2.0	8
8	The apoB-to-PCSK9 ratio: A new index for metabolic risk in humans. <i>Journal of Clinical Lipidology</i> , 2015, 9, 664-675.	0.6	20