

Kyu Jeung Ahn

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4620904/publications.pdf>

Version: 2024-02-01

58
papers

1,341
citations

394286
19
h-index

360920
35
g-index

59
all docs

59
docs citations

59
times ranked

2310
citing authors

#	ARTICLE	IF	CITATIONS
1	The efficacy and safety of <i>Dendropanax morbifera</i> leaf extract on the metabolic syndrome: a 12-week, placebo controlled, double blind, and randomized controlled trial. <i>Nutrition Research and Practice</i> , 2022, 16, 60.	0.7	1
2	Trends in the Prevalence of Obesity and Its Phenotypes Based on the Korea National Health and Nutrition Examination Survey from 2007 to 2017 in Korea. <i>Diabetes and Metabolism Journal</i> , 2022, 46, 808-812.	1.8	4
3	Pancreatic fat accumulation is associated with decreased β -cell function and deterioration in glucose tolerance in Korean adults. <i>Diabetes/Metabolism Research and Reviews</i> , 2021, 37, e3425.	1.7	9
4	The association between lipoprotein (a) and carotid atherosclerosis in patients with type 2 diabetes without pre-existing cardiovascular disease: A cross-sectional study. <i>Diabetes Research and Clinical Practice</i> , 2021, 171, 108622.	1.1	14
5	Allopurinol ameliorates high fructose diet induced hepatic steatosis in diabetic rats through modulation of lipid metabolism, inflammation, and ER stress pathway. <i>Scientific Reports</i> , 2021, 11, 9894.	1.6	14
6	Fenofibrate, a PPAR α agonist, reduces hepatic fat accumulation through the upregulation of TFEB-mediated lipophagy. <i>Metabolism: Clinical and Experimental</i> , 2021, 120, 154798.	1.5	44
7	Low Skeletal Muscle Mass Is Associated With the Presence, Incidence, and Progression of Coronary Artery Calcification. <i>Canadian Journal of Cardiology</i> , 2021, 37, 1480-1488.	0.8	8
8	Statin use for primary prevention in patients with type 2 diabetes: Can it benefit all ages? â€œ A nationwide propensity-matched cohort study. <i>Diabetes Research and Clinical Practice</i> , 2021, 180, 109044.	1.1	5
9	Increased Visit-to-Visit Liver Enzyme Variability Is Associated with Incident Diabetes: A Community-Based 12-Year Prospective Cohort Study. <i>Diabetes and Metabolism Journal</i> , 2021, 45, 890-898.	1.8	0
10	Clinical Characteristics and Prevalence of Comorbidities according to Metformin Use in Korean Patients with Type 2 Diabetes. <i>International Journal of Endocrinology</i> , 2020, 2020, 1-7.	0.6	5
11	Comparison of the Efficacy and Safety of Rosuvastatin/Ezetimibe Combination Therapy and Rosuvastatin Monotherapy on Lipoprotein in Patients With Type 2 Diabetes: Multicenter Randomized Controlled Study. <i>Diabetes Therapy</i> , 2020, 11, 859-871.	1.2	11
12	Lilly Insulin Glargine Versus Lantus [®] in Type 2 Diabetes Mellitus Patients: India and East Asia Subpopulation Analyses of the ELEMENT 5 Study. <i>Clinical Drug Investigation</i> , 2019, 39, 745-756.	1.1	1
13	Efficacy and Safety of Switching to Tenueligliptin in Patients with Type 2 Diabetes Inadequately Controlled with Dipeptidyl Peptidase-4 Inhibitors: A 12-Week Interim Report. <i>Diabetes Therapy</i> , 2019, 10, 1271-1282.	1.2	8
14	Use of Insulin Glargine 100 U/mL for the Treatment of Type 2 Diabetes Mellitus in East Asians: A Review. <i>Diabetes Therapy</i> , 2019, 10, 805-833.	1.2	4
15	Hospital-Based Korean Diabetes Prevention Study: A Prospective, Multi-Center, Randomized, Open-Label Controlled Study. <i>Diabetes and Metabolism Journal</i> , 2019, 43, 49.	1.8	13
16	Statins for primary prevention in adults aged 75 years and older: A nationwide population-based case-control study. <i>Atherosclerosis</i> , 2019, 283, 28-34.	0.4	19
17	Comparison of the Efficacy of Rosuvastatin Monotherapy 20 mg with Rosuvastatin 5 mg and Ezetimibe 10 mg Combination Therapy on Lipid Parameters in Patients with Type 2 Diabetes Mellitus. <i>Diabetes and Metabolism Journal</i> , 2019, 43, 582.	1.8	14
18	Metformin reduces the risk of cancer in patients with type 2 diabetes. <i>Medicine (United States)</i> , 2018, 97, e0036.	0.4	45

#	ARTICLE	IF	CITATIONS
19	Failure of monotherapy in clinical practice in patients with type 2 diabetes: The Korean National Diabetes Program. <i>Journal of Diabetes Investigation</i> , 2018, 9, 1144-1152.	1.1	10
20	Long-term effects on glycaemic control and β -cell preservation of early intensive treatment in patients with newly diagnosed type 2 diabetes: A multicentre randomized trial. <i>Diabetes, Obesity and Metabolism</i> , 2018, 20, 1121-1130.	2.2	13
21	Comparison of Adherence to Glimepiride/Metformin Sustained Release Once-daily Versus Glimepiride/Metformin Immediate Release BID Fixed-combination Therapy Using the Medication Event Monitoring System in Patients With Type 2 Diabetes. <i>Clinical Therapeutics</i> , 2018, 40, 752-761.e2.	1.1	3
22	Exendin-4, a glucagon-like peptide-1 receptor agonist, reduces hepatic steatosis and endoplasmic reticulum stress by inducing nuclear factor erythroid-derived 2-related factor 2 nuclear translocation. <i>Toxicology and Applied Pharmacology</i> , 2018, 360, 18-29.	1.3	13
23	Efficacy and safety of evogliptin monotherapy in patients with type 2 diabetes and moderately elevated glycated haemoglobin levels after diet and exercise. <i>Diabetes, Obesity and Metabolism</i> , 2017, 19, 1681-1687.	2.2	21
24	Efficacy and safety of gemigliptin, a dipeptidyl peptidase-4 inhibitor, in patients with type 2 diabetes mellitus inadequately controlled with combination treatment of metformin and sulphonylurea: 24-week, multicentre, randomized, double-blind, placebo-controlled study (TROICA study). <i>Diabetes, Obesity and Metabolism</i> , 2017, 19, 635-643.	2.2	11
25	Effects of foot complications in patients with Type 2 diabetes mellitus on public healthcare: An analysis based on the Korea National Diabetes Program Cohort. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 375-380.	1.2	3
26	Differential association between sarcopenia and metabolic phenotype in Korean young and older adults with and without obesity. <i>Obesity</i> , 2017, 25, 244-251.	1.5	21
27	Addition of Ipragliflozin to Metformin Treatment in Korean Patients with Type 2 Diabetes Mellitus: Subgroup Analysis of a Phase 3 Trial. <i>Diabetes and Metabolism Journal</i> , 2017, 41, 135.	1.8	14
28	The pharmacological stimulation of Nurr1 improves cognitive functions via enhancement of adult hippocampal neurogenesis. <i>Stem Cell Research</i> , 2016, 17, 534-543.	0.3	32
29	Effects of C-reactive protein on bone cells. <i>Life Sciences</i> , 2016, 145, 1-8.	2.0	20
30	Aster spathulifolius Maxim extract reduces body weight and fat mass in obese humans. <i>Nutrition Research</i> , 2016, 36, 671-678.	1.3	11
31	Hypoglycemia and Medical Expenses in Patients with Type 2 Diabetes Mellitus: An Analysis Based on the Korea National Diabetes Program Cohort. <i>PLoS ONE</i> , 2016, 11, e0148630.	1.1	18
32	Association between the Circulating Total Osteocalcin Level and the Development of Cardiovascular Disease in Middle-aged Men: A Mean 8.7-year Longitudinal Follow-up Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2015, 22, 136-143.	0.9	25
33	Acromegaly with Normal Insulin-Like Growth Factor-1 Levels and Congestive Heart Failure as the First Clinical Manifestation. <i>Endocrinology and Metabolism</i> , 2015, 30, 395.	1.3	5
34	Clinical and Economic Outcomes in Medication-Adherent and -Nonadherent Patients With Type 2 Diabetes Mellitus in the Republic of Korea. <i>Clinical Therapeutics</i> , 2014, 36, 245-254.	1.1	10
35	Optimal Serum Concentration of 25-Hydroxyvitamin D for Bone Health in Older Korean Adults. <i>Calcified Tissue International</i> , 2013, 92, 68-74.	1.5	29
36	Bisphenol A reduces differentiation and stimulates apoptosis of osteoclasts and osteoblasts. <i>Life Sciences</i> , 2013, 93, 367-372.	2.0	67

#	ARTICLE	IF	CITATIONS
37	Increasing achievement of the target goals for glycemic, blood pressure and lipid control for adults with diagnosed diabetes in Korea. Journal of Diabetes Investigation, 2013, 4, 460-465.	1.1	13
38	Risk Factors for the Progression of Intima-Media Thickness of Carotid Arteries: A 2-Year Follow-Up Study in Patients with Newly Diagnosed Type 2 Diabetes. Diabetes and Metabolism Journal, 2013, 37, 365.	1.8	8
39	Changes in Serum Osteocalcin are Not Associated with Changes in Glucose or Insulin for Osteoporotic Patients Treated with Bisphosphonate. Journal of Bone Metabolism, 2013, 20, 37.	0.5	24
40	Sarcopenia Is Independently Associated with Cardiovascular Disease in Older Korean Adults: The Korea National Health and Nutrition Examination Survey (KNHANES) from 2009. PLoS ONE, 2013, 8, e60119.	1.1	200
41	Circulating Osteocalcin Level Is Not Associated With Incident Type 2 Diabetes in Middle-Aged Male Subjects. Diabetes Care, 2012, 35, 1919-1924.	4.3	44
42	Direct Medical Costs for Patients with Type 2 Diabetes and Related Complications: A Prospective Cohort Study Based on the Korean National Diabetes Program. Journal of Korean Medical Science, 2012, 27, 876.	1.1	41
43	Optimal Range of Triglyceride Values to Estimate Serum Low Density Lipoprotein Cholesterol Concentration in Korean Adults: the Korea National Health and Nutrition Examination Survey, 2009. Journal of Korean Medical Science, 2012, 27, 1530.	1.1	11
44	Utilization patterns and cost of complementary and alternative medicine compared to conventional medicine in patients with type 2 diabetes mellitus. Diabetes Research and Clinical Practice, 2011, 93, 115-122.	1.1	13
45	Autoimmune Hypoglycemia in a Patient with Characterization of Insulin Receptor Autoantibodies. Diabetes and Metabolism Journal, 2011, 35, 80.	1.8	22
46	A Case of Primary Adrenal Insufficiency in a Patient with Acquired Immunodeficiency Syndrome. Endocrinology and Metabolism, 2011, 26, 253.	1.3	1
47	Prevalence of Chronic Complications in Korean Patients with Type 2 Diabetes Mellitus Based on the Korean National Diabetes Program. Diabetes and Metabolism Journal, 2011, 35, 504.	1.8	56
48	Comparison of the Efficacy of Glimepiride, Metformin, and Rosiglitazone Monotherapy in Korean Drug-Naïve Type 2 Diabetic Patients: The Practical Evidence of Antidiabetic Monotherapy Study. Diabetes and Metabolism Journal, 2011, 35, 26.	1.8	32
49	Assessment of glycemic lability and severity of hypoglycemia in Korean patients with type 1 diabetes. Endocrine Journal, 2011, 58, 433-440.	0.7	5
50	A Case of Adrenocortical Carcinoma Secreting Cortisol, Androgen and Aldosterone. Endocrinology and Metabolism, 2011, 26, 239.	1.3	1
51	A Case of Multiple Endocrine Neoplasia Type 1 with Primary Liver Gastrinoma. Journal of Korean Medical Science, 2010, 25, 953.	1.1	7
52	Characteristics of insulin resistance and insulin secretory capacity in Korean subjects with IFG and IGT. Diabetes Research and Clinical Practice, 2010, 89, 250-255.	1.1	22
53	The uncarboxylated form of osteocalcin is associated with improved glucose tolerance and enhanced β -cell function in middle-aged male subjects. Diabetes/Metabolism Research and Reviews, 2009, 25, 768-772.	1.7	204
54	The effects of Acanthopanax senticosus extract on bone turnover and bone mineral density in Korean postmenopausal women. Journal of Bone and Mineral Metabolism, 2009, 27, 584-590.	1.3	22

#	ARTICLE	IF	CITATIONS
55	Neuroprotective Effect of Ghrelin Is Associated with Decreased Expression of Prostate Apoptosis Response-4. Endocrine Journal, 2009, 56, 609-617.	0.7	48
56	Clinical Impacts of Iatrogenic Hypoglycemia. Korean Clinical Diabetes, 2008, 9, 106.	0.1	3
57	Differences in Insulin Sensitivity and Secretory Capacity Based on OGTT in Subjects with Impaired Glucose Regulation. Korean Journal of Internal Medicine, 2007, 22, 270.	0.7	18
58	A Case of Osteomalacia Caused by Severe Vitamin D Deficiency. Journal of Korean Endocrine Society, 2007, 22, 55.	0.1	1