Damien R Lannoy

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4625040/publications.pdf

Version: 2024-02-01

840119 794141 24 372 11 19 citations h-index g-index papers 28 28 28 414 docs citations times ranked citing authors all docs

#	Article	IF	Citations
1	Impact of Multiaccess Infusion Devices on In Vitro Drug Delivery During Multi-Infusion Therapy. Anesthesia and Analgesia, 2009, 109, 1147-1155.	1.1	46
2	Clinical implications of intravenous drug incompatibilities in critically ill patients. Anaesthesia, Critical Care & Dain Medicine, 2019, 38, 173-180.	0.6	39
3	Study of the complexation of risperidone and 9-hydroxyrisperidone with cyclodextrin hosts using affinity capillary electrophoresis and 1H NMR spectroscopy. Journal of Chromatography A, 2008, 1215, 185-193.	1.8	38
4	Cost-Effectiveness Analysis of Five Competing Strategies for the Management of Multiple Recurrent Community-Onset Clostridium difficile Infection in France. PLoS ONE, 2017, 12, e0170258.	1.1	38
5	Infusion Set Characteristics Such as Antireflux Valve and Dead-Space Volume Affect Drug Delivery. Anesthesia and Analgesia, 2010, 111, 1427-1431.	1.1	32
6	The Impact on Drug Mass Flow Rate of Interrupting and Resuming Carrier Fluid Flow. Anesthesia and Analgesia, 2012, 114, 328-332.	1.1	32
7	TRAnexamic acid in hemorrhagic CESarean section (TRACES) randomized placebo controlled dose-ranging pharmacobiological ancillary trial: study protocol for a randomized controlled trial. Trials, 2018, 19, 149.	0.7	16
8	Configurational stability of 9-hydroxyrisperidone. Kinetics and mechanism of racemization. Tetrahedron: Asymmetry, 2009, 20, 1125-1131.	1.8	15
9	Evaluation of the stability of vancomycin solutions at concentrations used in clinical services. European Journal of Hospital Pharmacy, 2020, 27, e87-e92.	0.5	15
10	Criteria for choosing an intravenous infusion line intended for multidrug infusion in anaesthesia and intensive care units. Anaesthesia, Critical Care & Dain Medicine, 2017, 36, 53-63.	0.6	13
11	Therapeutic and pharmaco-biological, dose-ranging multicentre trial to determine the optimal dose of TRAnexamic acid to reduce blood loss in haemorrhagic CESarean delivery (TRACES): study protocol for a randomised, double-blind, placebo-controlled trial. Trials, 2018, 19, 148.	0.7	13
12	In Vitro Assessment of Interaction Between Amino Acids and Copper in Neonatal Parenteral Nutrition. Journal of Parenteral and Enteral Nutrition, 2016, 40, 827-834.	1.3	10
13	Stability of 10 mg/mL cefuroxime solution for intracameral injection in commonly used polypropylene syringes and new ready-to-use cyclic olefin copolymer sterile vials using the LC-UV stability-indicating method. Drug Development and Industrial Pharmacy, 2016, 42, 166-174.	0.9	10
14	Economic assessment of aseptic compounding rooms in hospital pharmacies in five European countries. Journal of Oncology Pharmacy Practice, 2015, 21, 102-110.	0.5	8
15	Stability of frozen 1% voriconazole eye-drops in both glass and innovative containers. European Journal of Pharmaceutical Sciences, 2020, 141, 105102.	1.9	8
16	Intraventricular dopamine infusion alleviates motor symptoms in a primate model of Parkinson's disease. Neurobiology of Disease, 2020, 139, 104846.	2.1	8
17	Immunoassay quantification of human insulin added to ternary parenteral nutrition containers: comparison of two methods. Analytical and Bioanalytical Chemistry, 2017, 409, 3677-3684.	1.9	7
18	Administration des médicaments oraux chez le patient bénéficiant d'une nutrition entérale. Nutritic Clinique Et Metabolisme, 2013, 27, 255-262.	on 0.2	3

#	Article	IF	CITATION
19	Quality control and stability of ketamine, remifentanil, and sufentanil syringes in a pediatric operating theater. Paediatric Anaesthesia, 2019, 29, 193-199.	0.6	3
20	Instability of Insulin Aspart Diluted in Dextrose. Diabetes Care, 2020, 43, e77-e78.	4.3	3
21	Cost-effectiveness of taurolidine locks to prevent recurrent catheter-related blood stream infections in adult patients receiving home parenteral nutrition: a 2-year mirror-image study. Clinical Nutrition, 2021, 40, 4309-4315.	2.3	3
22	Long-term stability of ready-to-use norepinephrine solution at 0.2 and 0.5 mg/mL. European Journal of Hospital Pharmacy, 2020, 27, e93-e98.	0.5	2
23	Addition of Regular Insulin to Ternary Parenteral Nutrition: A Stability Study. Pharmaceutics, 2021, 13, 458.	2.0	2
24	Behavior of Regular Insulin in a Parenteral Nutrition Admixture: Validation of an LC/MS-MS Assay and the In Vitro Evaluation of Insulin Glycation. Pharmaceutics, 2022, 14, 1081.	2.0	2