
Maimana A Magdy

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4636031/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Quantitative determination of Dapoxetine Hydrochloride and Tadalafil using different validated spectrophotometric methods. Spectrochimica Acta - Part A: Molecular and Biomolecular Spectroscopy, 2020, 226, 117611.	2.0	25
2	Simultaneous HPTLC and RP-HPLC methods for determination of bumadizone in the presence of its alkaline-induced degradation product. Biomedical Chromatography, 2012, 26, 1143-1149.	0.8	20
3	A Validated Green HPTLC Method for Quantitative Determination of Dapoxetine Hydrochloride and Tadalafil in Bulk and Pharmaceutical Formulations. Journal of Chromatographic Science, 2020, 58, 303-308.	0.7	15
4	A green approach for simultaneous analysis of two natural hepatoprotective drugs in pure forms, capsules and human plasma using HPLC-UV method. Microchemical Journal, 2019, 151, 104258.	2.3	14
5	Kinetic study and mechanism of Niclosamide degradation. Spectrochimica Acta - Part A: Molecular and Biomolecular Spectroscopy, 2014, 132, 655-662.	2.0	11
6	Different chromatographic methods for determination of alogliptin benzoate, metformin hydrochloride, and metformin impurity in bulk and pharmaceutical dosage form. Journal of Separation Science, 2021, 44, 833-842.	1.3	9
7	Green validated HPTLC and HPLC methods for determination of ephedrine hydrochloride and naphazoline nitrate in the presence of methylparaben, in their pure forms and pharmaceutical formulation. Journal of Planar Chromatography - Modern TLC, 2020, 33, 141-148.	0.6	7
8	Validated ecofriendly chromatographic method for quantitative determination of antiâ€migraine quaternary mixture. Journal of Separation Science, 2020, 43, 2330-2337.	1.3	7
9	Linear Support Vector Regression and Partial Least-Squares for Determination of Dapoxetine Hydrochloride and Tadalafil in Binary Pharmaceutical Mixtures. Journal of AOAC INTERNATIONAL, 2020, 103, 132-139.	0.7	6
10	Quantitative Determination of Anti-Migraine Quaternary Mixture in Presence of <i>p</i> -Aminophenol and 4-Chloroacetanilide. Journal of Chromatographic Science, 2022, 60, 538-544.	0.7	5
11	US FDA–validated TLC method with four greenness assessment evaluations for simultaneous determination of prednisolone and esomeprazole in spiked human plasma. Biomedical Chromatography, 2022, 36, e5343.	0.8	5
12	Stability indicating spectrophotometric methods for determination of bumadizone in the presence of its alkaline degradation product. Drug Invention Today (discontinued), 2013, 5, 139-147.	0.6	4
13	A validated HPTLC method for the quantitative determination of duloxetine hydrochloride and 1-naphthol in bulk and pharmaceutical formulation. Journal of Planar Chromatography - Modern TLC, 2020, 33, 391-396.	0.6	4
14	Multivariate Model Update Chemometric Methods for Determination of Prednisolone and Esomeprazole in Spiked Human Plasma: A Comparative Study. Journal of AOAC INTERNATIONAL, 2022, 105, 317-322.	0.7	3
15	Three Spectrophotometric Methods for Quantitative Analysis of Duloxetine in Presence of its Toxic Impurity: 1-Naphthol. Journal of AOAC INTERNATIONAL, 2020, 103, 972-979.	0.7	1
16	Resolving a Binary Mixture of Hepatoprotective Drugs Using Different Validated Spectrophotometric Methods. Journal of Analytical Chemistry, 2020, 75, 176-183.	0.4	1
17	A Comparative Chemometric Study for Quantitative Determination of Duloxetine Hydrochloride in the Presence of its Toxic Impurity 1-Naphthol. Current Pharmaceutical Analysis, 2020, 16, 1030-1036.	0.3	1
18	HPTLC Separation of a Hepatoprotective Combination in Pharmaceutical Formulation and Human Plasma. Journal of Chromatographic Science, 2020, 58, 411-417.	0.7	0